


Graduate Texts in Mathematics

**G. Takeuti
W. M. Zaring**

Introduction to Axiomatic Set Theory




**Springer-Verlag
New York Heidelberg Berlin**

Graduate Texts in Mathematics 1

Managing Editor: P. R. Halmos

G. Takeuti W.M. Zaring

**Introduction to
Axiomatic Set Theory**



Springer-Verlag New York Heidelberg Berlin 

Gaisi Takeuti

Professor of Mathematics, University of Illinois

Wilson M. Zaring

Associate Professor of Mathematics, University of Illinois

AMS Subject Classifications (1970): 02K15, 04-01, 02K05

ISBN 0-387-05302-6 Springer-Verlag New York Heidelberg Berlin (soft cover)

ISBN 0-387-90024-1 Springer-Verlag New York Heidelberg Berlin (hard cover)

ISBN 3-540-05302-6 Springer-Verlag Berlin Heidelberg New York (soft cover)

This work is subject to copyright. All rights are reserved, whether the whole or part of the material is concerned, specifically those of translation, reprinting, re-use of illustrations, broadcasting, reproduction by photocopying machine or similar means, and storage in data banks. Under § 54 of the German Copyright Law where copies are made for other than private use, a fee is payable to the publisher, the amount of the fee to be determined by agreement with the publisher. © by Springer-Verlag New York 1971. Library of Congress Catalog Card Number 70-142387. Printed in U.S.A.

Preface

In 1963, the first author introduced a course in set theory at the University of Illinois whose main objectives were to cover Gödel's work on the consistency of the axiom of choice (AC) and the generalized continuum hypothesis (GCH), and Cohen's work on the independence of the AC and the GCH. Notes taken in 1963 by the second author were taught by him in 1966, revised extensively, and are presented here as an introduction to axiomatic set theory.

Texts in set theory frequently develop the subject rapidly moving from key result to key result and suppressing many details. Advocates of the fast development claim at least two advantages. First, key results are highlighted, and second, the student who wishes to master the subject is compelled to develop the details on his own. However, an instructor using a "fast development" text must devote much class time to assisting his students in their efforts to bridge gaps in the text.

We have chosen instead a development that is quite detailed and complete. For our slow development we claim the following advantages. The text is one from which a student can learn with little supervision and instruction. This enables the instructor to use class time for the presentation of alternative developments and supplementary material. Indeed, by presenting the student with a suitably detailed development, we enable him to move more rapidly to the research frontier and concentrate his efforts on original problems rather than expending that effort redoing results that are well-known.

Our main objective in this text is to acquaint the reader with Zermelo-Fraenkel set theory and bring him to a study of interesting results in one semester. Among the results that we consider interesting are the following: Sierpinski's proof that the GCH implies the AC, Rubin's proof that the aleph hypothesis (AH) implies the AC, Gödel's consistency results and Cohen's forcing techniques. We end the text with a section on Cohen's proof of the independence of the axiom of constructibility.

In a sequel to this text entitled *Axiomatic Set Theory*, we will discuss, in a very general framework, relative constructibility, general forcing, and their relationship.

We are indebted to so many people for assistance in the preparation of this text that we would not attempt to list them all. We do, however, wish to express our appreciation to Professors Kenneth Appel, W. W. Boone, Carl Jockusch, Thomas McLaughlin, and Nobuo Zama for their valuable suggestions and advice. We also wish to thank Professor H. L. Africk, Professor Kenneth Bowen, Paul E. Cohen, Eric Frankl, Charles Kahane, Donald Pelletier, George Sacerdote, Eric Schindler and Kenneth Slonneger, all students or former students of the authors, for their assistance at various stages in the preparation of the manuscript.

A special note of appreciation goes to Prof. Hisao Tanaka who made numerous suggestions for improving the text and to Dr. Klaus Gloede who through the cooperation of Springer-Verlag, provided us with valuable editorial advice and assistance.

We are also grateful to Mrs. Carolyn Bloemker for her care and patience in typing the final manuscript.

Urbana, January 1971

Gaisi Takeuti
Wilson M. Zaring

Contents

1	Introduction	1
2	Language and Logic.	4
3	Equality	6
4	Classes	9
5	The Elementary Properties of Classes	14
6	Functions and Relations	21
7	Ordinal Numbers	32
8	Ordinal Arithmetic	49
9	Relational Closure and the Rank Function	63
10	Cardinal Numbers	71
11	The Axiom of Choice, the Generalized Continuum Hypothesis and Cardinal Arithmetic	87
12	Models	102
13	Absoluteness	112
14	The Fundamental Operations.	133
15	The Gödel Model	143
16	The Arithmetization of Model Theory	175
17	Cohen's Method	196
18	Forcing	203
19	Languages, Structures, and Models	236
	Bibliography	239
	Problem List	241
	Appendix	243
	Index	245
	Index of Symbols	249

1 Introduction

In 1895 and 1897 Georg Cantor (1845–1918) published his master works on ordinal and cardinal numbers¹. Cantor's theory of ordinal and cardinal numbers was the culmination of three decades of research on number "aggregates". Beginning with his paper on the denumerability of infinite sets², published in 1874, Cantor had built a new theory of the infinite. In this theory a collection of objects, even an infinite collection, is conceived of as a single entity.

The notion of an infinite set as a complete entity was not universally accepted. Critics argued that logic is an extrapolation from experience that is necessarily finitistic. To extend the logic of the finite to the infinite entailed risks too grave to countenance. This prediction of logical disaster seemed vindicated when at the turn of the century paradoxes were discovered in the very foundations of the new discipline. Dedekind stopped publication of his *Was sind und was sollen die Zahlen?* Frege conceded that the foundation of his *Grundgesetze der Arithmetik* was destroyed.

Nevertheless set theory gained sufficient support to survive the crisis of the paradoxes. In 1908, speaking at the International Congress at Rome, the great Henri Poincaré (1854–1912) urged that a remedy be sought³. As a reward he promised "the joy of the physician called to treat a beautiful pathologic case." By that time Zermelo and Russell were already at work seeking fundamental principles on which a consistent theory could be built.

From this one might assume that the sole purpose for axiomatizing is to avoid the paradoxes. There are however reasons to believe that axiomatic set theory would have evolved even in the absence of paradoxes.

¹ *Beiträge zur Begründung der transfiniten Mengenlehre*. (Erster Artikel.) Math. Ann. Vol. 46, 1895, p. 481—512. (Zweiter Artikel) Math. Ann. Vol. 49, 1897, p. 207—246. For an English translation see Cantor, Georg. *Contributions to the Founding of the Theory of Transfinite Numbers*. Dover Publications, Inc.

² *Über eine Eigenschaft des Inbegriffes aller reellen algebraischen Zahlen*. J. Reine Angew. Math. Vol. 77, 1874, p. 258—262. In this paper Cantor proves that the set of all algebraic numbers is denumerable and that the set of all real numbers is not denumerable.

³ *Atti del IV Congresso Internazionale dei Matematici Roma 1909*, Vol. 1, p. 182.

Certainly the work of Dedekind and of Frege in the foundations of arithmetic was not motivated by fear of paradoxes but rather by a desire to see what foundational principles were required. In his *Begriffsschrift* Frege states:

“..., we divide all truths that require justification into two kinds, those for which the proof can be carried out purely by means of logic and those for which it must be supported by facts of experience. ... Now, when I came to consider the question to which of these two kinds the judgement of arithmetic belong, I first had to ascertain how far one could proceed in arithmetic by means of inferences alone, ...”⁴.

Very early in the history of set theory it was discovered that the Axiom of Choice, the Continuum Hypothesis, and the Generalized Continuum Hypothesis are principles of special interest and importance. The Continuum Hypothesis is Cantor’s conjectured solution to the question of how many points there are on a line in Euclidean space⁵. A formal statement of the Continuum Hypothesis and its generalization will be given later.

The Axiom of Choice, in one formulation, asserts that given any collection of pairwise disjoint non-empty sets, there exists a set that has exactly one element in common with each set of the given collection. The discovery that the Axiom of Choice has important implications for all major areas of mathematics provided compelling reasons for its acceptance. Its status as an axiom, and also that of the Generalized Continuum Hypothesis, was however not clarified until Kurt Gödel in 1938, proved them to be consistent with the axioms of general set theory and Paul Cohen, in 1963, proved that they are each independent of the axioms of general set theory. Our major objective in this text will be a study of the contributions of Gödel and Cohen. In order to do this we must first develop a satisfactory theory of sets.

For Cantor a set was “any collection into a whole M of definite and separate objects m of our intuition or our thought⁶.” This naive acceptance of any collection as a set leads us into the classical paradoxes, as for example Russell’s Paradox: If the collection of all sets that are not elements of themselves is a set then this set has the property that it is an element of itself if and only if it is not an element of itself.

In developing a theory of sets we then have two alternatives. Either we must abandon the idea that our theory is to encompass arbitrary

⁴ van Heijenoort, Jean. *From Frege to Gödel*. Cambridge: Harvard University Press, 1967. p. 5.

⁵ See *What is Cantor’s Continuum Problem?* by Kurt Gödel in the *Amer. Math. Monthly*. Vol. 54 (1947), p. 515—525. A revised and expanded version of this paper is also found in Benacerraf, Paul and Putnam, Hilary. *Philosophy of Mathematics Selected Readings*. Englewood Cliffs, Prentice-Hall, Inc. 1964.

⁶ Cantor, Georg. *Contributions to the Founding of the Theory of Transfinite Numbers*. New York: Dover Publications, Inc.

collections in the sense of Cantor, or we must distinguish between at least two types of collections, arbitrary collections that we call classes and certain special collections that we call sets. Classes, or arbitrary collections, are however so useful and our intuitive feelings about classes are so strong that we dare not abandon them. A satisfactory theory of sets must provide a means of speaking safely about classes. There are several ways of developing such a theory.

Bertrand Russell (1872–1970) and Alfred North Whitehead (1861—1947) in their *Principia Mathematica* (1910) resolved the difficulties with a theory of types. They established a hierarchy of types of collections. A collection x can be a member of a collection y only if y is one level higher in the hierarchy than x . In this system there are variables for each type level in the hierarchy and hence there are infinitely many primitive notions.

Two other systems, Gödel-Bernays (GB) set theory and Zermelo-Fraenkel (ZF) set theory, evolved from the work of Bernays (1937–1954), Fraenkel (1922), Gödel (1940), von Neumann (1925–1929), Skolem (1922), and Zermelo (1908). Our listing is alphabetical. We will not attempt to identify the specific contribution of each man. Following each name we have indicated the year or period of years of major contribution.

In Gödel-Bernays set theory the classical paradoxes are avoided by recognizing two types of classes, sets and proper classes. Sets are classes that are permitted to be members of other classes. Proper classes have sets as elements but are not themselves permitted to be elements of other classes. In this system we have three primitive notions; set, class and membership. In the formal language we have set variables, class variables and a binary predicate symbol “ \in ”.

In Zermelo-Fraenkel set theory we have only two primitive notions; set and membership. Class is introduced as a defined term. In the formal language we have only set variables and a binary predicate symbol “ \in ”. Thus in ZF quantification is permitted only on set variables while in GB quantification is permitted on both set and class variables. As a result there are theorems in GB that are not theorems in ZF. It can however be proved that GB is a conservative extension of ZF in the sense that every well formed formula of ZF is provable in ZF if and only if it is provable in GB.

Gödel’s⁷ work was done in Gödel-Bernays set theory. We, however, prefer Zermelo-Fraenkel theory in which Cohen⁸ worked.

⁷ Gödel, Kurt: *The Consistency of the Continuum Hypothesis*. Princeton: Princeton University Press, 1940.

⁸ Cohen, Paul J.: *The Independence of the Continuum Hypothesis*. Proceedings of the National Academy of Science of the United States of America, Vol. 50, 1963, pp. 1143—1148.

2 Language and Logic

The language of our theory consists of

individual variables: x_0, x_1, \dots ,

a predicate constant: \in .

logical symbols: $\neg, \vee, \wedge, \rightarrow, \leftrightarrow, \forall, \exists$,

auxiliary symbols: $(), []$.

The logical symbols in the order listed are for negation, disjunction, conjunction, implication, equivalence, universal quantification, and existential quantification.

We will not restrict ourselves to a minimal list of logical symbols, nor will we in general distinguish between primitive and defined logical symbols. When, in a given context, it is convenient to have a list of primitive symbols we will assume whatever list best suits our immediate need.

We will use

$$a, b, c, \dots, x, y, z$$

as variables in the meta-language whose domain is the collection of individual variables of the formal language. When we need many such variables we will use x_0, x_1, \dots and rely upon the context to make clear whether for example x_0 is a particular variable of the formal language or a meta-variable ranging over all individual variables of the formal language.

We will also use

$$\varphi, \psi, \eta$$

as meta-variables that range over well formed formulas (wffs).

Our rules for well formed formulas are the following:

- 1) If x and y are individual variables then $x \in y$ is a wff.
- 2) If φ and ψ are wffs then $\neg\varphi, \varphi \vee \psi, \varphi \wedge \psi, \varphi \rightarrow \psi$ and $\varphi \leftrightarrow \psi$ are wffs.
- 3) If φ is a wff and x is an individual variable then $(\forall x)\varphi$ and $(\exists x)\varphi$ are wffs.

A formula is well formed if and only if its being so is deducible from rules 1)–3). This means that every wff can be decomposed in a finite

number of steps into prime wffs, i.e., wffs that cannot be further decomposed into well formed parts; as for example $x_2 \in x_{10}$.

It is easily proved that there exists an effective procedure for determining whether or not a given formula is well formed. Consequently there is an effective procedure for determining the number of well formed parts contained in a given formula, i.e., the number of proper subformulas that are well formed. This fact will be of value to us in the proof of certain results in later sections.

From the language just described we obtain Zermelo-Fraenkel set theory by adjoining logical axioms, rules of inference, and nonlogical axioms. The nonlogical axioms for ZF will be introduced in context and collected on page 196. The logical axioms and rules of inference for our theory are the following.

Logical Axioms:

- 1) $\varphi \rightarrow [\psi \rightarrow \varphi]$.
- 2) $[\varphi \rightarrow [\psi \rightarrow \eta]] \rightarrow [[\varphi \rightarrow \psi] \rightarrow [\varphi \rightarrow \eta]]$.
- 3) $[\neg \varphi \rightarrow \neg \psi] \rightarrow [\psi \rightarrow \varphi]$.
- 4) $(\forall x) [\varphi \rightarrow \psi] \rightarrow [\varphi \rightarrow (\forall x)\psi]$ where x is not free in φ .
- 5) $(\forall x) \varphi(x) \rightarrow \varphi(a)$ where x has no free occurrence in a well formed part of φ of the form $(\forall a)\psi$.

Rules of Inference:

- 1) From φ and $\varphi \rightarrow \psi$ to infer ψ .
- 2) From φ to infer $(\forall x)\varphi$.

We will assume without proof those results from logic that we need except one theorem. We will use the conventional symbol “ \vdash ” to indicate that a wff is a theorem in our theory. That is, “ $\vdash \varphi$ ” is the meta-statement that the wff φ is deducible by the rules of inference from the logical axioms and the nonlogical axioms yet to be stated. To indicate that φ is deducible using only the logical axioms we will write “ $\vdash_{LA} \varphi$ ”. We say that two wffs φ and ψ are logically equivalent if and only if $\vdash_{LA} \varphi \leftrightarrow \psi$.

3 Equality

Definition 3.1. $a = b \stackrel{\Delta}{\longleftrightarrow} (\forall x) [x \in a \longleftrightarrow x \in b]$.

Remark. Definition 3.1 is incomplete in that the variable x does not occur in $a = b$, consequently we have not made clear which of the infinitely many formulas, equivalent under alphabetic change of variable, we intend $a = b$ to be an abbreviation for. This problem is easily resolved by specifying that x is the first variable on our list

$$x_0, x_1, \dots$$

that is distinct from a and from b . Having thereby shown that we can specify a particular formula we will not bother to do so either here or in similar definitions to follow.

Our intuitive idea of equality is of course identity. A basic property that we expect of equality is that paraphrased as “equals may be substituted for equals,” that is, if $a = b$ then anything that can be asserted of a can also be asserted of b . In particular if a certain wff holds for a it must also hold for b .

If φ_x is a wff in which the variable x occurs unbound zero or more times then the assertion that φ_x holds for a does not mean simply that the formula obtained from φ_x by replacing each free occurrence of x by a holds. A problem can arise if a occurs bound in φ_x . For example, in ZF

$$(\forall a) [a = x]$$

is false for each x in spite of the fact that

$$(\forall a) [a = a]$$

is true.

By “ φ_x holds for a ” we mean that the following formula holds. We first replace each bound occurrence of a in φ_x by a variable that does not occur in φ_x . We then replace each free occurrence of x by a . The resulting formula we will denote by $\varphi_x(a)$ or simply $\varphi(a)$.

Our substitution property for equality can then be stated as

$$a = b \rightarrow [\varphi(a) \longleftrightarrow \varphi(b)].$$

We, however, need not postulate such a substitution principle for, as we will now show, it can be deduced from Definition 3.1 and the following weaker principle.

Axiom 1 (Axiom of Extensionality).

$$(\forall a) (\forall x) (\forall y) [x = y \wedge x \in a \rightarrow y \in a].$$

Theorem 3.2.

- 1) $a = a.$
- 2) $a = b \rightarrow b = a.$
- 3) $a = b \wedge b = c \rightarrow a = c.$

Proof.

- 1) $(\forall x) [x \in a \leftrightarrow x \in a].$
- 2) $(\forall x) [x \in a \leftrightarrow x \in b] \rightarrow (\forall x) [x \in b \leftrightarrow x \in a].$
- 3) $(\forall x) [x \in a \leftrightarrow x \in b] \wedge (\forall x) [x \in b \leftrightarrow x \in c] \rightarrow (\forall x) [x \in a \leftrightarrow x \in c].$

Theorem 3.3. $x = y \rightarrow [x \in a \leftrightarrow y \in a].$

Proof. Axiom 1 and Theorem 3.2, 2).

Theorem 3.4. $a = b \rightarrow [\varphi(a) \leftrightarrow \varphi(b)].$

Proof. (By induction on n the number of well formed parts of φ .)
If $n = 0$ then $\varphi(a)$ is of the form $c \in d$, $c \in a$, $a \in c$, or $a \in a$. Clearly

$$a = b \rightarrow [c \in d \leftrightarrow c \in d].$$

From the definition of equality

$$a = b \rightarrow [c \in a \leftrightarrow c \in b].$$

From Theorem 3.3

$$a = b \rightarrow [a \in c \leftrightarrow b \in c].$$

Again from the definition of equality and Theorem 3.3 respectively

$$a = b \rightarrow [a \in a \leftrightarrow a \in b],$$

$$a = b \rightarrow [a \in b \leftrightarrow b \in b].$$

Therefore

$$a = b \rightarrow [a \in a \leftrightarrow b \in b].$$

As our induction hypothesis we assume the result true for each wff having fewer than n well formed parts. If $n > 0$ and $\varphi(a)$ has exactly n

well formed parts then $\varphi(a)$ must be of the form

$$1) \neg\psi(a), \quad 2) \psi(a) \wedge \eta(a), \quad \text{or} \quad 3) (\forall x)\psi(a).$$

In each case $\psi(a)$ and $\eta(a)$ have fewer than n well formed parts and hence from the induction hypothesis

$$a = b \rightarrow [\psi(a) \leftrightarrow \psi(b)],$$

$$a = b \rightarrow [\eta(a) \leftrightarrow \eta(b)].$$

From properties of negation and conjunction it then follows that

$$a = b \rightarrow [\neg\psi(a) \leftrightarrow \neg\psi(b)]$$

$$a = b \rightarrow [\psi(a) \wedge \eta(a) \leftrightarrow \psi(b) \wedge \eta(b)].$$

Thus if $\varphi(a)$ is $\neg\psi(a)$ or $\psi(a) \wedge \eta(a)$,

$$a = b \rightarrow [\varphi(a) \leftrightarrow \varphi(b)].$$

If $\varphi(a)$ is $(\forall x)\psi(a)$ we first choose an x that is distinct from a and b . By generalization on the induction hypothesis we then obtain

$$a = b \rightarrow [(\forall x)\psi(a) \leftrightarrow (\forall x)\psi(b)].$$

Extensionality assures us that a set is completely determined by its elements. From a casual acquaintance with this axiom one might assume that Extensionality is a substitution principle having more to do with logic than set theory. This suggests that if equality were taken as a primitive notion then perhaps this axiom could be dispensed with. Dana Scott¹ however, has proved that this cannot be done without weakening the system. Thus, even if we were to take equality as a primitive logical notion it would still be necessary to add an extensionality axiom².

¹ *Essays on the Foundations of Mathematics*. Amsterdam: North-Holland Publishing Company, 1962, pp. 115—131.

² See Quine, Van Orman: *Set Theory and its Logic*. Cambridge: Mass. Harvard University Press 1969, 30f.

4 Classes

We pointed out in the introduction that one objective of axiomatic set theory is to avoid the classical paradoxes. One such paradox, the Russell paradox, arose from the naive acceptance of the idea that given any property there exists a set whose elements are those objects having the given property, i.e., given a wff $\varphi(x)$ containing no free variables other than x , there exists a set that contains all objects for which $\varphi(x)$ holds and contains no object for which $\varphi(x)$ does not hold. More formally

$$(\exists a) (\forall x) [x \in a \longleftrightarrow \varphi(x)].$$

This principle, called the Axiom of Abstraction, was accepted by Frege in his *Grundgesetze der Arithmetik* (1893). In a letter¹ to Frege (1902) Bertrand Russell pointed out that the principle leads to the following paradox.

Consider the predicate $x \notin x$. If there exists a set a such that

$$(\forall x) [x \in a \longleftrightarrow x \notin x]$$

then in particular

$$a \in a \longleftrightarrow a \notin a.$$

The idea of the collection of all objects having a specified property is so basic that we could hardly abandon it. But if it is to be retained how shall the paradox be resolved? The Zermelo-Fraenkel approach is the following.

For each wff $\varphi(x, a_1, \dots, a_n)$ we will introduce a class symbol

$$\{x \mid \varphi(x, a_1, \dots, a_n)\}$$

which is read “the class of all x such that $\varphi(x, a_1, \dots, a_n)$.” Our principal interpretation is that the class symbol $\{x \mid \varphi(x)\}$ denotes the class of individuals x that have the property $\varphi(x)$. We will show that class is an extension of the notion of set in that every set is a class but not every class is a set.

¹ van Heijenoort, Jean: *From Frege to Gödel*. Cambridge: Harvard University Press 1967, pp. 124—125.

We will extend the \in -relation to class symbols in such a way that an object is an element of a class $\{x|\varphi(x)\}$ if and only if that object is a set and it has the defining property for the class. The Russell paradox is then resolved by showing that $\{x|x \notin x\}$ is a proper class i.e., a class that is not a set. It is then disqualified for membership in any class, including itself, on the grounds that it is not a set.

Were we to adjoin the symbols

$$\{x|\varphi(x)\}$$

to our object language it would be necessary to extend our rules for wffs and add axioms governing the new symbols. We choose instead to introduce classes as defined terms. It is of course essential that we provide an effective procedure for reducing to primitive symbols any formula that contains a defined term. We begin by defining the contexts in which class symbols are permitted to appear. Our only concern will be their appearance in wffs in the *wider sense* as defined by the following rules.

Definition 4.1.

- 1) If a and b are individual variables then $a \in b$ is a wff in the wider sense.
- 2) If φ and ψ are wffs in the wider sense and a and b are individual variables then $a \in \{x|\psi(x)\}$, $\{x|\varphi(x)\} \in b$, and $\{x|\varphi(x)\} \in \{x|\psi(x)\}$ are wffs in the wider sense.
- 3) If φ and ψ are wffs in the wider sense then $\neg\varphi$, $\varphi \wedge \psi$, $\varphi \vee \psi$, $\varphi \rightarrow \psi$, and $\varphi \longleftrightarrow \psi$ are wffs in the wider sense.
- 4) If φ is a wff in the wider sense and x is an individual variable then $(\exists x)\varphi$ and $(\forall x)\varphi$ are wffs in the wider sense.

A formula is a wff in the wider sense iff its being so is deducible from 1)–4).

It is our intention that every wff in the wider sense be an abbreviation for a wff in the original sense. It is also our intention that a set belong to a class iff it has the defining property of that class i.e.

$$a \in \{x|\varphi(x)\} \quad \text{iff} \quad \varphi(a).$$

Definition 4.2. If φ and ψ are wffs in the wider sense then

- 1) $a \in \{x|\varphi(x)\} \stackrel{\Delta}{\longleftrightarrow} \varphi(a)$.
- 2) $\{x|\varphi(x)\} \in a \stackrel{\Delta}{\longleftrightarrow} (\exists y \in a) (\forall z) [z \in y \longleftrightarrow \varphi(z)]$.
- 3) $\{x|\varphi(x)\} \in \{x|\psi(x)\} \stackrel{\Delta}{\longleftrightarrow} (\exists y \in \{x|\psi(x)\}) (\forall z) [z \in y \longleftrightarrow \varphi(z)]$.

Remark. From Definition 4.2 it is easily proved that each wff in the wider sense is reducible to a wff φ^* that is determined uniquely by the following rules.

Definition 4.3 If φ and ψ are wffs in the wider sense then

- 1) $[a \in b]^* \xleftrightarrow{\Delta} a \in b.$
- 2) $[a \in \{x | \varphi(x)\}]^* \xleftrightarrow{\Delta} [\varphi(a)]^* \xleftrightarrow{\Delta} \varphi^*(a).$
- 3) $[\{x | \varphi(x)\} \in a]^* \xleftrightarrow{\Delta} (\exists y \in a) (\forall z) [z \in y \longleftrightarrow \varphi^*(z)].$
- 4) $[\{x | \varphi(x)\} \in \{x | \psi(x)\}]^* \xleftrightarrow{\Delta} (\exists y) [\psi^*(y) \wedge (\forall z) [z \in y \longleftrightarrow \varphi^*(z)]].$
- 5) $[\neg \varphi]^* \xleftrightarrow{\Delta} \neg \varphi^*.$
- 6) $[\varphi \wedge \psi]^* \xleftrightarrow{\Delta} \varphi^* \wedge \psi^*.$
- 7) $[(\forall x) \varphi]^* \xleftrightarrow{\Delta} (\forall x) \varphi^*.$

Theorem 4.4. *Each wff in the wider sense φ , is reducible to one and only one wff φ^* determined from φ by the rules 1)–7) of Definition 4.3.*

Proof. (By induction on n the number of well formed parts, in the wider sense, in φ .) If $n = 0$, i.e., if φ has no well formed parts in the wider sense, then φ must be of the form $a \in b$. By 1) of Definition 4.3, φ^* is $a \in b$.

As our induction hypothesis we assume that each wff in the wider sense having fewer than n well formed parts, in the wider sense is reducible to one and only one wff that is determined by the rules 1)–7) of Definition 4.3. If φ is a wff in the wider sense having exactly n well formed parts in the wider sense and if $n > 0$ then φ must be of one of the following forms:

- 1) $a \in \{x | \psi(x)\},$
- 2) $\{x | \psi(x)\} \in a,$
- 3) $\{x | \psi(x)\} \in \{x | \eta(x)\},$
- 4) $\neg \psi,$
- 5) $\psi \wedge \eta,$
- 6) $(\forall x) \psi.$

In each case ψ and η have fewer than n well formed parts in the wider sense and hence there are unique wffs ψ^* and η^* determined by ψ and η respectively and the rules 1)–7) of Definition 4.3. Then by rules 2)–7) φ determines a unique wff φ^* .

Remark. From Theorem 4.4 every wff in the wider sense φ is an abbreviation for a wff φ^* . The proof tacitly assumes the existence of an effective procedure for determining whether or not a given formula is a wff in the wider sense. That such a procedure exists we leave as an

exercise for the reader. From such a procedure it is immediate that there is an effective procedure for determining φ^* from φ .

Theorem 4.4 also assures us that in Definitions 4.1 and 4.2 we have not extended the notion of class but have only extended the notation for classes for if $\varphi(x)$ is a wff in the wider sense then

$$\{x | \varphi(x)\} .$$

and

$$\{x | \varphi^*(x)\}$$

are the same class. This is immediate from Theorem 4.4 and equality for classes which we now define. We wish this definition to encompass not only equality between class and class but also between set and class. For this, and other purposes, we introduce the notion of a *term*.

By a term we mean an individual variable or a class symbol. We shall use capital Roman letters

$$A, B, C, \dots$$

as meta-variables on terms.

Definition 4.5 If A and B are terms then

$$A = B \stackrel{\Delta}{\longleftrightarrow} (\forall x) [x \in A \longleftrightarrow x \in B] .$$

Theorem 4.6. $A \in B \longleftrightarrow (\exists x) [x = A \wedge x \in B]$.

Proof. Definitions 4.2 and 4.5.

Theorem 4.7. *If $A, B,$ and C are terms then*

- 1) $A = A,$
- 2) $A = B \rightarrow B = A,$
- 3) $A = B \wedge B = C \rightarrow A = C.$

The proof is similar to that of Theorem 3.2 and is left to the reader.

Theorem 4.8. *If A and B are terms and φ is a wff in the wider sense, then*

$$A = B \rightarrow [\varphi(A) \longleftrightarrow \varphi(B)] .$$

The proof is by induction on the number of well formed parts, in the wider sense, in φ . It is similar to the proof of Theorem 3.4 and is left to the reader.

Theorem 4.9. $a = \{x | x \in a\}.$

Proof. $x \in a \longleftrightarrow x \in a.$

Remark. Theorem 4.9 establishes that every set is a class. We now wish to establish that not all classes are sets. We introduce the predicates $\mathcal{M}(A)$ and $\mathcal{P}_r(A)$ for “ A is a set” and “ A is a proper class” respectively.

Definition 4.10. $\mathcal{M}(A) \stackrel{\Delta}{\longleftrightarrow} (\exists x) [x = A]$.

$$\mathcal{P}_r(A) \stackrel{\Delta}{\longleftrightarrow} \neg \mathcal{M}(A).$$

Theorem 4.11. $(\forall a) \mathcal{M}(a)$.

Proof. $(\forall a) [a = a]$

Theorem 4.12. $A \in \{x | \varphi(x)\} \longleftrightarrow \mathcal{M}(A) \wedge \varphi(A)$.

Proof. Definitions 4.2 and 4.10 and Theorems 4.6 and 4.8.

Definition 4.13. $\text{Ru} = \{x | x \notin x\}$.

Theorem 4.14. $\mathcal{P}_r(\text{Ru})$.

Proof. From Theorem 4.12

$$\mathcal{M}(\text{Ru}) \rightarrow [\text{Ru} \in \text{Ru} \longleftrightarrow \text{Ru} \notin \text{Ru}].$$

Therefore Ru is a proper class.

Remark. Since the Russell class, Ru, is a proper class the Russell paradox is resolved. It should be noted that the Russell class is the first non-set we have encountered. Others will appear in the sequel.

We now have examples establishing that the class of individuals for which a given wff φ holds may be a set or a proper class. Those sets, $\{x | \varphi(x)\}$, for which $\varphi(x)$ has no free variables other than x we call *definable sets*.

Definition 4.15. A set a is *definable* iff there is a wff $\varphi(x)$ containing no free variables other than x such that $a = \{x | \varphi(x)\}$.

5 The Elementary Properties of Classes

In this section we will introduce certain properties of classes with which the reader is already familiar. The immediate consequences of the definitions are for the most part elementary and easily proved; consequently they will be left to the reader as exercises.

We begin with the notion of unordered pair, $\{a, b\}$, and ordered pair $\langle a, b \rangle$.

Definition 5.1. $\{a, b\} \triangleq \{x \mid x = a \vee x = b\}$.
 $\{a\} \triangleq \{a, a\}$.

Remark. We postulate that pairs are sets.

Axiom 2. (*Axiom of Pairing*). $(\forall a)(\forall b) \mathcal{M}(\{a, b\})$.

Definition 5.2. $\langle a, b \rangle \triangleq \{x \mid x = \{a\} \vee x = \{a, b\}\}$.

Exercises. *Prove the following.*

- 1) $x \in \{a, b\} \longleftrightarrow x = a \vee x = b$.
- 2) $x \in \{a\} \longleftrightarrow x = a$.
- 3) $x \in \langle a, b \rangle \longleftrightarrow x = \{a\} \vee x = \{a, b\}$.
- 4) $\{a\} = \{b\} \longleftrightarrow a = b$.
- 5) $\{a\} = \{b, c\} \longleftrightarrow a = b = c$.
- 6) $\langle a, b \rangle = \langle c, d \rangle \longleftrightarrow a = c \wedge b = d$.
- 7) $\mathcal{M}(\langle a, b \rangle)$.
- 8) $(\forall x) [a \in x \rightarrow b \in x] \rightarrow a = b$.
- 9) $A \in B \rightarrow \mathcal{M}(A)$.

Remark. The notion of unordered pair and ordered pair have natural generalizations to unordered n-tuple, $\{a_1, a_2, \dots, a_n\}$ and ordered n-tuple, $\langle a_1, a_2, \dots, a_n \rangle$.

Definition 5.3. $\{a_1, a_2, \dots, a_n\} = \{x \mid x = a_1 \vee x = a_2 \vee \dots \vee x = a_n\}$.

Definition 5.4. $\langle a_1, a_2, \dots, a_n \rangle = \langle \langle a_1, \dots, a_{n-1} \rangle, a_n \rangle, n \geq 3$.

Remark. Since ordered pairs are sets it follows by induction that ordered n-tuples are also sets. From the fact that unordered pairs are sets we might also hope to prove by induction that unordered n-tuples are sets. For such a proof however we need certain properties of set union.

Definition 5.5. $\cup(A) \triangleq \{x \mid (\exists y) [x \in y \wedge y \in A]\}$.

Axiom 3 (*Axiom of Unions*). $(\forall a) \mathcal{M}(\cup(a))$.

Definition 5.6. $A \cup B \triangleq \{x \mid x \in A \vee x \in B\}$.

$A \cap B \triangleq \{x \mid x \in A \wedge x \in B\}$.

Theorem 5.7. $a \cup b = \cup\{a, b\}$.

Proof. $x \in a \cup b \iff [x \in a \vee x \in b]$
 $\iff (\exists y) [x \in y \wedge y \in \{a, b\}]$
 $\iff x \in \cup\{a, b\}$.

Corollary 5.8. $\mathcal{M}(a \cup b)$.

Proof. Theorem 5.7, the Axiom of Unions, and the Axiom of Pairing.

Exercises. *Prove the following.*

- 1) $b \in \{a_1, a_2, \dots, a_n\} \iff b = a_1 \vee b = a_2 \vee \dots \vee b = a_n$.
- 2) $\{a_1, a_2, \dots, a_{n+1}\} = \{a_1, a_2, \dots, a_n\} \cup \{a_{n+1}\}, n \geq 1$.
- 3) $\mathcal{M}(\{a_1, a_2, \dots, a_n\})$.
- 4) $\mathcal{M}(\langle a_1, a_2, \dots, a_n \rangle)$.
- 5) $a \in b \cup \{b\} \iff a \in b \vee a = b$.
- 6) $A \cup B = B \cup A$.
- 7) $A \cap B = B \cap A$.
- 8) $(A \cup B) \cup C = A \cup (B \cup C)$.
- 9) $(A \cap B) \cap C = A \cap (B \cap C)$.
- 10) $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$.
- 11) $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$.

Remark. We next introduce the notions of subclass $A \subseteq B$, and power set, $\mathcal{P}(a)$.

Definition 5.9. $A \subseteq B \stackrel{\Delta}{\iff} (\forall x) [x \in A \rightarrow x \in B]$.

$A \subset B \stackrel{\Delta}{\iff} A \subseteq B \wedge A \neq B$.

Definition 5.10. $\mathcal{P}(a) \triangleq \{x \mid x \subseteq a\}$.

Axiom 4 (*Axiom of Powers*). $(\forall a) \mathcal{M}(\mathcal{P}(a))$.

Exercises. Reduce the following wffs in the wider sense to wffs.

- 1) $\{a, b\} \in \cup(c),$
- 2) $\mathcal{M}(\{a, b\}),$
- 3) $\mathcal{M}(\cup(a)),$
- 4) $\mathcal{M}(\mathcal{P}(a)).$

Prove the following.

- 5) $A \subseteq B \wedge B \subseteq C \rightarrow A \subseteq C.$
- 6) $A \subseteq B \rightarrow C \cap A \subseteq C \cap B.$
- 7) $A \subseteq B \rightarrow C \cup A \subseteq C \cup B.$
- 8) $A \subseteq B \leftrightarrow B = (A \cup B).$
- 9) $A \subseteq B \leftrightarrow A = (A \cap B).$
- 10) $A \subseteq A.$
- 11) $A \subseteq A \cup B.$
- 12) $A \cap B \subseteq A \wedge A \cap B \subseteq B.$
- 13) $A \not\subseteq B \leftrightarrow (\exists x) [x \in A \wedge x \notin B].$
- 14) $A \subset B \rightarrow [\cup(A) \subseteq \cup(B)].$
- 15) $A = B \rightarrow [\cup(A) = \cup(B)].$
- 16) $a \in \mathcal{P}(a).$
- 17) $\cup(\mathcal{P}(a)) = a.$
- 18) $a \subset b \leftrightarrow \mathcal{P}(a) \subset \mathcal{P}(b).$
- 19) $a = b \leftrightarrow \mathcal{P}(a) = \mathcal{P}(b).$

Remark. Since there exist classes that are not sets we must reject the Axiom of Abstraction. Zermelo proposed as its replacement the Axiom of Separation that asserts that the class of all objects in a set that have a given property is a set i.e.

$$\mathcal{M}(a \cap A).$$

Fraenkel in turn chose to replace Zermelo's Axiom of Separation by a principle that asserts that functions map sets onto sets. The condition that a wff $\varphi(u, v)$ should define a function i.e., that

$$\{\langle u, v \rangle \mid \varphi(u, v)\}$$

should be a single valued relation is simply that

$$\varphi(u, v) \wedge \varphi(u, w) \rightarrow v = w.$$

If this is the case and if

$$A = \{u | (\exists v) \varphi(u, v)\} \quad \text{and} \quad B = \{v | (\exists u) \varphi(u, v)\}$$

then the function in question maps A onto B and by Fraenkel's Axiom maps $a \cap A$ onto a subset of B . That is

$$(\forall a) \mathcal{M}(\{y | (\exists z \in a) \varphi(z, y)\})$$

or equivalently

$$(\forall a) (\exists b) (\forall y) [y \in b \leftrightarrow (\exists z \in a) \varphi(z, y)].$$




Fig. 1.

From Fraenkel's Axiom we can easily deduce Zermelo's. The two are however not equivalent. Indeed Richard Montague has proved that ZF is not a finite extension of Zermelo set theory¹.

Axiom 5 (*Axiom Schema of Replacement*).

$$(\forall a) [(\forall u) (\forall v) (\forall w) [\varphi(u, v) \wedge \varphi(u, w) \rightarrow v = w] \\ \rightarrow (\exists b) (\forall y) [y \in b \leftrightarrow (\exists x) [x \in a \wedge \varphi(x, y)]]].$$

Theorem 5.11. (*Zermelo's Schema of Separation*).

$$(\forall a) (\exists b) (\forall x) [x \in b \leftrightarrow x \in a \wedge \varphi(x)].$$

Proof. Applying Axiom 5 to the wff $\varphi(u) \wedge u = v$ where v does not occur in $\varphi(u)$ we have that

$$[\varphi(u) \wedge u = v] \wedge [\varphi(u) \wedge u = w] \rightarrow v = w.$$

Therefore

$$(\exists b) (\forall x) [x \in b \leftrightarrow (\exists u) [\varphi(u) \wedge u = x \wedge x \in a]]$$

i.e.

$$(\exists b) (\forall x) [x \in b \leftrightarrow \varphi(x) \wedge x \in a].$$

¹ *Essays on the Foundations of Mathematics*. Amsterdam, North-Holland Publishing Company.

Corollary 5.12. $(\forall a) \mathcal{M}(\{x | x \in a \wedge \varphi(x)\})$.

Proof. By Theorem 5.11 $(\exists b)[b = \{x | x \in a \wedge \varphi(x)\}]$.

Remark. Hereafter we will write $\{x \in a | \varphi(x)\}$ for $\{x | x \in a \wedge \varphi(x)\}$.

Corollary 5.13. $\mathcal{M}(a \cap A)$.

Proof. $a \cap A = \{x \in a | x \in A\}$.

Definition 5.14. $A - B = \{x | x \in A \wedge x \notin B\}$.

Theorem 5.15. $\mathcal{M}(a - A)$.

Proof. $a - A = \{x \in a | x \notin A\}$.

Definition 5.16. $0 \triangleq \{x | x \neq x\}$.

Theorem 5.17. $(\forall a) [a - a = 0]$.

Proof. $x \in a - a \iff x \in a \wedge x \notin a$
 $\iff x \neq x$
 $\iff x \in 0$.

Corollary 5.18. $\mathcal{M}(0)$.

Proof. $(\forall a) [a - a = 0] \rightarrow (\exists a) [a - a = 0]$. From Theorems 5.17 and 5.15 we then conclude that 0 is a set.

Theorem 5.19.

- 1) $(\forall a) [a \notin 0]$.
- 2) $a \neq 0 \iff (\exists x) [x \in a]$.

Proof.

- 1) $(\forall a) [a = a]$. Therefore $a \notin 0$.
- 2) $a \neq 0 \iff (\exists x) [x \in 0 \wedge x \notin a] \vee (\exists x) [x \in a \wedge x \notin 0]$.

Since $(\forall x) [x \notin 0]$ we conclude that $a \neq 0 \iff (\exists x) [x \in a]$.

Remark. To exclude the possibility that a set can be an element of itself and also to exclude the possibility of having “ \in -loops” i.e., $a_1 \in a_2 \in \dots \in a_n \in a_1$, Zermelo introduced his Axiom of Regularity, also known as the Axiom of Foundation, which asserts that every non-empty set a contains an element x with the property that no element of x is also an element of a . A stronger form of this axiom asserts the same property of non-empty classes. Later we will prove that the weak and strong forms are in fact equivalent.

Axiom 6 (*Axiom of Regularity, weak form*).

$$a \neq 0 \rightarrow (\exists x) [x \in a \wedge x \cap a = 0].$$

Axiom 6' (*Axiom of Regularity, strong form*).

$$A \neq 0 \rightarrow (\exists x) [x \in A \wedge x \cap A = 0].$$

Theorem 5.20. $\neg [a_1 \in a_2 \in \dots \in a_n \in a_1]$.

Proof. $(\exists a) [a = \{a_1, a_2, \dots, a_n\}]$. Therefore if $a_1 \in a_2 \in \dots \in a_n \in a_1$ then $(\forall x) [x \in a \rightarrow x \cap a \neq 0]$. This contradicts Regularity.

Corollary 5.21. $(\forall a) [a \notin a]$.

Proof. Theorem 5.20 with $n = 1$.

Definition 5.22. $V = \{x \mid x = x\}$.

Theorem 5.23. $\mathcal{P}_r(V)$.

Proof. Since $V = V$ it follows that if $\mathcal{M}(V)$ then $V \in V$.

Remark. From the strong form of Regularity we can deduce the following induction principle.

Theorem 5.24. $(\forall a) [a \subseteq A \rightarrow a \in A] \rightarrow A = V$.

Proof. If $B = V - A$ and if $B \neq 0$ then by (strong) Regularity $(\exists x) [x \in B \wedge x \cap B = 0]$ i.e., $(\forall y) [y \in x \rightarrow y \notin B]$. But $y \notin B \rightarrow y \in A$. Therefore $x \subseteq A$ and hence $x \in A$. But this contradicts the fact that $x \in B$. Therefore $B = 0$ and $A = V$. (See Exer. 1) below.)

Remark. Theorem 5.24 assures us that if every set a has a certain property, $\varphi(a)$, whenever each element of a has that property then every set does indeed have the property. Consider the following example. If each element of a set a has no infinite descending \in -chain then clearly a has no infinite descending \in -chain. Therefore there are no infinite descending \in -chains.

Exercises. *Prove the following.*

- 1) $0 \subseteq A \wedge A \subseteq V$.
- 2) $(\forall x) [x \notin A] \rightarrow A = 0$.
- 3) $A \subseteq a \rightarrow \mathcal{M}(A)$.
- 4) $\mathcal{M}(A) \rightarrow \mathcal{M}(A \cap B)$.
- 5) $Ru = V$.
- 6) $A \notin A$.
- 7) $A - B = 0 \leftrightarrow A \subseteq B$.

- 8) $A - B = A \cap \{x \mid x \notin B\}$.
- 9) $A - (B \cup C) = (A - B) \cap (A - C)$.
- 10) $A - (B \cap C) = (A - B) \cup (A - C)$.
- 11) $A - (B - A) = A$.
- 12) $A \cap (B - C) = (A \cap B) - (A \cap C) = (A \cap B) - C$.
- 13) $A \cup (B - C) = (A \cup B) - (C - A) = (A \cup B) - ((A \cap B) - C)$.
- 14) $A - B \subseteq A$.
- 15) $A \subseteq B \rightarrow [C - B \subseteq C - A]$.

6 Functions and Relations

Definition 6.1. $A \times B \triangleq \{x | (\exists a) (\exists b) [a \in A \wedge b \in B \wedge x = \langle a, b \rangle]\}$.

Theorem 6.2. $\mathcal{M}(a \times b)$.

Proof.

$$\begin{aligned} c \in a \times b &\iff (\exists x) (\exists y) [x \in a \wedge y \in b \wedge c = \langle x, y \rangle]. \\ &\rightarrow (\exists x) (\exists y) [\{x\} \subseteq a \cup b \wedge \{x, y\} \subseteq a \cup b \wedge c = \langle x, y \rangle]. \\ &\rightarrow (\exists x) (\exists y) [\{x\} \in \mathcal{P}(a \cup b) \wedge \{x, y\} \in \mathcal{P}(a \cup b) \wedge c = \langle x, y \rangle]. \\ &\rightarrow (\exists x) (\exists y) [\langle x, y \rangle \in \mathcal{P}(\mathcal{P}(a \cup b)) \wedge c = \langle x, y \rangle]. \\ &\rightarrow c \in \mathcal{P}(\mathcal{P}(a \cup b)). \end{aligned}$$

Therefore $a \times b \subseteq \mathcal{P}(\mathcal{P}(a \cup b))$; hence $\mathcal{M}(a \times b)$.

Definition 6.3.

- 1) $A^2 \triangleq A \times A$.
- 2) $A^{-1} \triangleq \{\langle x, y \rangle | \langle y, x \rangle \in A\}$.

Remark. If A contains elements that are not ordered pairs, for example, if $A = \{\langle 0, 1 \rangle, 0\}$ then $(A^{-1})^{-1} \neq A$; indeed for the example at hand $A^{-1} = \{\langle 1, 0 \rangle\}$ and $(A^{-1})^{-1} = \{\langle 0, 1 \rangle\}$.

Definition 6.4.

- 1) $\mathcal{R}el(A) \xleftrightarrow{\Delta} A \subseteq V^2$.
- 2) $\mathcal{U}n(A) \xleftrightarrow{\Delta} (\forall u) (\forall v) (\forall w) [\langle u, v \rangle \in A \wedge \langle u, w \rangle \in A \rightarrow v = w]$.
- 3) $\mathcal{U}n_2(A) \xleftrightarrow{\Delta} \mathcal{U}n(A) \wedge \mathcal{U}n(A^{-1})$.
- 4) $\mathcal{F}nc(A) \xleftrightarrow{\Delta} \mathcal{R}el(A) \wedge \mathcal{U}n(A)$.
- 5) $\mathcal{F}nc_2(A) \xleftrightarrow{\Delta} \mathcal{R}el(A) \wedge \mathcal{U}n_2(A)$.

Definition 6.5.

- 1) $\mathcal{D}(A) \triangleq \{x | (\exists y) [\langle x, y \rangle \in A]\}$.
- 2) $\mathcal{W}(A) \triangleq \{y | (\exists x) [\langle x, y \rangle \in A]\}$.

Definition 6.6.

- 1) $A \upharpoonright B \triangleq A \cap (B \times V).$
- 2) $A''B \triangleq \mathcal{W}(A \upharpoonright B).$
- 3) $A \circ B = \{\langle x, y \rangle \mid (\exists z) [\langle x, z \rangle \in B \wedge \langle z, y \rangle \in A]\}.$

Remark. The intended interpretation of the predicates in Definition 6.4 and the terms in Definitions 6.5 and 6.6 are the following:

$\mathcal{R}el(A)$ means “ A is a relation.”

$\mathcal{U}n(A)$ means “ A is single valued.”

$\mathcal{U}n_2(A)$ means “ A is one-to-one.”

$\mathcal{F}nc(A)$ means “ A is a function.”

$\mathcal{F}nc_2(A)$ means “ A is a one-to-one function.”

$\mathcal{D}(A)$ denotes the domain of A .

$\mathcal{W}(A)$ denotes the range of A .

$A \upharpoonright B$ denotes the restriction of A to B .

$A''B$ denotes the image of B under A .

$A \circ B$ denotes the composite of A with B .

It should be noted that a class does not have to be a relation in order to have a domain and a range. Indeed every class has both. The domain of A is simply the class of first entries of those ordered pairs that are in A and the range of A is the class of second entries of those ordered pairs that are in A . In addition every class has certain function-like properties: $A \upharpoonright B$ is the class of ordered pairs in A having first entry in B and $A''B$ is the class of second entries of those ordered pairs in A that have first entry in B .

Exercises. *Prove the following.*

- 1) $(A^{-1})^{-1} \subseteq A.$
- 2) $(A^{-1})^{-1} = A \iff \mathcal{R}el(A).$
- 3) $A \times B \subseteq V^2.$
- 4) $V^2 \subset V.$
- 5) $\mathcal{R}el(A) \wedge \mathcal{R}el(B) \rightarrow \mathcal{R}el(A \cup B).$
- 6) $(\forall x) [x \in A \rightarrow \mathcal{R}el(x)] \rightarrow \mathcal{R}el(\cup(A)).$
- 7) $\mathcal{U}n_2(A) \rightarrow (\forall w) (\forall x) (\forall y) (\forall z) [\langle w, x \rangle \in A \wedge \langle z, y \rangle \in A \rightarrow [w = z \iff x = y]].$
- 8) $\mathcal{F}nc_2(A) \rightarrow \mathcal{F}nc(A) \wedge \mathcal{F}nc(A^{-1}).$
- 9) $(A \times B) \cap (C \times D) = (A \cap C) \times (B \cap D).$
- 10) $A \subseteq B \rightarrow \mathcal{D}(A) \subseteq \mathcal{D}(B).$

- 11) $A \subseteq B \rightarrow \mathcal{W}(A) \subseteq \mathcal{W}(B).$
- 12) $\mathcal{R}el(A \Gamma B).$
- 13) $\langle x, y \rangle \in A \Gamma B \leftrightarrow \langle x, y \rangle \in A \wedge x \in B.$
- 14) $\mathcal{D}(A \Gamma B) = B \cap \mathcal{D}(A).$
- 15) $A \Gamma B \subseteq A.$
- 16) $[A = A \Gamma \mathcal{D}(A)] \leftrightarrow \mathcal{R}el(A).$
- 17) $B \subseteq C \rightarrow [(A \Gamma C) \Gamma B = A \Gamma B].$
- 18) $\mathcal{U}n(A) \rightarrow \mathcal{U}n(A \Gamma B).$
- 19) $b \in A \text{“} B \leftrightarrow (\exists a) [\langle a, b \rangle \in A \wedge a \in B].$
- 20) $[A \subseteq B \wedge C \subseteq D] \rightarrow A \text{“} C \subseteq B \text{“} D.$
- 21) If $A = \{\langle \langle x, y \rangle, \langle y, x \rangle \rangle \mid x \in V \cap y \in V\}$ then
 $\mathcal{U}n(A) \wedge A \text{“} B = B^{-1}.$
- 22) If $A = \{\langle \langle x, y \rangle, x \rangle \mid x \in V \wedge y \in V\}$ then $\mathcal{U}n(A) \wedge A \text{“} B = \mathcal{D}(B).$
- 23) If $A = \{\langle \langle x, y \rangle, y \rangle \mid x \in V \wedge y \in V\}$ then $\mathcal{U}n(A) \wedge A \text{“} B = \mathcal{W}(B).$
- 24) $\mathcal{R}el(A \circ B).$
- 25) $\mathcal{U}n(A) \wedge \mathcal{U}n(B) \rightarrow \mathcal{U}n(A \circ B).$
- 26) $\mathcal{U}n(A) \wedge \mathcal{U}n(B) \rightarrow \mathcal{F}nc(A \circ B).$
- 27) $\mathcal{U}n_2(A) \wedge \mathcal{U}n_2(B) \rightarrow \mathcal{U}n_2(A \circ B).$
- 28) $\mathcal{U}n_2(A) \wedge \mathcal{U}n_2(B) \rightarrow \mathcal{F}nc_2(A \circ B).$
- 29) $\mathcal{F}nc(A) \wedge \mathcal{F}nc(B) \rightarrow \mathcal{F}nc(A \circ B).$
- 30) $\mathcal{F}nc_2(A) \wedge \mathcal{F}nc_2(B) \rightarrow \mathcal{F}nc_2(A \circ B).$
- 31) $A^{-1} = (A \cap V^2)^{-1}.$

Theorem 6.7. $\mathcal{U}n(A) \rightarrow \mathcal{M}(A \text{“} a).$

Proof. $\mathcal{U}n(A) \leftrightarrow (\forall u) (\forall v) (\forall w) [\langle u, v \rangle \in A \wedge \langle u, w \rangle \in A \rightarrow v = w].$

From the Axiom Schema of Replacement it then follows that

$$(\exists b) (\forall x) [x \in b \leftrightarrow (\exists y) [\langle y, x \rangle \in A \wedge y \in a]]$$

i.e.,

$$(\exists b) (\forall x) [x \in b \leftrightarrow x \in A \text{“} a].$$

Thus $(\exists b) [b = A \text{“} a].$

Remark. Theorem 6.7 assures us that single valued relations *i.e.*, functions, map sets onto sets.

Corollary 6.8.

- 1) $\mathcal{M}(a^{-1})$.
- 2) $\mathcal{M}(\mathcal{D}(a))$.
- 3) $\mathcal{M}(\mathcal{W}(a))$.

Proof.

- 1) If $A = \{\langle \langle x, y \rangle, \langle y, x \rangle \rangle \mid x \in V \wedge y \in V\}$ then $\mathcal{U}_n(A) \wedge A^{\ast}a = a^{-1}$.
- 2) If $A = \{\langle \langle x, y \rangle, x \rangle \mid x \in V \wedge y \in V\}$ then $\mathcal{U}_n(A) \wedge A^{\ast}a = \mathcal{D}(a)$.
- 3) If $A = \{\langle \langle x, y \rangle, y \rangle \mid x \in V \wedge y \in V\}$ then $\mathcal{U}_n(A) \wedge A^{\ast}a = \mathcal{W}(a)$.

Corollary 6.9.

- 1) $\mathcal{M}(A \times B) \longleftrightarrow \mathcal{M}(B \times A)$.
- 2) $\mathcal{P}_r(A) \wedge B \neq 0 \rightarrow \mathcal{P}_r(A \times B) \wedge \mathcal{P}_r(B \times A)$.

Proof.

1) If $C = \{\langle \langle x, y \rangle, \langle y, x \rangle \rangle \mid x \in V \wedge y \in V\}$ then $\mathcal{U}_n(C) \wedge C^{\ast}(A \times B) = (B \times A) \wedge C^{\ast}(B \times A) = (A \times B)$.

2) If $B \neq 0$ then $(\exists b) [b \in B]$. If $C = \{\langle \langle x, b \rangle, x \rangle \mid x \in A\}$ then $\mathcal{U}_n(C) \wedge \mathcal{D}(C) \subseteq A \times B$.

Therefore $\mathcal{M}(A \times B) \rightarrow \mathcal{M}(\mathcal{D}(C))$. But $A = C^{\ast}\mathcal{D}(C)$ and hence $\mathcal{M}(\mathcal{D}(C)) \rightarrow \mathcal{M}(A)$. From this contradiction we conclude that $A \times B$ is a proper class and hence by 1) so is $B \times A$.

Definition 6.10.

$$(\exists! x) \varphi(x) \xleftrightarrow{\Delta} (\exists x) \varphi(x) \wedge (\forall x) (\forall y) [\varphi(x) \wedge \varphi(y) \rightarrow x = y].$$

Definition 6.11. $A^{\ast}b = \{x \mid (\exists y) [x \in y \wedge \langle b, y \rangle \in A] (\exists! y) [\langle b, y \rangle \in A]\}$.

Theorem 6.12.

- 1) $\langle b, y \rangle \in A \wedge (\exists! y) [\langle b, y \rangle \in A] \rightarrow A^{\ast}b = y$.
- 2) $\neg(\exists! y) [\langle b, y \rangle \in A] \rightarrow A^{\ast}b = 0$.

Proof.

1) From Definition 6.11, $\langle b, y \rangle \in A \wedge (\exists! y) [\langle b, y \rangle \in A]$ implies

$$x \in A^{\ast}b \longleftrightarrow x \in y$$

i.e., $A^{\ast}b = y$.

2) From Definition 6.11, $\neg(\exists! y) [\langle b, y \rangle \in A]$ implies

$$(\forall x) [x \notin A^{\ast}b]$$

i.e., $A^{\ast}b = 0$.

Remark. From Theorem 6.12 we see that Definition 6.11 is an extension of the notion of function value. If A is a function and if b is in $\mathcal{D}(A)$ then $A'b$ is the value of A at b in the usual sense. If b is not in $\mathcal{D}(A)$ then $A'b = 0$. If A is not a function $A'b$ is still defined. Indeed if b is not in $\mathcal{D}(A)$ then $A'b = 0$. If b is in $\mathcal{D}(A)$ but there are two different ordered pairs in A with first entry b then again $A'b = 0$. If b is in $\mathcal{D}(A)$ and $\langle b, y \rangle$ is the only ordered pair in A with first entry b then $A'b = y$.

Corollary 6.13. $\mathcal{M}(A'b)$.

Proof. Theorem 6.12 assures us that $(\exists y) [A'b = y]$.

Definition 6.14.

- 1) $A \mathcal{F}n B \xleftrightarrow{\Delta} \mathcal{F}nc(A) \wedge \mathcal{D}(A) = B.$
- 2) $A \mathcal{F}n_2 B \xleftrightarrow{\Delta} \mathcal{F}nc_2(A) \wedge \mathcal{D}(A) = B.$
- 3) $F : A \longrightarrow B \xleftrightarrow{\Delta} F \mathcal{F}n A \wedge \mathcal{W}(F) \subseteq B.$
- 4) $F : A \xrightarrow{\text{onto}} B \xleftrightarrow{\Delta} F \mathcal{F}n A \wedge \mathcal{W}(F) = B.$
- 5) $F : A \xrightarrow{1-1} B \xleftrightarrow{\Delta} F \mathcal{F}n_2 A \wedge \mathcal{W}(F) \subseteq B.$
- 6) $F : A \xrightarrow[1-1]{\text{onto}} B \xleftrightarrow{\Delta} F \mathcal{F}n_2 A \wedge \mathcal{W}(F) = B.$

Theorem 6.15.

- 1) $A \mathcal{F}n a \rightarrow \mathcal{M}(A).$
- 2) $A \mathcal{F}n_2 a \rightarrow \mathcal{M}(A).$

Proof. 1) $A \mathcal{F}n a \rightarrow A \subseteq (a \times A''a) \wedge \mathcal{U}n(A).$ But $\mathcal{U}n(A) \rightarrow \mathcal{M}(A''a).$ Therefore $\mathcal{M}(a \times A''a)$ and hence $\mathcal{M}(A).$

- 2) $A \mathcal{F}n_2 a \rightarrow A \mathcal{F}n a.$

Theorem 6.16. $\mathcal{U}n(A) \rightarrow \mathcal{M}(A \Gamma a).$

Proof. $\mathcal{U}n(A) \rightarrow \mathcal{U}n(A \Gamma a).$ But we also have $\mathcal{R}el(A \Gamma a) \wedge \mathcal{D}(A \Gamma a) \subseteq a.$ Therefore $(A \Gamma a) \mathcal{F}n \mathcal{D}(A \Gamma a) \wedge \mathcal{M}(\mathcal{D}(A \Gamma a)).$ Hence $\mathcal{M}(A \Gamma a).$

Exercises. Prove the following.

- 1) $\mathcal{P}r(A) \rightarrow \mathcal{P}r(A^2).$
- 2) $\mathcal{P}r(V^2).$
- 3) $\mathcal{U}n(B) \wedge (\exists a) [A - \mathcal{W}(B \Gamma a) = 0] \rightarrow \mathcal{M}(A).$
- 4) $A \mathcal{F}n C \wedge B \mathcal{F}n D \rightarrow [A = B \leftrightarrow [C = D \wedge (\forall x)[x \in C \rightarrow A'x = B'x]]].$
- 5) $\mathcal{U}n(A) \wedge \mathcal{U}n(B) \wedge a \in \mathcal{D}(A \circ B) \rightarrow (A \circ B)'a = A'B'a.$
- 6) $A_1 \mathcal{F}n B_1 \wedge A_2 \mathcal{F}n B_2 \wedge \mathcal{W}(A_2) \subseteq B_1 \rightarrow A_1 \circ A_2 \mathcal{F}n B_2.$

- 7) $A_1 \mathcal{F}n_2 B_1 \wedge A_2 \mathcal{F}n_2 B_2 \wedge \mathcal{W}(A_2) = B_1 \rightarrow A_1 \circ A_2 \mathcal{F}n_2 B_2$
 $\wedge \mathcal{W}(A_1 \circ A_2) = \mathcal{W}(A_1).$
- 8) $A \mathcal{F}n_2 B \wedge \mathcal{W}(A) = C \rightarrow A^{-1} \mathcal{F}n_2 C \wedge \mathcal{W}(A^{-1}) = B.$
- 9) $Un(A) \wedge \mathcal{M}(B) \rightarrow \mathcal{M}(A \ulcorner B).$
- 10) $Un(A) \rightarrow A \ulcorner B = \{A \ulcorner b \mid b \in B \cap \mathcal{D}(A)\}.$
- 11) $A \mathcal{F}n B \vee A \mathcal{F}n_2 B \rightarrow A = A \Gamma B.$
- 12) $(\exists x) (\exists y) [x \neq y \wedge \langle b, x \rangle \in A \wedge \langle b, y \rangle \in A] \rightarrow A \ulcorner b = 0.$
- 13) $\neg(\exists x) [\langle b, x \rangle \in A] \rightarrow A \ulcorner b = 0.$

Remark. In later sections we will study structures consisting of a class A on which is defined a relation R i.e. $R \subseteq A^2$. Since for any class B , $B \cap A^2 \subseteq A^2$ we see that every class B determines a relation on A in a very natural way. We therefore choose to begin our discussion with a very general theory of ordered pairs of classes (A, R) that we will call relational systems.

Definition 6.17. $x R y \stackrel{\Delta}{\longleftrightarrow} \langle x, y \rangle \in R.$

Remark. In the material ahead we will be interested in several types of relational systems. In the next section for example we will study a special class of objects called ordinal numbers. On this class there is an important order relation. Later we will be interested in partial ordering of certain sets.

Definition 6.18.

- 1) $R \text{ Or } A \stackrel{\Delta}{\longleftrightarrow} (\forall x, y \in A) [x R y \longleftrightarrow \neg [x = y \vee y R x]]$
 $\wedge (\forall x, y, z \in A) [x R y \wedge y R z \rightarrow x R z].$
- 2) $R \text{ Po } A \stackrel{\Delta}{\longleftrightarrow} (\forall x \in A) [x R x] \wedge (\forall x, y \in A) [x R y \wedge y R x \rightarrow x = y]$
 $\wedge (\forall x, y, z \in A) [x R y \wedge y R z \rightarrow x R z].$

Remark. There are several properties of relational systems whose proofs depend upon the classes of R -predecessors:

$$\{x \mid x R y\}$$

Theorem 6.19. $(R^{-1}) \ulcorner \{y\} = \{x \mid x R y\}.$

Proof. $x R y \longleftrightarrow \langle x, y \rangle \in R$
 $\longleftrightarrow \langle y, x \rangle \in R^{-1}$
 $\longleftrightarrow x \in (R^{-1}) \ulcorner \{y\}.$

Definition 6.20. $R^{-1} A \triangleq (R^{-1}) \ulcorner A.$

Remark. From Theorem 6.19 we see that $A \cap R^{-1}\{y\} = 0$ means that no element of A precedes y in the sense of R . If, in addition, $y \in A$ then y is an R -minimal element of A . We wish to consider relations with respect to which each subclass of a given class has an R -minimal element. Such a relation we call a *foundational* relation. Since we cannot quantify on class symbols we must formulate our definition in terms of subsets and impose additional conditions that will enable us to deduce the property for subclasses. Later we will show that these additional conditions are not essential.

Definition 6.21.

$$R \text{ Fr } A \stackrel{\Delta}{\longleftrightarrow} (\forall a) [a \subseteq A \wedge a \neq 0 \rightarrow (\exists x \in a) [a \cap R^{-1}\{x\} = 0]].$$

Definition 6.22. $E = \{\langle a, b \rangle \mid a \in b\}$.

Remark. From the Axiom of Regularity we see that the \in -relation E is a foundational relation on V . As in the case of the \in -relation, foundational relations have no relational loops and, as we will prove later, no infinite descending relational chains.

Theorem 6.23.

$$R \text{ Fr } A \wedge a_1 \in A \wedge \cdots \wedge a_n \in A \rightarrow \neg [a_1 R a_2 \wedge a_2 R a_3 \wedge \cdots \wedge a_n R a_1].$$

The proof is left to the reader.

Remark. There are two types of foundational relations that are of special interest, the well founded relations and the well ordering relations.

Definition 6.24.

- 1) $R \text{ Wfr } A \stackrel{\Delta}{\longleftrightarrow} R \text{ Fr } A \wedge (\forall x \in A) [\mathcal{M}(A \cap R^{-1}\{x\})]$.
- 2) $R \text{ We } A \stackrel{\Delta}{\longleftrightarrow} R \text{ Fr } A \wedge (\forall x \in A) (\forall y \in A) [x R y \vee x = y \vee y R x]$.
- 3) $R \text{ Wfwe } A \stackrel{\Delta}{\longleftrightarrow} R \text{ Wfr } A \wedge R \text{ We } A$.

Remark. R is a foundational relation on A iff each nonempty subset of A has an R -minimal element. R is a well founded relation on A iff each nonempty subset of A has an R -minimal element and each R -initial segment of A is a set. By an R -initial segment of A we mean the class of all elements in A that R -precede a given element of A i.e., $A \cap R^{-1}\{x\}$, $x \in A$. For example, each E -initial segment of V is a set, indeed

$$E^{-1}\{a\} = \{x \mid x \in a\} = a.$$

Then $a \cap x = a \cap E^{-1}\{x\}$ and hence from the Axiom of Regularity

$$a \neq 0 \rightarrow (\exists x \in a) [a \cap E^{-1}\{x\} = 0]$$

i.e., E is well founded on V .

There do exist foundational relations that are not well founded. Let A be the class of all finite sets and for $a, b \in A$ define aRb to mean that a has fewer elements than b . Given any nonempty collection of finite sets there is a set in the collection that has the least number of elements. Thus R is foundational. However the R -initial segment of A that contains all finite sets that R -precede a given doubleton set contains all singleton sets hence is a proper class. Thus R is not well founded.

R is a well ordering of A iff R determines an R -minimal element for each nonempty subset of A and the elements in A are pairwise R -comparable. If there were elements $a, b \in A$ that were not R -comparable i.e., neither aRb nor bRa , then both a and b would be R -minimal elements of $\{a, b\}$. Conversely if a and b are R -comparable then a and b cannot both be R -minimal elements of the same set. Thus if R well orders A then R determines a unique R -minimal element for each nonempty subset of A . That R is a transitive relation satisfying trichotomy we leave to the reader:

Theorem 6.25. $R \text{ We } A \rightarrow R \text{ Or } A.$

Exercises.

- 1) $R \text{ Fr } A \wedge B \subseteq A \rightarrow R \text{ Fr } B.$
- 2) $R \text{ Fr } a \rightarrow R \text{ Wfr } a.$
- 3) $R \text{ Wfr } A \wedge B \subseteq A \rightarrow R \text{ Wfr } B.$
- 4) $R \text{ We } A \wedge B \subseteq A \rightarrow R \text{ We } B.$
- 5) $R \text{ We } A \rightarrow (\forall x \in A) (\forall y \in A) [xRy \rightarrow \neg[x = y \vee yRx]].$
- 6) $R \text{ We } A \rightarrow (\forall x \in A) (\forall y \in A) (\forall z \in A) [xRy \wedge yRz \rightarrow xRz].$

Remark. If a relation R well orders a class A does it follow that R determines an R -minimal element for every nonempty subclass of A ? If R is a well founded well ordering of A , i.e. $R \text{ Wfwe } A$ then the answer is, Yes:

Theorem 6.26. $R \text{ Wfwe } A \wedge B \subseteq A \wedge B \neq 0 \rightarrow (\exists a \in B) [B \cap R^{-1}\{a\} = 0].$

Proof. Since $B \neq 0$, $\exists x \in B$. If $B \cap R^{-1}\{x\} \neq 0$ then since $B \cap R^{-1}\{x\}$ is a set

$$(\exists a \in B \cap R^{-1}\{x\}) [B \cap R^{-1}\{x\} \cap R^{-1}\{a\} = 0]$$

then $a \in B$ and aRx . Furthermore

$$y \in B \cap R^{-1}\{a\} \rightarrow yRa$$

and hence yRx . Thus

$$B \cap R^{-1}\{x\} \cap R^{-1}\{a\} = 0 \rightarrow B \cap R^{-1}\{a\} = 0.$$

Theorem 6.27.

$R \text{ Wfwe } A \wedge B \subseteq A \wedge (\forall x \in A) [A \cap R^{-1}\{x\} \subseteq B \rightarrow x \in B] \rightarrow A = B.$

Proof. If $A - B \neq 0$ then by Theorem 6.26

$$(\exists x \in A - B) [(A - B) \cap R^{-1}\{x\} = 0]$$

then

$$A \cap R^{-1}\{x\} \subseteq B.$$

Since $x \in A$ it follows from hypothesis that $x \in B$. But this contradicts the fact that $x \in A - B$.

Therefore $A - B = 0$ that is $A \subseteq B$. Since by hypothesis $B \subseteq A$ we conclude that $A = B$.

Remark. Theorem 6.27 is an induction principle. To prove that $(\forall x \in A) \varphi(x)$, we consider

$$B = \{x \in A \mid \varphi(x)\}.$$

If for any R that is a well founded well ordering on A we can prove

$$(\forall x \in A) [A \cap R^{-1}\{x\} \subseteq B \rightarrow x \in B]$$

it then follows that $A = B$ i.e. $(\forall x \in A) \varphi(x)$.

Later it will be shown that Theorems 6.26 and 6.27 are over hypothesized. We will prove that the hypothesis that $R \text{ Wfwe } A$ can be replaced by $R \text{ Fr } A$. See Theorem 9.20.

Clearly two relational systems (A, R_1) and (A, R_2) are essentially the same if R_1 and R_2 have the same relational part in common with A^2 i.e. if $A^2 \cap R_1 = A^2 \cap R_2$. Even if R_1 and R_2 do not have the same relational part there is a sense in which the two relational systems are equivalent. They may be equivalent in the sense that there exists an isomorphism between the two relational systems.

Definition 6.28. $H \text{ Isom}_{R_1, R_2}(A_1, A_2) \stackrel{\Delta}{\longleftrightarrow} H : A_1 \xrightarrow[\text{onto}]{1-1} A_2$
 $\wedge (\forall x \in A_1) (\forall y \in A_1) [x R_1 y \longleftrightarrow H(x) R_2 H(y)].$

Definition 6.29. $I \triangleq \{\langle x, x \rangle \mid x \in V\}.$

Theorem 6.30.

- 1) $(I \Gamma A) \text{ Isom}_{R, R}(A, A).$
- 2) $H \text{ Isom}_{R_1, R_2}(A_1, A_2) \rightarrow H^{-1} \text{ Isom}_{R_2, R_1}(A_2, A_1).$
- 3) $H_1 \text{ Isom}_{R_1, R_2}(A_1, A_2) \wedge H_2 \text{ Isom}_{R_2, R_3}(A_2, A_3)$
 $\rightarrow H_2 \circ H_1 \text{ Isom}_{R_1, R_3}(A_1, A_3).$

The proof is left to the reader.

Theorem 6.31. *If $H \text{ Isom}_{R_1, R_2}(A_1, A_2) \wedge B \subseteq A_1 \wedge x \in A_1$, then*

- 1) $B \cap R_1^{-1}\{x\} = 0 \iff H'' B \cap R_2^{-1}\{H'x\} = 0,$
- 2) $H''(A_1 \cap R_1^{-1}\{x\}) = A_2 \cap R_2^{-1}\{H'x\}.$

Proof.

$$1) \quad y \in B \cap R_1^{-1}\{x\} \rightarrow y \in B \wedge y R_1 x$$

$$\rightarrow H'y \in H'' B \wedge H'y R_2 H'x$$

$$\rightarrow H'y \in H'' B \cap R_2^{-1}\{H'x\}.$$

$$z \in H'' B \cap R_2^{-1}\{H'x\} \rightarrow (\exists y \in B) [z = H'y \wedge z R_2 H'x]$$

$$\rightarrow (\exists y \in B) [H'y R_2 H'x]$$

$$\rightarrow (\exists y \in B) [y R_1 x]$$

$$\rightarrow \exists y \in B \cap R_1^{-1}\{x\}.$$

$$2) \quad z \in H''(A_1 \cap R_1^{-1}\{x\}) \iff (\exists y) [y \in A_1 \wedge y R_1 x \wedge z = H'y]$$

$$\iff (\exists y \in A_1) [z = H'y \wedge H'y R_2 H'x]$$

$$\iff z \in A_2 \cap R_2^{-1}\{H'x\}.$$

Remark. Theorem 6.30 assures us that isomorphism between relational systems is an equivalence relation. From Theorem 6.31 we see that such isomorphisms preserve minimal elements and preserve initial segments. From this it is easy to prove the following. Details are left to the reader.

Theorem 6.32. *If $H \text{ Isom}_{R_1, R_2}(A_1, A_2)$ then*

$$1) \quad R_1 \text{ Fr } A_1 \iff R_2 \text{ Fr } A_2,$$

$$2) \quad R_1 \text{ Wfr } A_1 \iff R_2 \text{ Wfr } A_2,$$

$$3) \quad R_1 \text{ We } A_1 \iff R_2 \text{ We } A_2.$$

Remark. From Theorem 6.32 we see that if in a given equivalence class there is a relational system that is foundational then every relational system in that equivalence class is foundational. Similarly if there is a relational system that is well founded then all systems in that class are well founded; if one system is a well ordering all are well orderings.

Each equivalence class represents a particular type of ordering. Suppose that we are given a particular type of ordering, $[A_1, R_1]$ with $R_1 \subseteq A_1^2$ and a class A_2 , can we define an ordering on A_2 of the same type, that is, can we define a relation $R_2 \subseteq A_2^2$ such that the ordering $[A_2, R_2]$ is order isomorphic to the ordering $[A_1, R_1]$?

From the definition of order isomorphism we see that it is necessary that there exist a one-to-one correspondence between A_1 and A_2 . This is also sufficient.

Theorem 6.33.

If $H : A_1 \xrightarrow{\text{onto}} A_2 \wedge R_2 = \{\langle H'x, H'y \rangle \mid x \in A_1 \wedge y \in A_1 \wedge \langle x, y \rangle \in R_1\}$
 then

$$H \text{ Isom}_{R_1, R_2}(A_1, A_2).$$

The proof is left to the reader.

Remark. The relation R_2 in Theorem 6.33 is said to be induced on A_2 by the one-to-one function H and the relation R_1 , on A_1 . The theorem assures us that if a one-to-one correspondence exists between two classes then any type of ordering that can be defined on one class can also be defined on the other class. While this is a very useful result it leaves unanswered the question of what types of relations are definable on a given class A . Are there foundational relations definable on A ? Are there well founded relations on A ? Can A be well ordered? The first two questions are easily answered because the \in -relation is well founded on A .

The last question is the most interesting. From the work of Paul Cohen we know that the question of whether or not every set can be well ordered is undecidable in ZF. We will have more to say on this subject later.

Exercises.

1) If R_2 We A_2 then

$$H \text{ Isom}_{R_1, R_2}(A_1, A_2) \longleftrightarrow H : A_1 \xrightarrow{\text{onto}} A_2 \wedge$$

$$(\forall x \in A_1) [H'x \in A_2 - H''R_1^{-1}\{x\} \wedge (A_2 - H''R_1^{-1}\{x\}) \cap R_2^{-1}\{H'x\} = \emptyset].$$

7 Ordinal Numbers

The theory of ordinal numbers is essentially a theory of well ordered sets. For Cantor an ordinal number was “the general concept which results from (a well-ordered aggregate) M if we abstract from the nature of its elements while retaining their order of precedence” It was Gottlob Frege (1848–1925) and Bertrand Russell (1872–1970), working independently, who removed Cantor’s numbers from the realm of psychology. In 1903 Russell defined an ordinal number to be an equivalence class of well ordered sets under order isomorphism.

Our approach is that of von Neumann. We choose to define ordinal numbers to be particular members of equivalence classes rather than the equivalence classes themselves. We will show that every well ordered set is order isomorphic to exactly one set that is well ordered by the \in -relation and that is transitive in the following sense.

Definition 7.1. $\text{Tr}(A) \stackrel{\Delta}{\longleftrightarrow} (\forall x) [x \in A \rightarrow x \subseteq A]$.

Theorem 7.2. $\text{Tr}(A) \wedge B \in A \rightarrow B \subset A$.

Proof. $B \in A \rightarrow \mathcal{M}(B)$. By Definition 7.1, $\mathcal{M}(B) \wedge B \in A \wedge \text{Tr}(A) \rightarrow B \subseteq A$. But $B = A \rightarrow A \in A$ which is a contradiction. Therefore $B \subset A$.

Remark. We next define ordinal classes.

Definition 7.3. $\text{Ord}(A) \stackrel{\Delta}{\longleftrightarrow} \text{Tr}(A) \wedge E \text{ We } A$.

Remark. Since the \in -relation E is foundational, indeed well founded, on any class it follows that in order for E to well order A it is sufficient that E linearly order A .

Theorem 7.4.

$\text{Ord}(A) \longleftrightarrow \text{Tr}(A) \wedge (\forall x \in A) (\forall y \in A) [x \in y \vee x = y \vee y \in x]$.

Proof. Obvious from the definition of ordinal, the definition of well ordering and the fact that $E \text{ Fr } V$.

Theorem 7.5. $\text{Ord}(A) \wedge B \subseteq A \wedge B \neq 0 \rightarrow (\exists x \in B) [B \cap x = 0]$.

Proof. From Definition 7.3, E well orders A . Since E is also well founded on A i.e., E is foundational and E -initial segments of A are sets, it follows from Theorem 6.26 that B has an E -minimal element, i.e.,

$$(\exists x \in B) [B \cap E^{-1}\{x\} = 0].$$

But $E^{-1}\{x\} = x$.

Theorem 7.6. $\text{Ord}(A) \wedge a \in A \rightarrow \text{Ord}(a)$.

Proof. Since A is transitive $a \in A$ implies $a \subseteq A$. Then, since E well orders A , E well orders a . Furthermore, from transitivity $x \in y \wedge y \in a \rightarrow x \in A$. Since E well orders A we must have $x \in a \vee x = a \vee a \in x$. But $x = a \wedge x \in y \wedge y \in a$ and $a \in x \wedge x \in y \wedge y \in a$ each contradict Theorem 6.23. Hence $x \in y \wedge y \in a \rightarrow x \in a$, i.e., $y \in a \rightarrow y \subseteq a$.

Since a is transitive and well ordered by E , a is an ordinal.

Remark. We now wish to prove that the \in -relation also well orders the class of ordinal sets. From this and Theorem 7.6 it will then follow that the class of ordinal sets is an ordinal class.

Theorem 7.7. $\text{Ord}(A) \wedge \text{Tr}(B) \rightarrow [B \subset A \leftrightarrow B \in A]$.

Proof. By Theorem 7.2, $B \in A \rightarrow B \subset A$. Conversely if $B \subset A$ then $A - B \neq 0$. From Theorem 7.5, $A - B$ has an E -minimal element, i.e.,

$$(\exists x \in A - B) [(A - B) \cap x = 0].$$

Since $x \in A$ and A is transitive $x \subset A$. Since $(A - B) \cap x = 0$, $x \subseteq B$.

If $y \in B$ then since $B \subset A$, $y \in A$. But A is an ordinal class and $x \in A$. Therefore

$$y \in x \vee y = x \vee x \in y.$$

From the transitivity of B

$$[x \in y \vee x = y] \wedge y \in B \rightarrow x \in B$$

which contradicts the fact that $x \in A - B$. We conclude that $y \in x$, i.e., $B \subseteq x$.

Then $x = B \wedge x \in A$; hence $B \in A$.

Corollary 7.8. $\text{Ord}(A) \wedge \text{Ord}(B) \rightarrow [B \subset A \leftrightarrow B \in A]$.

Proof. $\text{Ord}(B) \rightarrow \text{Tr}(B)$.

Remark. Among other things Theorems 7.6 and 7.7 assure us that a transitive subclass of an ordinal is an ordinal.

Theorem 7.9. $\text{Ord}(A) \wedge \text{Ord}(B) \rightarrow \text{Ord}(A \cap B)$.

Proof. Since $A \cap B \subseteq A$ and $E \text{ We } A$, $E \text{ We } (A \cap B)$. Furthermore since A and B are transitive

$$\begin{aligned} x \in A \cap B &\rightarrow x \in A \wedge x \in B \\ &\rightarrow x \subset A \wedge x \subset B \\ &\rightarrow x \subset A \cap B. \end{aligned}$$

Therefore $A \cap B$ is transitive, and hence $A \cap B$ is an ordinal class.

Theorem 7.10. $\text{Ord}(A) \wedge \text{Ord}(B) \rightarrow [A \in B \vee A = B \vee B \in A]$.

Proof. $A \cap B \subseteq A \wedge A \cap B \subseteq B$. If $A \cap B \subset A \wedge A \cap B \subset B$ then $A \cap B \in A \wedge A \cap B \in B$ (Theorems 7.9 and 7.7) hence $A \cap B \in A \cap B$. But since by hypothesis E is foundational on A this contradicts Theorem 6.23.

Therefore $A \cap B = A$ or $A \cap B = B$, i.e., $B \subseteq A$ or $A \subseteq B$. Hence, by Corollary 7.8

$$A \in B \vee A = B \vee B \in A.$$

Definition 7.11. $\text{On} \triangleq \{x \mid \text{Ord}(x)\}$.

Theorem 7.12. $\text{Ord}(\text{On})$.

Proof. From Theorem 7.6,

$$x \in \text{On} \rightarrow x \subseteq \text{On}$$

i.e., On is transitive. From Theorems 7.6 and 7.10

$$x \in \text{On} \wedge y \in \text{On} \rightarrow x \in y \vee x = y \vee y \in x.$$

If $a \subseteq \text{On} \wedge a \neq 0$ then $(\exists x) [x \in a]$. But $x \in a \rightarrow \text{Ord}(x)$. If $a \cap x = 0$ then x is an E-minimal element of a . If $a \cap x \neq 0$ then since $a \cap x \subseteq x$ it follows that $a \cap x$ has an E-minimal element, i.e.,

$$(\exists y \in a \cap x) [(a \cap x) \cap y = 0].$$

Then $z \in a \cap y \rightarrow z \in a \wedge z \in y$. But since $y \in x \wedge \text{Ord}(x)$, $z \in y \rightarrow z \in x$, i.e.,

$$z \in a \cap y \rightarrow z \in (a \cap x) \cap y.$$

Therefore $a \cap y = 0$, i.e., y is an E-minimal element of a .

Thus E is a foundational relation on On and hence $E \text{ We } \text{On}$. Since On is transitive it is an ordinal class.

Remark. With the exception of Theorem 7.4 we have purposely avoided using the Axiom of Regularity in establishing our theory of ordinals. By so doing we will be able, at a later point, to prove the relative consistence of the Axiom of Regularity and the other axioms of ZF.

Theorem 7.13. $\mathcal{P}_r(\text{On})$.

Proof. $\mathcal{M}(\text{On}) \rightarrow \text{On} \in \text{On}$.

Corollary 7.14. $\text{Ord}(A) \rightarrow A \in \text{On} \vee A = \text{On}$.

Proof. From Theorems 7.10 and 7.12, $A \in \text{On} \vee A = \text{On} \vee \text{On} \in A$.
But $\mathcal{P}_r(\text{On}) \rightarrow \text{On} \notin A$.

Corollary 7.15. $\text{Ord}(A) \rightarrow A \subseteq \text{On}$.

Proof. Corollary 7.14, Theorem 7.12 and Corollary 7.8.

Remark. The elements of On are the ordinal numbers in our system. We have proved that every ordinal class is an ordinal number except one, On. The ordinal numbers play such an important role in the theory ahead that we find it convenient to use the symbols

$$\alpha, \beta, \gamma, \dots$$

as variables on ordinal numbers.

Definition 7.16

- 1) $(\forall \alpha) \varphi(\alpha) \stackrel{\Delta}{\leftrightarrow} (\forall x) [\text{Ord}(x) \rightarrow \varphi(x)]$.
- 2) $(\exists \alpha) \varphi(\alpha) \stackrel{\Delta}{\leftrightarrow} (\exists x) [\text{Ord}(x) \wedge \varphi(x)]$.

Remark. To make clear in what sense Definition 7.16 is a definition, that is to make clear precisely what wff $(\forall \alpha) \varphi(\alpha)$ is an abbreviation for, we could standardize our list of ordinal variables:

$$\alpha_0, \alpha_1, \dots$$

and insist that our intention is that

$$\begin{aligned} (\forall \alpha_n) \varphi(\alpha_n) & \text{ is } (\forall x_n) [\text{Ord}(x_n) \rightarrow \varphi(x_n)]. \\ (\exists \alpha_n) \varphi(\alpha_n) & \text{ is } (\exists x_n) [\text{Ord}(x_n) \wedge \varphi(x_n)]. \end{aligned}$$

Theorem 7.17. (*The Principle of Transfinite Induction.*) If 1) $A \subseteq \text{On}$ and 2) $(\forall \alpha) [\alpha \subseteq A \rightarrow \alpha \in A]$ then $A = \text{On}$.

Proof. If $\text{On} - A \neq 0$ then $(\exists \alpha \in \text{On} - A) [(\text{On} - A) \cap \alpha = 0]$. Since $\alpha \subset \text{On}$ it follows that $\alpha \subseteq A$. Then by 2), $\alpha \in A$. But this contradicts the fact that $\alpha \in \text{On} - A$. Therefore $\text{On} - A = 0$, i.e., $\text{On} \subseteq A$. Then from 1), $A = \text{On}$.

Definition 7.18.

- 1) $\alpha < \beta \stackrel{\Delta}{\leftrightarrow} \alpha \in \beta$.
- 2) $\alpha \leq \beta \stackrel{\Delta}{\leftrightarrow} \alpha < \beta \vee \alpha = \beta$.
- 3) $\max(\alpha, \beta) = \alpha \cup \beta$.

Exercises. Prove the following.

- 1) $(\forall \alpha) [\alpha \subset \text{On}]$.
- 2) $(\forall \alpha) (\forall \beta) [\alpha < \beta \vee \alpha = \beta \vee \beta < \alpha]$.
- 3) $\alpha \leq \beta \wedge \beta < \gamma \rightarrow \alpha < \gamma$.
- 4) $\alpha < \beta \wedge \beta \leq \gamma \rightarrow \alpha < \gamma$.
- 5) $\alpha \leq \beta \leftrightarrow \alpha \subseteq \beta$.
- 6) $(\forall \alpha) (\exists \beta) [\beta = \alpha \cup \{\alpha\}]$.
- 7) $\text{Tr}(A) \leftrightarrow (\forall x) (\forall y) [x \in y \wedge y \in A \rightarrow x \in A]$.
- 8) $\text{Ord}(A) \wedge a \in A \rightarrow a = a \cap A$.
- 9) $A \subseteq \text{On} \wedge A \neq 0 \rightarrow (\exists \alpha) [\alpha \in A \wedge (\forall \beta) [\beta \in A \rightarrow \alpha \leq \beta]]$.
- 10) $\text{Tr}(A) \rightarrow [\cup(A) \subseteq A]$.
- 11) $\cup(\text{On}) = \text{On}$.
- 12) $\text{Ord}(\max(\alpha, \beta))$.
- 13) $\alpha \leq \max(\alpha, \beta) \wedge \beta \leq \max(\alpha, \beta)$.
- 14) $\alpha = \max(\alpha, \beta) \vee \beta = \max(\alpha, \beta)$.

Theorem 7.19. $A \subseteq \text{On} \rightarrow \text{Ord}(\cup(A))$.

Proof. $x \in \cup(A) \rightarrow (\exists y) [x \in y \wedge y \in A]$. But $y \in A \rightarrow y \subseteq \cup(A)$ and, since $A \subseteq \text{On}$, $x \in y \wedge y \in A \rightarrow x \subseteq y$. Therefore $x \subseteq \cup(A)$, i.e., $\cup(A)$ is transitive. Furthermore since $A \subseteq \text{On} \wedge \text{Tr}(\text{On})$

$$x \in A \rightarrow x \subseteq \text{On}$$

i.e., $\cup(A) \subseteq \text{On}$. Since $\cup(A)$ is a transitive subclass of the ordinal On it follows that $\cup(A)$ is an ordinal class.

Theorem 7.20. $A \subseteq \text{On} \wedge \alpha \in A \rightarrow \alpha \leq \cup(A)$.

Proof. $\alpha \in A \rightarrow \alpha \subseteq \cup(A)$
 $\rightarrow \alpha \leq \cup(A)$.

Theorem 7.21. $A \subseteq \text{On} \wedge (\forall \beta) [\beta \in A \rightarrow \beta \leq \alpha] \rightarrow \cup(A) \leq \alpha$.

Proof. $\beta \in \cup(A) \rightarrow (\exists \gamma) [\beta < \gamma \wedge \gamma \in A]$. Therefore $\beta < \gamma \wedge \gamma \leq \alpha$. Hence $\beta < \alpha$, i.e., $\cup(A) \subseteq \alpha$.

Remark. Theorem 7.20 assures us that $\cup(A)$ is an upper bound for the class of ordinals A . Theorem 7.21 assures us that $\cup(A)$ is also the smallest upper bound for A . Furthermore if A has a maximal ordinal, i.e., if $(\exists \alpha) [\alpha \in A \wedge (\forall \beta) [\beta \in A \rightarrow \beta \leq \alpha]]$ then $\alpha = \cup(A)$ and hence $\cup(A) \in A$. If A has no maximal element then $\cup(A) \notin A$. In particular if A

is an ordinal number, i.e., $(\exists \alpha) [\alpha = A]$ and if $\cup(\alpha) \notin \alpha$ then $\cup(\alpha) = \alpha$. Such an ordinal is called a *limit ordinal*.

Definition 7.22. $\alpha + 1 \triangleq \alpha \cup \{\alpha\}$.

Example. $0 + 1 = 0 \cup \{0\} = \{0\} \triangleq 1$

$1 + 1 = 1 \cup \{1\} = \{0\} \cup \{1\} = \{0, 1\} \triangleq 2$.

Theorem 7.23. $(\forall \alpha) [\alpha \in \alpha + 1 \wedge \alpha \subseteq \alpha + 1]$.

Proof. $\alpha \in (\alpha \cup \{\alpha\}) \wedge \alpha \subseteq (\alpha \cup \{\alpha\})$.

Theorem 7.24. $(\forall \alpha) [\alpha + 1 \in \text{On}]$.

Proof. $(\forall \alpha) [\alpha \in \text{On} \wedge \alpha \subseteq \text{On}]$. Therefore $\alpha + 1 \subseteq \text{On}$. Furthermore

$$\begin{aligned} x \in \alpha + 1 &\rightarrow x \in \alpha \vee x = \alpha \\ &\rightarrow x \subseteq \alpha. \end{aligned}$$

But $\alpha \subseteq \alpha + 1$, hence $x \subseteq \alpha + 1$. Thus $\alpha + 1$ is a transitive subset of an ordinal class hence an ordinal, i.e., $\alpha + 1 \in \text{On}$.

Theorem 7.25. $\neg [\alpha < \beta < \alpha + 1]$.

Proof. If $\alpha < \beta \wedge \beta < \alpha + 1$ then $\alpha \in \beta \wedge [\beta \in \alpha \vee \beta = \alpha]$. But this contradicts the fact that $E \text{ Fr On}$.

Theorem 7.26. $a \subset \text{On} \rightarrow (\forall \alpha) [\alpha \in a \rightarrow \alpha < \cup(a) + 1]$.

Proof. From Theorems 7.19 and 7.20 we have $\text{Ord}(\cup(a)) \wedge \alpha \leq \cup(a)$. From Theorem 7.23 we conclude that $\alpha < \cup(a) + 1$.

Remark. Theorem 7.26 assures us that given an ordinal number there is an ordinal number that is larger, indeed given any set of ordinal numbers there is an ordinal number that is larger than each element of the given set. The naive acceptance of On as a set then leads to the paradox of the largest ordinal, i.e., an ordinal number that is larger than every ordinal including itself.

This paradox was discovered by Cantor in 1895. It first appeared in print in 1897 having been rediscovered by Burali-Forti, whose name it now bears.

Definition 7.27. $K_1 \triangleq \{\alpha \mid \alpha = 0 \vee (\exists \beta) [\alpha = \beta + 1]\}$

$K_{II} \triangleq \text{On} - K_1$.

Definition 7.28. $\omega \triangleq \{\alpha \mid \alpha \cup \{\alpha\} \subseteq K_1\}$.

Exercises. Prove the following.

- 1) $\alpha \in K_{II} \leftrightarrow \alpha \neq 0 \wedge \alpha = \cup(\alpha).$
- 2) $\omega \neq 0.$
- 3) $(\forall \alpha) [\alpha + 1 \in K_I].$
- 4) $\omega \subseteq K_I.$
- 5) $\beta \in \omega \leftrightarrow (\forall \gamma) [\gamma \leq \beta \rightarrow \gamma \in K_I].$
- 6) $\beta < \alpha \wedge \alpha \in K_{II} \rightarrow (\exists \gamma) [\beta < \gamma < \alpha].$
- 7) $\alpha \subseteq K_I \rightarrow \alpha + 1 \subseteq K_I.$

Remark. From Exercise 1 we see that K_{II} is the class of all limit ordinals. We will refer to the elements of K_{II} as limit ordinals and to the elements of K_I as non-limit ordinals. The elements in ω are the natural numbers or non-negative integers as we will now show by proving that they satisfy the Peano postulates. As a notational convenience we will use

$$i, j, k, \dots, m, n$$

as variables on ω .

Definition 7.29. $(\forall i) \varphi(i) \stackrel{\Delta}{\leftrightarrow} (\forall x) [x \in \omega \rightarrow \varphi(x)]$
 $(\exists i) \varphi(i) \stackrel{\Delta}{\leftrightarrow} (\exists x) [x \in \omega \wedge \varphi(x)].$

Theorem 7.30 (*Peano's Postulates*).

- 1) $0 \in \omega.$
- 2) $(\forall i) [i + 1 \in \omega].$
- 3) $(\forall i) [i + 1 \neq 0].$
- 4) $(\forall i) (\forall j) [i + 1 = j + 1 \leftrightarrow i = j].$
- 5) $0 \in A \wedge (\forall i) [i \in A \rightarrow i + 1 \in A] \rightarrow \omega \subseteq A.$

Proof.

1) From Definition 7.27 we have that $0 \in K_I$. Therefore $0 \cup \{0\} \subseteq K_I$ hence $0 \in \omega$.

2) Since $i + 1 \in K_I$ and since $i \in \omega \rightarrow i + 1 \subseteq K_I$ it follows that $(i + 1) \cup \{i + 1\} \subseteq K_I$, i.e., $i + 1 \in \omega$.

3) $i \in i + 1$. Therefore $i + 1 \neq 0$.

4) $i = j \rightarrow (i + 1) = (j + 1)$. Conversely if $i + 1 = j + 1$ then since $i \in i + 1$ we have $i \in j \vee i = j$ and $j \in i \vee j = i$. Since $i \in j \wedge j \in i$ and $i \in j \wedge j = i$ each contradict the fact that E is foundational on On we conclude that $i = j$.

5) If $\omega - A \neq 0$ then there is a smallest element i in $\omega - A$. Since $0 \in A$ it follows that $0 \neq i$. But $i \in \omega \wedge \omega \subseteq K_1$; therefore $(\exists \beta) [i = \beta + 1]$. Furthermore $i \subseteq i + 1$ and $i + 1 \subseteq K_1$. Therefore $\beta + 1 \subseteq K_1$, i.e., $\beta \in \omega$. Since $\beta < i$ and i is the smallest element in $\omega - A$ we must have $\beta \notin \omega - A$, i.e., $\beta \in A$. But $\beta \in \omega \wedge \beta \in A \rightarrow \beta + 1 \in A$ that is $i \in A$. This is a contradiction that forces us to conclude that $\omega \subseteq A$.

Corollary 7.31. (*The Principle of Finite Induction.*)

If $A \subseteq \omega \wedge 0 \in A \wedge (\forall i) [i \in A \rightarrow i + 1 \in A]$ then $A = \omega$.

Proof. Obvious from Theorem 7.30.

Theorem 7.32. $\text{Ord}(\omega)$.

Proof. Since $\omega \subseteq K_1 \wedge K_1 \subseteq \text{On}$ we have that $\omega \subseteq \text{On}$. Furthermore $x \in y \wedge y \in \omega \rightarrow \text{Ord}(y) \wedge y + 1 \subseteq K_1$. Therefore $x \subseteq y \wedge y \subseteq y + 1 \wedge y + 1 \subseteq K_1$ hence $x \in K_1 \wedge x \subseteq K_1$. Consequently $x \cup \{x\} \subseteq K_1$, that is $x \in \omega$. Thus $\omega \subseteq \text{On} \wedge \text{Tr}(\omega)$; therefore $\text{Ord}(\omega)$.

Remark. From Theorem 7.32 we see that either $\omega \in \text{On} \vee \omega = \text{On}$. But which of these alternatives is true? The question is whether or not ω is a set. This cannot be resolved by the axioms stated thus far. We choose to resolve the issue by postulating that ω is a set.

Axiom 7 (Axiom of Infinity). $\mathcal{M}(\omega)$.

Theorem 7.33. $\omega \in K_{II}$.

Proof. From Axiom 7 and Theorem 7.31 we have that $\omega \in \text{On}$. Since $\omega \subseteq K_1$ it follows that if $\omega \in K_1$ then $\omega + 1 \subseteq K_1$ and hence $\omega \in \omega$. Therefore $\omega \in K_{II}$.

Theorem 7.34. $F \text{ Fn } \omega \rightarrow (\exists n) [F(n + 1) \notin F(n)]$.

Proof. From the Axiom of Infinity, F is a function whose domain is a set. Therefore the range of F , $\mathcal{W}(F)$, is a set. Then from the Axiom of Regularity

$$(\exists x \in \mathcal{W}(F)) [\mathcal{W}(F) \cap x = 0].$$

But $x \in \mathcal{W}(F) \rightarrow (\exists n) [x = F(n)]$. Since $F(n + 1) \in \mathcal{W}(F)$ we have $F(n + 1) \notin F(n)$.

Remark. Theorem 7.34 assures us that there are no infinite descending \in -chains. Given a non-empty set a_0 , $\exists a_1 \in a_0$. If $a_1 \neq 0$, $\exists a_2 \in a_1$, etc. However, by Theorem 7.34, we see that after a finite number of steps we must arrive at a set $a_n \in a_{n-1}$ with $a_n = 0$.

While every descending \in -chain must be of finite length it does not follow that for a given set a there is a bound on the length of the \in -chains descending from a . Consider, for example, ω :

$$(\forall n) 0 \in 1 \in 2 \in \dots \in n \in \omega.$$

Exercises. Prove the following.

- 1) $\alpha \in K_{\text{II}} \rightarrow \omega \leq \alpha.$
- 2) $A \subseteq \omega \wedge A \neq 0 \rightarrow (\exists k \in A) (\forall i \in A) [k \leq i].$
- 3) $F : \omega \rightarrow A \wedge R \text{ Fr } A \rightarrow (\exists n) [F(n+1) \notin R^{-1}\{F \cdot n\}].$

Definition 7.35. $\cap(A) \triangleq \{x \mid (\forall y \in A) [x \in y]\}.$

Definition 7.36.

$$\begin{aligned} \sup(A) &\triangleq \cup(A \cap \text{On}) \text{ if } A \cap \text{On} \neq 0. \\ &\triangleq 0 \quad \text{if } A \cap \text{On} = 0. \end{aligned}$$

$$\begin{aligned} \inf(A) &\triangleq \cap(A \cap \text{On}) \text{ if } A \cap \text{On} \neq 0. \\ &\triangleq 0 \quad \text{if } A \cap \text{On} = 0. \end{aligned}$$

$$\begin{aligned} \sup_{<\beta}(A) &\triangleq \cup(A \cap \beta) \text{ if } A \cap \beta \neq 0. \\ &\triangleq 0 \quad \text{if } A \cap \beta = 0 \end{aligned}$$

$$\begin{aligned} \inf_{>\beta}(A) &\triangleq \cap(A \cap \text{On}) - \beta \text{ if } (A \cap \text{On}) - \beta \neq 0. \\ &\triangleq 0 \quad \text{if } (A \cap \text{On}) - \beta = 0. \end{aligned}$$

Definition 7.37. $\mu_\alpha(\varphi(\alpha)) = \inf\{\alpha \mid \varphi(\alpha)\}.$

Exercises. Prove the following.

- 1) $\cap\{a, b\} = a \cap b.$
- 2) $\text{Tr}(A) \rightarrow \cap(A) \subseteq A.$
- 3) $a \in A \rightarrow \cap(A) \subseteq a.$
- 4) $\mathcal{M}(\cap a).$
- 5) $A = 0 \rightarrow \cap(A) = 0.$
- 6) $\sup(A) \in \text{On} \vee \sup(A) = \text{On}.$
- 7) $\sup_{<\beta}(A) \in \text{On}.$
- 8) $\inf(A) \in \text{On} \wedge \inf_{>\beta}(A) \in \text{On}.$
- 9) $\inf(A) \in A \iff A \cap \text{On} \neq 0.$
- 10) $(\exists \alpha) \varphi(\alpha) \rightarrow \varphi(\mu_\alpha(\varphi(\alpha))).$
- 11) $\text{Ord}(A) \wedge A - \alpha \neq 0 \rightarrow \alpha = \inf(A - \alpha).$

Remark. Every ordinal is well ordered by the \in -relation, that is $(\forall \alpha) [E \Gamma (\alpha \times \alpha) \text{ We } \alpha].$ Thus $\langle E \Gamma (\alpha \times \alpha), \alpha \rangle$ is a member of some equivalence class determined by order isomorphism. We now wish to prove that there is exactly one such pair in each equivalence class of well

ordered sets, that is, if $r \in a$ then there is one and only one ordinal α for which

$$(\exists f) [f \text{ Isom}_{E,r}(\alpha, a)]$$

We first prove that there is at most one such ordinal.

Theorem 7.38. $\text{Ord}(A) \wedge \text{Ord}(B) \wedge F \text{ Isom}_{E,E}(A, B) \rightarrow A = B.$

Proof. It is sufficient to prove that F is the identity function I restricted to A . This we prove by transfinite induction. If $\beta \in A \wedge (\forall \alpha < \beta) [F' \alpha = \alpha]$ then $F' \beta = \beta$. Furthermore since β is the E -minimal element in $A - \beta$ and order isomorphisms map minimal elements onto minimal elements it follows that $F' \beta$ is the E -minimal element in $B - F' \beta = B - \beta$, i.e., $F' \beta = \beta$.

Theorem 7.39.

$$\text{Ord}(A) \wedge \text{Ord}(B) \wedge F_1 \text{ Isom}_{E,R}(A, C) \wedge F_2 \text{ Isom}_{E,R}(B, C) \rightarrow A = B.$$

Proof. Theorem 7.38 and the fact that $F_2 \circ F_1^{-1} \text{ Isom}_{E,E}(A, B)$.

Remark. Theorem 7.39 assures us that every well ordered class is order isomorphic to at most one ordinal well ordered by the \in -relation. We will prove that while not all well ordered classes are order isomorphic to an ordinal every well ordered set is.

Since order isomorphisms must map minimal elements into minimal elements and initial segments into initial segments we must show that if $R \in a$ then there exists an ordinal number α and a function F mapping α onto a in such a way that $\forall \beta < \alpha, F' \beta$ is the R -minimal element in $a - F' \beta$. But this means that F must be so defined that its value at β depends on β , the values F assumes at all ordinals smaller than β , and the additional requirement that $F' \beta$ be the R -minimal element in $a - F' \beta$. Does there exist a function fulfilling all of these requirements? We will prove that such a function does exist. Indeed we will prove that there exists exactly one function F defined on On in such a way that its value at β depends upon β , and upon the values F assumes at all ordinals smaller than β , hence upon $F \upharpoonright \beta$, and also depends upon any previously given condition G . Such a function F is said to be defined by transfinite recursion.

We need the following lemma.

Lemma 1. $f \mathcal{F}_n \beta \wedge (\forall \alpha < \beta) [f(\alpha) = G(f \upharpoonright \alpha)] \wedge g \mathcal{F}_n \gamma$
 $\wedge (\forall \alpha < \gamma) [g(\alpha) = G(g \upharpoonright \alpha)] \wedge \beta \leq \gamma \rightarrow (\forall \alpha < \beta) [f' \alpha = g' \alpha].$

Proof (by induction). If $(\forall \gamma < \alpha) [f' \gamma = g' \gamma]$ then $f \upharpoonright \alpha = g \upharpoonright \alpha$ and hence $f' \alpha = G(f \upharpoonright \alpha) = G(g \upharpoonright \alpha) = g' \alpha$.

Theorem 7.40 (*Principle of Transfinite Recursion*). If $K = \{f \mid (\exists \beta) [f \mathcal{F}_n \beta \wedge (\forall \alpha < \beta) [f' \alpha = G'(f \upharpoonright \alpha)]]\}$ and if $F = \cup(K)$ then

- 1) $F \mathcal{F}_n \text{ On}$.
- 2) $(\forall \alpha) [F' \alpha = G'(F \upharpoonright \alpha)]$.
- 3) $A \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [A' \alpha = G'(A \upharpoonright \alpha)] \rightarrow A = F$.

Proof.

- 1) Since $f \in K \rightarrow \mathcal{R}el(f)$, we have that F is a relation. Furthermore $\langle a, b \rangle \in F \wedge \langle a, c \rangle \in F \rightarrow (\exists f \in K) (\exists g \in K) [\langle a, b \rangle \in f \wedge \langle a, c \rangle \in g]$.

From Lemma 1 we have that $f(a) = g(a)$, i.e., $b = c$. Thus F is single valued hence is a function.

If $x \in a \wedge a \in \mathcal{D}(F)$ then from the definition of F and K it follows that $(\exists f) (\exists \beta) [f \mathcal{F}_n \beta \wedge a \in \beta]$. But ordinals are transitive and the elements of ordinals are ordinals, i.e., $\text{Ord}(a) \wedge x \in \beta$. Thus $\mathcal{D}(F) \subseteq \text{On} \wedge \text{Tr}(\mathcal{D}(F))$. Therefore $\text{Ord}(\mathcal{D}(F))$, i.e., $\mathcal{D}(\mathcal{F}) = \text{On}$ or $(\exists \gamma) [\gamma = \mathcal{D}(F)]$. If $\gamma = \mathcal{D}(F)$ and if

$$g = F \cup \{\langle \gamma, G'(F \upharpoonright \gamma) \rangle\}$$

then $g \mathcal{F}_n(\gamma + 1) \wedge (\forall \alpha < \gamma + 1) [g' \alpha = G'(g \upharpoonright \alpha)]$. Thus $g \in K$ and $\gamma \in \mathcal{D}(F)$. But $\gamma = \mathcal{D}(F)$. This is a contradiction; hence $\mathcal{D}(F) = \text{On}$ and $F \mathcal{F}_n \text{ On}$.

2) $(\forall \alpha) (\exists f) [f \subseteq F \wedge F' \alpha = f' \alpha \wedge f' \alpha = G'(f \upharpoonright \alpha)]$. It then follows that $F \upharpoonright \alpha = f \upharpoonright \alpha$ and hence $F' \alpha = f' \alpha = G'(f \upharpoonright \alpha) = G'(F \upharpoonright \alpha)$.

3) Since $\mathcal{D}(A) = \mathcal{D}(F)$ it is sufficient to prove that $(\forall \alpha) [A' \alpha = F' \alpha]$. This we do by transfinite induction. If $(\forall \alpha) [\alpha < \gamma \rightarrow A' \alpha = F' \alpha]$ then $A \upharpoonright \gamma = F \upharpoonright \gamma$. Therefore $A' \gamma = G(A \upharpoonright \gamma) = G(F \upharpoonright \gamma) = F' \gamma$.

Corollary 7.41. $(\exists! f) [f \mathcal{F}_n \alpha \wedge (\forall \beta < \alpha) [f' \beta = G(f \upharpoonright \beta)]]$.

The proof is left to the reader.

Remark. Theorem 7.40 is a theorem schema and hence a metatheorem. With quantification in the metalanguage it could be stated as

$$(\forall G) (\exists! F) [F \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [F' \alpha = G(F \upharpoonright \alpha)]] .$$

While this statement more readily conveys the content of the theorem as it will be used it is nevertheless of interest to note that Theorem 7.40 as stated is stronger. It not only asserts the existence of the function F but prescribes a method for exhibiting F when G is given.

For certain types of problems it is sufficient to know that functions can be defined recursively on the natural numbers in the following sense. Given any function h and any set a there is a function f defined on ω in such a way that

$$\begin{aligned} f'0 &= a \\ f'(n+1) &= h' f'n . \end{aligned}$$

This type of recursion can be extended to functions on On by requiring that at limit ordinals α the value of f be the supremum of its values at the ordinals preceding α , i.e.,

$$f' a = \bigcup_{\gamma < \alpha} f' \gamma, \quad \alpha \in K_{\text{II}}.$$

We wish to prove the following:

$$(\forall H) (\forall a) (\exists F) \left[F \mathcal{F}_n \text{On} \wedge F'0 = a \wedge F'(\beta + 1) = H' F' \beta \right. \\ \left. \wedge \left[\beta \in K_{\text{II}} \rightarrow F' \beta = \bigcup_{\gamma < \beta} F' \gamma \right] \right].$$

For the statement of our theorem we choose to retain the format of a theorem schema and hence we resort to the standard circumlocutions to avoid quantification on class symbols.

Theorem 7.42. *If $G = \{ \langle x, y \rangle \mid [x = 0 \wedge y = a] \vee [x \neq 0 \wedge \sup(\mathcal{D}(x)) \neq \mathcal{D}(x) \wedge y = H' x' \sup(\mathcal{D}(x))] \vee [x \neq 0 \wedge \sup(\mathcal{D}(x)) = \mathcal{D}(x) \wedge y = \cup \mathcal{W}(x)] \}$ and $F \mathcal{F}_n \text{On} \wedge (\forall \alpha) [F' \alpha = G'(F \upharpoonright \alpha)]$ then*

- 1) $F'0 = a.$
- 2) $F'(\beta + 1) = H' F' \beta.$
- 3) $F' \beta = \bigcup_{\gamma < \beta} F' \gamma, \beta \in K_{\text{II}}.$
- 4) F is unique.

Proof.

- 1) $F'0 = G'(F \upharpoonright 0) = G'(0) = a.$
- 2) $F'(\beta + 1) = G'(F \upharpoonright (\beta + 1)).$ Since $\mathcal{D}(F \upharpoonright (\beta + 1)) = \beta + 1$ and $\sup(\beta + 1) = \beta \neq \beta + 1$ we have that

$$G'(F \upharpoonright (\beta + 1)) = H'(F \upharpoonright (\beta + 1))' \sup(\beta + 1) = H' F' \beta,$$

i.e., $F'(\beta + 1) = H' F' \beta.$

- 3) $F' \beta = G'(F \upharpoonright \beta).$ Since $\mathcal{D}(F \upharpoonright \beta) = \beta$ and $\beta \in K_{\text{II}}$ we have that $\sup(\beta) = \beta.$ Hence $G'(F \upharpoonright \beta) = \cup(\mathcal{W}(F \upharpoonright \beta)) = \bigcup_{\gamma < \beta} F' \gamma,$ i.e., $F' \beta = \bigcup_{\gamma < \beta} F' \gamma.$

- 4) Obvious by induction.

Corollary 7.43 (*Principle of Finite Recursion*).

$$(\exists! f) [f \mathcal{F}_n \omega \wedge f'0 = a \wedge (\forall k) [f'(k + 1) = H' f'k]].$$

Proof. If in Theorem 7.42 we restrict F to ω then $(F \upharpoonright \omega) \mathcal{F}_n \omega$ hence $\mathcal{M}(F \upharpoonright \omega),$ i.e., $(\exists f) [f = F \upharpoonright \omega].$ Then $f \mathcal{F}_n \omega \wedge f'0 = a \wedge (\forall k) [f'(k + 1) = H' f'k].$

It is obvious, by induction, that f is unique.

Remark. In the study of order isomorphisms we are especially interested in those order preserving functions that map ordinals onto ordinals. Any function whose domain is an ordinal and whose range is a class of ordinal numbers we will call an *ordinal function*. If in addition an ordinal function is order preserving we say that it is strictly monotone.

Definition 7.44.

$$\text{Orf}(G) \stackrel{\Delta}{\longleftrightarrow} G \mathcal{F}_n \mathcal{D}(G) \wedge \text{Ord}(\mathcal{D}(G)) \wedge \mathcal{W}(G) \subseteq \text{On}.$$

$$\text{Smo}(G) \stackrel{\Delta}{\longleftrightarrow} \text{Orf}(G) \wedge (\forall \alpha \in \mathcal{D}(G)) (\forall \beta \in \mathcal{D}(G)) [\alpha < \beta \rightarrow G'\alpha < G'\beta].$$

Exercises. *Prove the following.*

- 1) $\text{Smo}(G) \rightarrow (\forall \alpha) [\alpha \in \mathcal{D}(G) \rightarrow \alpha \leq G'\alpha].$
- 2) If in Theorem 7.42, $a \in \text{On}$ and H is a strictly monotone ordinal function on On then F is a strictly monotone ordinal function on On .
- 3) $F \text{ Isom}_{E,E}(A, B) \wedge \text{Ord}(A) \wedge B \subseteq \text{On} \rightarrow \text{Smo}(F).$
- 4) State and prove a generalization of Theorem 7.40 in which On is replaced by a well ordered class A .

Remark. The principle of transfinite recursion assures us that we can define a function F on On in such a way that its value at α is dependent on its values at all ordinals less than α and on any given condition G . If $R \text{ We } A$ and if F is to be an order preserving isomorphism from some ordinal onto A then $F'\alpha$ must be the R -minimal element in $A - F''\alpha$. Suppose then that we could define G in such a way that $G(F \upharpoonright \alpha)$ is the R -minimal element in $A - F''\alpha$. Then clearly F would be an order preserving map from ordinals into A . It would then only remain to be proved that A is exhausted, i.e., $\mathcal{W}(F) = A$. In fact we will discover that there are two cases of interest. If R is a well founded relation then $\mathcal{W}(F) = A$; if R is not well founded then $\mathcal{W}(F)$ will be an R -initial segment of A .

Note that $\mathcal{W}(F)$ is R -transitive, that is $x R y \wedge y \in \mathcal{W}(F) \rightarrow x \in \mathcal{W}(F)$.

Theorem 7.45.

If $R \text{ We } A \wedge R \text{ Wfr } A \wedge B \subseteq A \wedge (\forall x \in A) (\forall y \in B) [x R y \rightarrow x \in B]$ then

$$A = B \vee (\exists x \in A) [B = A \cap R^{-1}\{x\}].$$

Proof. If $A \neq B$ then, since $B \subseteq A$, $A - B \neq 0$. Thus $A - B$ has an R -minimal element, a (Theorem 6.26). Since $a \in A \wedge a \notin B$ and, by hypothesis

$$x \in A \wedge x R y \wedge y \in B \rightarrow x \in B$$

it follows that

$$(\forall x \in B) [x R a]$$

for otherwise $a \in B$. Therefore $B \subseteq [A \cap R^{-1}\{a\}]$.

Furthermore $x \in [A \cap R^{-1}\{a\}] \rightarrow x \in A \wedge x R a$. But a is the R -minimal element in $A - B$. Therefore $x \notin A - B$, i.e., $x \in B$. Thus $[A \cap R^{-1}\{a\}] \subseteq B$ and hence $B = [A \cap R^{-1}\{a\}]$.

Remark. In Theorem 7.45 the requirement that R be well founded is only used to establish that $A - B$ has an R -minimal element. Later we will show that it is sufficient for R to be foundational. Consequently this result follows if R is a well ordering relation that is not well founded.

Theorem 7.46.

If $F \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [F'\alpha = G'(F \Gamma \alpha)] \wedge (\forall \alpha) [G'(F \Gamma \alpha) \in [A - F''\alpha]]$ then

- 1) $\mathcal{W}(F) \subseteq A$.
- 2) $\mathcal{U}_{n_2}(F)$.
- 3) $\mathcal{P}_r(A)$.

Proof.

- 1) $(\forall \alpha) [F'\alpha = G'(F \Gamma \alpha) \wedge G'(F \Gamma \alpha) \in [A - F''\alpha]]$, i.e., $(\forall \alpha) [F'\alpha \in A]$.
- 2) $\alpha < \beta \rightarrow F'\alpha \in F''\beta$. Furthermore $F'\beta \in [A - F''\beta]$, i.e., $F'\beta \notin F''\beta$. Therefore $F'\alpha \neq F'\beta$. Consequently $\alpha \neq \beta \rightarrow F'\alpha \neq F'\beta$, i.e., $F'\alpha = F'\beta \rightarrow \alpha = \beta$.

3) Since $\mathcal{W}(F) \subseteq A$ we have that $\mathcal{M}(A) \rightarrow \mathcal{M}(\mathcal{W}(F))$. Furthermore $\mathcal{U}_{n_2}(F) \rightarrow [\mathcal{M}(\mathcal{W}(F)) \leftrightarrow \mathcal{M}(\text{On})]$. Thus $\mathcal{P}_r(A)$.

Theorem 7.47. If $F \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [F'\alpha = G'(F \Gamma \alpha)] \wedge (\forall \alpha) [A - F''\alpha \neq 0 \rightarrow G'(F \Gamma \alpha) \in A - F''\alpha] \wedge \mathcal{M}(A)$ then

$$(\exists \alpha) (\forall \beta < \alpha) [A - F''\beta \neq 0] \wedge F''\alpha = A \wedge \mathcal{U}_{n_2}(F \Gamma \alpha).$$

Proof. If $(\forall \alpha) [A - F''\alpha \neq 0]$ then $(\forall \alpha) [G'(F \Gamma \alpha) \in A - F''\alpha]$ and by Theorem 7.46 A is a proper class. Since by hypothesis A is a set we conclude that $(\exists \alpha) [A - F''\alpha = 0]$. There is then a smallest such α , i.e.,

$$(\exists \alpha) [A - F''\alpha = 0 \wedge (\forall \beta < \alpha) [A - F''\beta \neq 0]].$$

If $x \in F''\alpha$ then $(\exists \beta < \alpha) [x = F'\beta]$. But $\beta < \alpha \rightarrow [A - F''\beta] \neq 0$ and hence $F'\beta \in [A - F''\beta]$, i.e., $x \in A$. Thus $F''\alpha \subseteq A$. But since $[A - F''\alpha] = 0$, $A \subseteq F''\alpha$. Therefore $A = F''\alpha$.

Since F is a function $F \Gamma \alpha$ is single valued. Furthermore

$$\gamma < \beta < \alpha \rightarrow F'\gamma \in F''\beta \wedge F'\beta \in A - F''\beta$$

i.e., $F'\gamma \in F''\beta \wedge F'\beta \notin F''\beta$. Then $F'\beta \neq F'\gamma$. Thus

$$\gamma < \alpha \wedge \beta < \alpha \wedge F'\gamma = F'\beta \rightarrow \gamma = \beta$$

that is $F \Gamma \alpha$ is one-to-one.

Remark. In Theorem 7.46 we see that the requirement that $(\forall \alpha) [G'(F \uparrow \alpha) \in A - F''\alpha]$ assures us that the function F defined by transfinite induction will be one-to-one. Conversely if F is one-to-one then $\mathcal{W}(F)$ will be a proper class and $(\forall \alpha) [G'(F \uparrow \alpha) \in A - F''\alpha]$. Furthermore if $\mathcal{W}(F)$ is a set then F cannot be one-to-one. In this case Theorem 7.47 assures us that if F fulfills the requirements for one-to-oneness "as long as it can" i.e., until $\mathcal{W}(F)$ is exhausted then the restriction of F to some ordinal α will map α one-to-one onto $\mathcal{W}(F)$. From this we can prove that every well ordered set is order isomorphic to an ordinal number.

Theorem 7.48. *If R is well founded on A and well orders A , and if $G = \{\langle x, y \rangle \mid y \in [A - \mathcal{W}(x)] \wedge [(A - \mathcal{W}(x)) \cap R^{-1}\{y\}] = 0\} \wedge F \mathcal{F}_m \text{ On}$
 $\wedge (\forall \alpha) [F'\alpha = G(F \uparrow \alpha)]$ then G is single valued and*

$$G'x \in [A - \mathcal{W}(x)] \longleftrightarrow [A - \mathcal{W}(x) \neq 0].$$

Proof. If $\langle x, y_1 \rangle, \langle x, y_2 \rangle \in G$ then

$$y_1 \in [A - \mathcal{W}(x)] \wedge y_2 \in [A - \mathcal{W}(x)]$$

and $[(A - \mathcal{W}(x)) \cap R^{-1}\{y_1\}] = 0 \wedge [(A - \mathcal{W}(x)) \cap R^{-1}\{y_2\}] = 0$. Therefore $y_1 \notin R^{-1}\{y_2\} \wedge y_2 \notin R^{-1}\{y_1\}$. Since R We A we must have $y_1 = y_2$.

If $G'x \in [A - \mathcal{W}(x)]$ then clearly $[A - \mathcal{W}(x)] \neq 0$. Conversely if $[A - \mathcal{W}(x)] \neq 0$ then since R is well founded on A and well orders A , $A - \mathcal{W}(x)$ has an R -minimal element, y . But $G'x = y$, i.e., $G'x \in [A - \mathcal{W}(x)]$.

Theorem 7.49. *If A is a proper class that is well ordered by R , if R is well founded on A and if*

$$G = \{\langle x, y \rangle \mid y \in [A - \mathcal{W}(x)] \wedge [(A - \mathcal{W}(x)) \cap R^{-1}\{y\}] = 0\} \wedge F \mathcal{F}_m \text{ On} \\ \wedge (\forall \alpha) [F'\alpha = G(F \uparrow \alpha)]$$

then $F \text{ Isom}_{E,R}(\text{On}, A)$.

Proof. If $(\exists \alpha) [A - F''\alpha = 0]$ then $A \subseteq F''\alpha$. Since $F''\alpha$ is a set it would then follow that A is a set. Since A is a proper class it follows that $(\forall \alpha) [A - F''\alpha \neq 0]$. From Theorem 7.48 and the defining properties of F and G it follows that $F'\alpha$ is the R -minimal element in $A - F''\alpha$ i.e.,

$$F'\alpha \in [A - F''\alpha] \quad \text{and} \quad [(A - F''\alpha) \cap R^{-1}\{F'\alpha\}] \neq 0.$$

From Theorem 7.46 it then follows that $\mathcal{W}(F) \subseteq A$ and F is one-to-one. To prove that F is onto we note that if $y \in \mathcal{W}(F)$ then $(\exists \alpha) [y = F'\alpha]$. Furthermore since $F'\alpha$ is the R -minimal element in $A - F''\alpha$

$$x R y \rightarrow x \notin [A - F''\alpha]$$

then

$$x \in A \wedge x R y \rightarrow x \in F''\alpha \\ \rightarrow x \in \mathcal{W}(F).$$

We then have fulfilled the hypotheses of Theorem 7.45:

$$R \text{ We } A \wedge \mathcal{W}(F) \subseteq A \wedge (\forall x \in A) (\forall y \in \mathcal{W}(F)) [x R y \rightarrow x \in \mathcal{W}(F)].$$

We conclude that $\mathcal{W}(F) = A \vee (\exists x \in A) [\mathcal{W}(F) = A \cap R^{-1}\{x\}]$. But $\mathcal{W}(F)$ cannot be an R-initial segment of A because R-initial segments of A are sets, and $\mathcal{W}(F)$ being the one-to-one image of the proper class On cannot be a set. Therefore $\mathcal{W}(F) = A$ and

$$F : \text{On} \xrightarrow[\text{onto}]{1-1} A.$$

Finally $\alpha < \beta \rightarrow F''\alpha \subseteq F''\beta$ and hence $[A - F''\beta] \subseteq [A - F''\alpha]$. Since $F'\beta \in [A - F''\beta]$ it follows that $F'\beta \in [A - F''\alpha]$. But $F'\alpha$ is the R-minimal element of $A - F''\alpha$. Hence

$$F'\alpha R F'\beta \vee F'\alpha = F'\beta.$$

Since F is one-to-one $F'\alpha \neq F'\beta$. Then

$$\alpha < \beta \rightarrow F'\alpha R F'\beta$$

i.e., $F \text{ Isom}_{E,R}(\text{On}, A)$.

Corollary 7.50. *If A is a proper class of ordinals and if*

$$G = \{\langle x, y \rangle \mid y \in [A - \mathcal{W}(x)] \wedge [(A - \mathcal{W}(x)) \cap R^{-1}\{y\}] = 0\} \wedge \\ F \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [F'\alpha = G(F \upharpoonright \alpha)]$$

then $F \text{ Isom}_{E,E}(\text{On}, A)$.

Proof. $E \text{ We } A \wedge E \text{ Wfr } A$.

Theorem 7.51. $R \text{ We } A \wedge \mathcal{M}(A) \rightarrow (\exists! \alpha) (\exists! f) [f \text{ Isom}_{E,R}(\alpha, A)]$.

Proof. If $G = \{\langle x, y \rangle \mid y \in [A - \mathcal{W}(x)] \wedge [(A \cap \mathcal{W}(x)) \cap R^{-1}\{y\}] = 0\}$ and if $F \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [F'\alpha = G(F \upharpoonright \alpha)]$ then by Theorem 7.47, $(\exists \alpha) [F''\alpha = A \wedge \mathcal{U}_2(F \upharpoonright \alpha)]$, i.e., $(F \upharpoonright \alpha) : \alpha \xrightarrow[\text{onto}]{1-1} A$.

That $F \upharpoonright \alpha$ is order preserving is proved as in the proof of Theorem 7.49 and is left to the reader. Then $(F \upharpoonright \alpha) \text{ Isom}_{E,R}(\alpha, A)$. But $F \upharpoonright \alpha$ is a set, i.e., $(\exists f) [f = F \upharpoonright \alpha]$. Then $(\exists f) [f \text{ Isom}_{E,R}(\alpha, A)]$.

The uniqueness argument is left to the reader.

Corollary 7.52. $A \subseteq \text{On} \wedge \mathcal{M}(A) \rightarrow (\exists! \alpha) (\exists! f) [f \text{ Isom}_{E,E}(\alpha, A)]$.

Proof. $A \subseteq \text{On} \rightarrow E \text{ We } A$.

Remark. Since E is a well founded relation and well foundedness is preserved under order isomorphism it follows that the requirement in Theorem 7.49 that R be well founded on A cannot be removed. In its absence we can only prove that On is order isomorphic to some R-initial segment of A . That this can occur we show by an example, the so called lexicographical ordering on $\text{On} \times \text{On}$.

Definition 7.53. $Le \triangleq \{\langle \langle \alpha, \beta \rangle, \langle \gamma, \delta \rangle \rangle \mid \alpha < \gamma \vee [\alpha = \gamma \wedge \beta < \delta]\}$.

Theorem 7.54.

- 1) $Le \text{ We } On^2 \wedge [B \subseteq On^2 \wedge B \neq 0 \rightarrow (\exists x \in B) [B \cap Le^{-1}\{x\} = 0]]$.
- 2) $\neg Le \text{ Wfr } On^2$.

Proof.

1) The proof is left to the reader.

2) If $F'\alpha = \langle 0, \alpha \rangle$ then $F : On \xrightarrow[\text{onto}]{1-1} Le^{-1}\{\langle 1, 0 \rangle\}$, hence $Le^{-1}\{\langle 1, 0 \rangle\}$ is a proper class.

Remark. From the lexicographical ordering we in turn define a relation R_0 that will be of value to us in later sections. We will show that this relation R_0 not only well orders On^2 it is well founded on On^2 .

Definition 7.55. $R_0 \triangleq \{\langle \langle \alpha, \beta \rangle, \langle \gamma, \delta \rangle \rangle \mid \max(\alpha, \beta) < \max(\gamma, \delta) \vee [\max(\alpha, \beta) = \max(\gamma, \delta) \wedge \langle \alpha, \beta \rangle Le \langle \gamma, \delta \rangle]\}$.

Theorem 7.56.

- 1) $R_0 \text{ We } On^2 \wedge [B \subseteq On^2 \wedge B \neq 0 \rightarrow (\exists x \in B) [B \cap R_0^{-1}\{x\} = 0]]$.
- 2) $R_0 \text{ Wfr } On^2$.

Proof.

1) The proof is left to the reader.

2) If $\gamma = \max(\alpha, \beta) + 1$,

$$\langle \delta, \tau \rangle R_0 \langle \alpha, \beta \rangle \rightarrow \max(\delta, \tau) \leq \max(\alpha, \beta) < \gamma.$$

Then $\langle \delta, \tau \rangle \in \gamma \times \gamma$ and hence $On^2 \cap R_0^{-1}\{\langle \alpha, \beta \rangle\} \subseteq \gamma \times \gamma$. Since $\gamma \times \gamma$ is a set $On^2 \cap R_0^{-1}\{\langle \alpha, \beta \rangle\}$ is a set. Thus R_0 is well founded on On^2 .

Remark. R_0 well orders On^2 and is well founded on On^2 . Consequently the ordering (R_0, On^2) is order isomorphic to (E, On) . By Theorem 7.49 there exists an order isomorphism, indeed a unique order isomorphism between the two order systems.

Definition 7.57. $J_0 \text{ Isom}_{R_0, E}(On^2, On)$.

8 Ordinal Arithmetic

In Section 7 we defined $\alpha + 1$ to be $\alpha \cup \{\alpha\}$. We proved that $\alpha + 1$ is an ordinal, that is, $\alpha + 1$ is a transitive set that is well ordered by the \in -relation. As a well ordered set $\alpha + 1$ has an initial segment α and its “terminal” segment beginning with α consists of just a single element, namely α .

If we add 1 to $\alpha + 1$ we obtain an ordinal with an initial segment α and a terminal segment, beginning with α , consisting of two elements α and $\alpha + 1$. Since this terminal segment $\{\alpha, \alpha + 1\}$ is order isomorphic to $2 \triangleq 1 + 1$ we call the sum of $\alpha + 1$ and 1, $\alpha + 2$.

In general by $\alpha + \beta$ we mean an ordinal obtained from α by adding 1, β times. That is, $\alpha + \beta$ is an ordinal with an initial segment α and a terminal segment, beginning with α , that is order isomorphic to β . That such an ordinal number exists is clear from the fact that $(\{0\} \times \alpha) \cup (\{1\} \times \beta)$ is well ordered by the lexicographical ordering Le . With respect to Le $\{0\} \times \alpha$ is an initial segment order isomorphic to α and $\{1\} \times \beta$ is a terminal segment order isomorphic to β .

It would then seem reasonable to define $\alpha + \beta$ as the ordinal that is order isomorphic to $(\{0\} \times \alpha) \cup (\{1\} \times \beta)$. However for certain purposes it is preferable to define $\alpha + \beta$ recursively in the following way.

Definition 8.1.

$$\begin{aligned}\alpha + 0 &= \alpha, \\ \alpha + (\beta + 1) &= (\alpha + \beta) + 1, \\ \alpha + \beta &= \bigcup_{\gamma < \beta} (\alpha + \gamma), \beta \in K_{II}.\end{aligned}$$

Remark. Definition 8.1 is an example of a very convenient form of definition by transfinite recursion. To define the addition of β to α i.e. $\alpha + \beta$ we specify the result of adding 0 to α , we define the sum of α and $(\beta + 1)$ as an operation on $\alpha + \beta$ namely the operation of adding one, and we define $\alpha + \beta$ for $\beta \in K_{II}$ as the supremum of the set of sums $\alpha + \gamma$ for $\gamma < \beta$.

That this is sufficient to define $\alpha + \beta$ for all α and β is clear from Theorem 7.42. If in Theorem 7.42 $H = \{\langle \alpha, \alpha + 1 \rangle \mid \alpha \in \text{On}\}$ and $a = \alpha$

then

$$\begin{aligned} F'0 &= \alpha, \\ F'(\beta + 1) &= H'F'\beta, \\ F'\beta &= \bigcup_{\gamma < \beta} F'\gamma, \beta \in K_{II}, \end{aligned}$$

i.e. $\alpha + \beta = F'\beta$.

Theorem 8.2. $\alpha + \beta \in \text{On}$.

Proof (by transfinite induction). $\alpha + 0 = \alpha \in \text{On}$. If $\alpha + \beta \in \text{On}$ then $\alpha + (\beta + 1) = (\alpha + \beta) + 1 \in \text{On}$. If $\beta \in K_{II}$ and $(\forall \gamma) [\gamma < \beta \rightarrow \alpha + \gamma \in \text{On}]$ then $\alpha + \beta = \bigcup_{\gamma < \beta} (\alpha + \gamma) \in \text{On}$.

Theorem 8.3. $0 + \alpha = \alpha + 0 = \alpha$.

Proof. $\alpha + 0 = \alpha$ by definition. If $0 + \alpha = \alpha$ then $0 + (\alpha + 1) = (0 + \alpha) + 1 = \alpha + 1$. If $\alpha \in K_{II}$ and $(\forall \beta) [\beta < \alpha \rightarrow 0 + \beta = \beta]$ then $0 + \alpha = \bigcup_{\beta < \alpha} (0 + \beta) = \bigcup_{\beta < \alpha} \beta = \alpha$.

Theorem 8.4. $\alpha < \beta \rightarrow \gamma + \alpha < \gamma + \beta$.

Proof (by transfinite induction on β). $\gamma + \alpha < (\gamma + \alpha) + 1 = \gamma + (\alpha + 1)$. If $\alpha < \beta \rightarrow \gamma + \alpha < \gamma + \beta$ and if $\alpha < \beta + 1$ then $\alpha < \beta \vee \alpha = \beta$. In either case we have that $\gamma + \alpha \leq \gamma + \beta < \gamma + \beta + 1$. If $\beta \in K_{II}$ and $(\forall \delta) [\delta < \beta \wedge \alpha < \delta \rightarrow \gamma + \alpha < \gamma + \delta]$ then $\gamma + \alpha < \bigcup_{\delta < \beta} (\gamma + \delta) = \gamma + \beta$.

Corollary 8.5. $\gamma + \alpha = \gamma + \beta \leftrightarrow \alpha = \beta$.

Proof.

$\alpha = \beta \rightarrow \gamma + \alpha = \gamma + \beta$; $\alpha < \beta \rightarrow \gamma + \alpha < \gamma + \beta$; $\beta < \alpha \rightarrow \gamma + \beta < \gamma + \alpha$.

Remark. For the proofs of several results on ordinal arithmetic we need the following property of suprema.

Theorem 8.6. $(\forall \alpha \in A) (\exists \beta \in B) [\alpha \leq \beta] \rightarrow \sup(A) \leq \sup(B)$.

Proof. $\gamma \in \sup(A) \rightarrow (\exists \alpha) [\gamma \in \alpha \wedge \alpha \in A]$. But $\alpha \in A \rightarrow (\exists \beta) [\beta \in B \wedge \alpha \leq \beta]$ i.e. $(\exists \beta) [\gamma \in \beta \wedge \beta \in B]$. Therefore $\gamma \in \sup(B)$ and hence $\sup(A) \leq \sup(B)$.

Remark. That Theorem 8.6 can not be an iff result is established by the counter-example

$$\sup(\omega) = \sup(\omega + 1).$$

Theorem 8.7. $\alpha \leq \beta \rightarrow \alpha + \gamma \leq \beta + \gamma$.

Proof (by transfinite induction on γ). $\alpha \leq \beta \rightarrow \alpha + 0 \leq \beta + 0$. If $\alpha + \gamma \leq \beta + \gamma$ then $\alpha + (\gamma + 1) \leq \beta + (\gamma + 1)$. If $\gamma \in K_{II}$ and $(\forall \delta) [\delta < \gamma \rightarrow \alpha + \delta \leq \beta + \delta]$ then $\alpha + \gamma = \bigcup_{\delta < \gamma} (\alpha + \delta) \leq \bigcup_{\delta < \gamma} (\beta + \delta) = \beta + \gamma$.

Theorem 8.8. $\alpha \leq \beta \rightarrow (\exists! \gamma) [\alpha + \gamma = \beta]$.

Proof. Since $\alpha \geq 0$ it follows from Theorems 8.7 and 8.3 that $\alpha + \beta \geq 0 + \beta = \beta$. Thus there exists a smallest ordinal γ such that $\alpha + \gamma \geq \beta$. If $\gamma \in K_I$ then $\gamma = 0 \vee (\exists \delta) [\gamma = \delta + 1]$. If $\gamma = 0$ then $\alpha \geq \beta \wedge \alpha \leq \beta$. Therefore $\alpha = \beta$ and $\alpha + \gamma = \beta$. If $\gamma = \delta + 1$ then $\delta < \gamma$ and $\alpha + \delta < \beta$. Then $\alpha + \delta + 1 \leq \beta$ i.e. $\alpha + \gamma \leq \beta$. But $\alpha + \gamma \geq \beta$; therefore $\alpha + \gamma = \beta$. If $\gamma \in K_{II}$ then $(\forall \delta) [\delta < \gamma \rightarrow \alpha + \delta < \beta]$. Therefore $\alpha + \gamma = \bigcup_{\delta < \beta} (\alpha + \delta) \leq \beta$. Again since $\alpha + \gamma \geq \beta$ we have that $\alpha + \gamma = \beta$.

From Corollary 8.5 we see that if $\alpha + \gamma = \beta$ and $\alpha + \delta = \beta$ then $\gamma = \delta$.

Theorem 8.9. $m + n \in \omega$.

Proof (by finite induction). $m + 0 = m \wedge m \in \omega$. If $m + n \in \omega$ then $m + (n + 1) = (m + n) + 1 \wedge (m + n) + 1 \in \omega$.

Theorem 8.10. $n < \omega \wedge \omega \leq \alpha \rightarrow n + \alpha = \alpha$.

Proof (by transfinite induction on α). $n + \omega = \bigcup_{\gamma < \omega} (n + \gamma)$. By Theorem 8.9, $\gamma < \omega \rightarrow n + \gamma < \omega$. Hence $\bigcup_{\gamma < \omega} (n + \gamma) \leq \omega$. On the other hand, by Theorem 8.9, $(\forall \gamma < \omega) (\exists \beta \in \omega) [n + \gamma \leq \beta]$. Then $\omega \leq \bigcup_{\gamma < \omega} (n + \gamma)$ by Theorem 8.6. Thus $n + \omega = \omega$. By Definition 8.1 and the induction hypothesis

$$n + (\alpha + 1) = (n + \alpha) + 1 = \alpha + 1.$$

Finally, if $\alpha \in K_{II}$ then from the induction hypothesis

$$n + \alpha = \bigcup_{\beta < \alpha} (n + \beta) = \bigcup_{\beta < \alpha} \beta = \alpha.$$

Remark. From Theorem 8.10 we see that ordinal addition is not commutative:

$$1 + \omega = \omega \neq \omega + 1.$$

Furthermore $1 + \omega = 2 + \omega$ but $1 \neq 2$. Thus we do *not* have a right hand cancellation law. From Corollary 8.5 we see that we do however have a left hand cancellation law.

Theorem 8.4 assures us of the additivity property for inequalities for addition from the left. Theorem 8.6 however suggests that addition from the right may not preserve strict inequality. This is the case as we see from the following example.

$$1 < 2 \quad \text{but} \quad 1 + \omega = 2 + \omega.$$

Theorem 8.8 shows that subtraction, when permitted, is unique.

Finally ordinal addition is associative. The proof requires

Theorem 8.11. $\beta \in K_{II} \rightarrow \alpha + \beta \in K_{II}$.

Proof. $\beta \in K_{II} \rightarrow \beta \neq 0$. Therefore $\alpha + \beta \neq 0$. Thus $\alpha + \beta \in K_{II}$ or $(\exists \delta) [\alpha + \beta = \delta + 1]$. But $\beta \in K_{II} \rightarrow \alpha + \beta = \bigcup_{\gamma < \beta} (\alpha + \gamma)$. Since $\delta \in \delta + 1$, $\alpha + \beta = \delta + 1 \rightarrow \delta \in \bigcup_{\gamma < \beta} (\alpha + \gamma)$ i.e. $(\exists \gamma) [\gamma < \beta \wedge \delta \in \alpha + \gamma]$. But $\delta \in \alpha + \gamma \rightarrow (\delta + 1) \in (\alpha + \gamma + 1)$. Since $\beta \in K_{II}$, $\gamma < \beta \rightarrow \gamma + 1 < \beta$. Therefore $\delta + 1 \in \bigcup_{\gamma < \beta} (\alpha + \gamma)$ i.e. $\delta + 1 \in \delta + 1$.

Since this is a contradiction we conclude that $\alpha + \beta \in K_{II}$.

Theorem 8.12. $(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma)$.

Proof (by transfinite induction on γ). $(\alpha + \beta) + 0 = \alpha + \beta = \alpha + (\beta + 0)$. If $(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma)$ then $(\alpha + \beta) + (\gamma + 1) = ((\alpha + \beta) + \gamma) + 1 = (\alpha + (\beta + \gamma)) + 1 = \alpha + ((\beta + \gamma) + 1) = \alpha + (\beta + (\gamma + 1))$. If $\gamma \in K_{II}$ and $(\alpha + \beta) + \delta = \alpha + (\beta + \delta)$ for $\delta < \gamma$ then

$$(\alpha + \beta) + \gamma = \bigcup_{\delta < \gamma} ((\alpha + \beta) + \delta) = \bigcup_{\delta < \gamma} (\alpha + (\beta + \delta)).$$

Furthermore since $\gamma \in K_{II}$ we have by Theorem 8.11 that $\beta + \gamma \in K_{II}$. Therefore

$$\alpha + (\beta + \gamma) = \bigcup_{\eta < \beta + \gamma} (\alpha + \eta).$$

If $\delta < \gamma$ and $\eta = \beta + \delta$ then $\eta < \beta + \gamma$ and $\alpha + (\beta + \delta) \leq \alpha + \eta$. Conversely if $\eta < \beta + \gamma$ then $\eta < \beta \vee (\exists \delta) [\eta = \beta + \delta]$. If $\eta < \beta$ then $\alpha + \eta \leq \alpha + (\beta + 0) \wedge 0 < \gamma$. If $\eta = \beta + \delta$ then $\alpha + \eta \leq \alpha + (\beta + \delta)$ and since $\eta < \beta + \gamma$ we have that $\delta < \gamma$. Thus by Theorem 8.6

$$\bigcup_{\delta < \gamma} (\alpha + (\beta + \delta)) = \bigcup_{\eta < \beta + \gamma} (\alpha + \eta)$$

i.e.

$$(\alpha + \beta) + \gamma = \alpha + (\beta + \gamma).$$

Theorem 8.13. $\alpha \geq \omega \rightarrow (\exists! \beta) (\exists! n) [\beta \in K_{II} \wedge \alpha = \beta + n]$.

Proof. If $A = \{\gamma \in K_{II} \mid \gamma \leq \alpha\}$ and if $\beta = \cup(A)$ then $\beta \in K_{II}$ and $\beta < \alpha$. Therefore, by Theorem 8.8, $(\exists \gamma) [\beta + \gamma = \alpha]$. If $\gamma \geq \omega$ then $(\exists \delta) [\gamma = \omega + \delta]$ and $\alpha = \beta + (\omega + \delta) = (\beta + \omega) + \delta$. But $\beta + \omega \in K_{II}$ and $\beta + \omega < \alpha$. Thus $\beta + \omega \in A$; but $\beta < \beta + \omega$. This contradicts the definition of β ; hence $\gamma < \omega$.

If $\alpha = \beta_1 + n_1 = \beta_2 + n_2$ with $\beta_1 \leq \beta_2$ then $(\exists \gamma) [\beta_1 + \gamma = \beta_2]$ i.e.

$$\beta_1 + n_1 = \beta_1 + \gamma + n_2,$$

$$n_1 = \gamma + n_2.$$

Since $\gamma \leq \gamma + n_2$ we must have that $\gamma < \omega$. Furthermore since $\beta_1 + \gamma = \beta_2$ and $\beta_2 \in K_{II}$ it follows that $\gamma = 0$ i.e. $\beta_1 = \beta_2$ and $n_1 = n_2$.

Definition 8.14. $\alpha \div \beta = \cap \{\gamma \mid \beta + \gamma \geq \alpha\}$.

Exercises. Prove the following.

- 1) $\alpha + \beta \in \omega \rightarrow \alpha \in \omega \wedge \beta \in \omega.$
- 2) $\alpha \leq \beta \rightarrow \alpha + (\beta - \alpha) = \beta.$
- 3) $\omega - n = \omega.$
- 4) $[m + n = n + m] \wedge [m + n = k + n \rightarrow m = k].$
- 5) $\alpha \leq \alpha + \beta \wedge [\beta > 0 \rightarrow \alpha < \alpha + \beta].$
- 6) $\alpha \leq \beta + \alpha.$
- 7) $\alpha + \beta \in K_{II} \leftrightarrow \beta \in K_{II} \vee [\beta = 0 \wedge \alpha \in K_{II}].$
- 8) $\beta \in K_{II} \wedge \alpha < \beta \rightarrow (\forall n) [\alpha + n < \beta].$
- 9) $\alpha + \beta$ is order isomorphic to $(\{0\} \times \alpha) \cup (\{1\} \times \beta)$ where the order on the later set is Le i.e. $(\exists f) [f \text{ Isom}_{E, Le}(\alpha + \beta, (\{0\} \times \alpha) \cup (\{1\} \times \beta))].$
- 10) Prove Theorem 8.8 by transfinite induction on $\beta.$

Remark. From the foregoing we see that ordinal addition on ω has all of the arithmetic properties that we expect. Addition on On is however not commutative and the right hand cancellation law fails.

In very much the same way as we define integer multiplication as repeated addition we can also define ordinal multiplication as repeated addition. For the justification of our definition we again appeal to Theorem 7.42.

If in Theorem 7.42, $H_\alpha = \{\langle \beta, \beta + \alpha \rangle \mid \beta \in \text{On}\}$ and if $a = 0$ then

$$\begin{aligned} F'_\alpha 0 &= 0, \\ F'_\alpha(\beta + 1) &= H'_\alpha F'_\alpha \beta, \\ F'_\alpha \beta &= \bigcup_{\gamma < \beta} F'_\alpha \gamma, \beta \in K_{II}. \end{aligned}$$

We define the product of α and β , i.e. $\alpha\beta$, as $F'_\alpha \beta.$

Definition 8.15.

$$\begin{aligned} \alpha \cdot 0 &= 0, \\ \alpha(\beta + 1) &= \alpha\beta + \alpha, \\ \alpha\beta &= \bigcup_{\gamma < \beta} \alpha\gamma, \beta \in K_{II}. \end{aligned}$$

Theorem 8.16. $\alpha\beta \in \text{On}.$

Proof (by transfinite induction). $\alpha \cdot 0 = 0 \in \text{On}.$ If $\alpha\beta \in \text{On}$ then $\alpha(\beta + 1) = \alpha\beta + \alpha \in \text{On}.$ If $\beta \in K_{II}$ and $\alpha\gamma \in \text{On}$ for $\gamma < \beta$ then $\bigcup_{\delta < \beta} \alpha\delta \in \text{On}.$

Theorem 8.17. $mn \in \omega.$

Proof (by finite induction). $m \cdot 0 = 0 \in \omega.$ If $mn \in \omega$ then $m(n + 1) = mn + n \in \omega.$

Theorem 8.18.

1) $0 \cdot \alpha = \alpha \cdot 0 = 0.$

2) $1 \cdot \alpha = \alpha \cdot 1 = \alpha.$

Proof (by transfinite induction).

1) $0 \cdot 0 = 0.$ If $0 \cdot \alpha = 0$ then $0(\alpha + 1) = 0 \cdot \alpha + 0 = 0.$ If $\alpha \in K_{II}$ and $0 \cdot \gamma = 0$ for $\gamma < \alpha$ then $0 \cdot \alpha = \bigcup_{\gamma < \alpha} 0 \cdot \gamma = 0.$ From Definition 8.15 we have that $\alpha \cdot 0 = 0.$

2) $1 \cdot 0 = 0.$ If $1 \cdot \alpha = \alpha$ then $1(\alpha + 1) = 1 \cdot \alpha + 1 = \alpha + 1.$ If $\alpha \in K_{II}$ and $1 \cdot \gamma = \gamma$ for $\gamma < \alpha$ then $1 \cdot \alpha = \bigcup_{\gamma < \alpha} 1 \cdot \gamma = \alpha.$ From 1) above $\alpha \cdot 1 = \alpha(0 + 1) = \alpha \cdot 0 + \alpha = 0 + \alpha = \alpha.$

Theorem 8.19. $\alpha < \beta \wedge \gamma > 0 \leftrightarrow \gamma\alpha < \gamma\beta.$

Proof. (by transfinite induction on β). $\gamma > 0 \rightarrow \gamma\alpha < \gamma\alpha + \gamma = \gamma(\alpha + 1).$ If $\gamma > 0 \wedge \alpha < \beta \rightarrow \gamma\alpha < \gamma\beta$ and if $\alpha < \beta + 1$ then $\alpha < \beta \vee \alpha = \beta.$ In either case $\gamma\alpha \leq \gamma\beta < \gamma\beta + \gamma = \gamma(\beta + 1).$ If $\beta \in K_{II}$ and if $(\forall \delta) [\delta < \beta \wedge \alpha < \delta \rightarrow \gamma\alpha < \gamma\delta]$ then $\gamma\alpha < \bigcup_{\delta < \beta} \gamma\delta = \gamma\beta.$

Conversely $\gamma\alpha < \gamma\beta \rightarrow \gamma > 0.$ Since $\alpha = \beta \rightarrow \gamma\alpha = \gamma\beta$ and $\beta < \alpha \wedge \gamma > 0 \rightarrow \gamma\beta < \gamma\alpha$ we conclude that $\gamma\alpha < \gamma\beta \rightarrow \alpha < \beta \wedge \gamma > 0.$

Theorem 8.20. $\gamma\alpha = \gamma\beta \wedge \gamma > 0 \rightarrow \alpha = \beta.$

Proof. $\alpha < \beta \wedge \gamma > 0 \rightarrow \gamma\alpha < \gamma\beta;$ $\beta < \alpha \wedge \gamma > 0 \rightarrow \gamma\beta < \gamma\alpha.$

Theorem 8.21. $\alpha \leq \beta \rightarrow \alpha\gamma \leq \beta\gamma.$

Proof (by transfinite induction). $\alpha \cdot 0 = 0 \leq \beta \cdot 0.$ If $\alpha\gamma \leq \beta\gamma$ then $\alpha(\gamma + 1) = \alpha\gamma + \alpha \leq \beta\gamma + \beta = \beta(\gamma + 1).$ If $\gamma \in K_{II}$ and $(\forall \delta) [\delta < \gamma \rightarrow \alpha\delta \leq \beta\delta]$ then $\alpha\gamma = \bigcup_{\delta < \gamma} \alpha\delta \leq \bigcup_{\delta < \gamma} \beta\delta = \beta\gamma.$

Theorem 8.22. $\alpha\beta = 0 \leftrightarrow \alpha = 0 \vee \beta = 0.$

Proof. $0 \cdot \beta = 0 \wedge \alpha \cdot 0 = 0.$ Furthermore $\alpha \geq 1 \wedge \beta \geq 1 \rightarrow \alpha \leq \alpha\beta$ i.e. $\alpha\beta \neq 0.$

Theorem 8.23. $\alpha \neq 0 \wedge \beta \in K_{II} \rightarrow \alpha\beta \in K_{II}.$

Proof. $\alpha \neq 0 \wedge \beta \in K_{II} \rightarrow \alpha\beta \neq 0.$ Therefore $\alpha\beta \in K_{II}$ or $(\exists \gamma) [\gamma + 1 = \alpha\beta].$ Since $\gamma \in \gamma + 1$ and since $\beta \in K_{II}$ it follows that if $\gamma + 1 = \alpha\beta$ then

$$\gamma \in \bigcup_{\delta < \beta} \alpha\delta$$

i.e. $(\exists \delta) [\delta < \beta \wedge \gamma < \alpha\delta]$ (see Theorem 8.24). Then $\gamma + 1 < \alpha\delta + 1 \leq \alpha\delta + \alpha = \alpha(\delta + 1).$ But $\beta \in K_{II}$ and $\delta < \beta$ implies $\delta + 1 < \beta$ i.e. $\gamma + 1 \in \alpha(\delta + 1)$ and

$\delta + 1 < \beta$. Thus

$$\gamma + 1 \in \bigcup_{\delta < \beta} \alpha \delta = \gamma + 1.$$

From this contradiction we conclude that $\alpha \beta \in K_{II}$.

Theorem 8.24. $\beta \in K_{II} \wedge \gamma < \alpha \beta \rightarrow (\exists \delta) [\delta < \beta \wedge \gamma < \alpha \delta]$.

Proof. Definition 8.15.

Theorem 8.25. $\alpha(\beta + \gamma) = \alpha \beta + \alpha \gamma$.

Proof (by transfinite induction on γ). $\alpha(\beta + 0) = \alpha \beta = \alpha \beta + \alpha \cdot 0$. If $\alpha(\beta + \gamma) = \alpha \beta + \alpha \gamma$, then $\alpha(\beta + (\gamma + 1)) = \alpha((\beta + \gamma) + 1) = \alpha(\beta + \gamma) + \alpha = (\alpha \beta + \alpha \gamma) + \alpha = \alpha \beta + (\alpha \gamma + \alpha) = \alpha \beta + \alpha(\gamma + 1)$. If $\gamma \in K_{II}$ and $\alpha(\beta + \delta) = \alpha \beta + \alpha \delta$ for $\delta < \gamma$ then we consider two cases $\alpha = 0 \wedge \alpha \neq 0$. If $\alpha = 0$ then

$$\alpha(\beta + \gamma) = 0 = \alpha \beta + \alpha \gamma.$$

If $\alpha \neq 0$ then since $\gamma \in K_{II}$ it follows that $\beta + \gamma \in K_{II}$ and $\alpha \gamma \in K_{II}$.

$$\begin{aligned} \alpha(\beta + \gamma) &= \bigcup_{\delta < \beta + \gamma} \alpha \delta, \\ \alpha \beta + \alpha \gamma &= \bigcup_{\eta < \alpha \gamma} (\alpha \beta + \eta). \end{aligned}$$

If $\delta < \beta + \gamma$ then $\delta < \beta \vee (\exists \tau) [\tau < \gamma \wedge \delta = \beta + \tau]$. Therefore $\alpha \delta < \alpha \beta$ or

$$\alpha \delta = \alpha(\beta + \tau) = \alpha \beta + \alpha \tau = \alpha \beta + \eta$$

where $\eta = \alpha \tau$. Since $\tau < \gamma$ we have that $\alpha \tau < \alpha \gamma$ i.e. $\eta < \alpha \gamma$. Thus

$$\alpha(\beta + \gamma) \leq \alpha \beta + \alpha \gamma.$$

If $\eta < \alpha \gamma$ then $(\exists \delta) [\delta < \gamma \wedge \eta < \alpha \delta]$. Therefore $\beta + \delta < \beta + \gamma$ and hence

$$\alpha \beta + \eta < \alpha \beta + \alpha \delta = \alpha(\beta + \delta).$$

Thus $\alpha \beta + \alpha \gamma \leq \alpha(\beta + \gamma)$ and hence $\alpha(\beta + \gamma) = \alpha \beta + \alpha \gamma$.

Remark. $(\omega + 1) 2 = (\omega + 1) + (\omega + 1) = \omega + (1 + \omega) + 1 = \omega + \omega + 1 = \omega 2 + 1 \neq \omega 2 + 2$. We do not have a right hand distributive law.

Theorem 8.26. $(\alpha \beta) \gamma = \alpha(\beta \gamma)$.

Proof (by induction on γ). $(\alpha \beta) \cdot 0 = 0 = \alpha \cdot 0 = \alpha(\beta \cdot 0)$. If $(\alpha \beta) \gamma = \alpha(\beta \gamma)$ then $(\alpha \beta) (\gamma + 1) = (\alpha \beta) \gamma + \alpha \beta = \alpha(\beta \gamma) + \alpha \beta = \alpha(\beta \gamma + \beta) = \alpha(\beta(\gamma + 1))$. If $\gamma \in K_{II}$ and $\alpha \beta = 0$ then $\alpha = 0 \vee \beta = 0$ and $(\alpha \beta) \gamma = 0 = \alpha(\beta \gamma)$. If $\alpha \beta \neq 0$ then $\beta \gamma \in K_{II}$ and hence

$$\begin{aligned} (\alpha \beta) \gamma &= \bigcup_{\delta < \gamma} (\alpha \beta) \delta, \\ \alpha(\beta \gamma) &= \bigcup_{\eta < \beta \gamma} \alpha \eta. \end{aligned}$$

But $\delta < \gamma \iff \beta\delta < \beta\gamma$. Therefore $(\alpha\beta)\gamma = \alpha(\beta\gamma)$.

Theorem 8.27. $\beta \neq 0 \rightarrow (\exists! \gamma) (\exists! \delta) [\alpha = \beta\gamma + \delta \wedge \delta < \beta]$.

Proof. If $\alpha < \beta$ then $\alpha = \beta \cdot 0 + \alpha \wedge \alpha < \beta$. If $\beta \leq \alpha$ and $\gamma = \sup\{\delta \mid \beta\delta \leq \alpha\}$ then $\gamma \geq 1$. If $\alpha < \beta v$ then $\beta\delta \leq \alpha \rightarrow \delta < v$ thus $\gamma \leq v$. Consequently $\delta < \gamma \rightarrow \beta\delta \leq \alpha$. If $(\exists \tau) [\gamma = \tau + 1]$ then $\tau < \gamma$ hence $\tau \in \{\delta \mid \beta\delta \leq \alpha\}$ therefore $(\exists v) [v \in \{\delta \mid \beta\delta \leq \alpha\} \wedge \tau < v]$. Thus $v = \gamma$ i.e. $\beta\gamma \leq \alpha$. If $\gamma \in K_{\text{II}}$ then $\beta\gamma = \bigcup_{\delta < \gamma} \beta\delta \leq \bigcup_{\delta < \alpha} \delta = \alpha$. Thus $\beta\gamma \leq \alpha$ and hence $(\exists \delta) [\alpha = \beta\gamma + \delta]$. If $\delta \geq \beta$, then $(\exists \mu) [\delta = \beta + \mu]$ then $\alpha = \beta\gamma + \beta + \mu = \beta(\gamma + 1) + \mu$ hence $\beta(\gamma + 1) \leq \alpha$ and $\gamma + 1 \leq \gamma$. From this we conclude that $\delta < \beta$ i.e.

$$\alpha = \beta\gamma + \delta \wedge \delta < \beta.$$

If $\alpha = \beta\gamma_1 + \delta_1 = \beta\gamma_2 + \delta_2$ with $\delta_1 < \beta \wedge \delta_2 < \beta \wedge \gamma_1 \leq \gamma_2$ then $(\exists v) [\gamma_2 = \gamma_1 + v]$ and

$$\begin{aligned} \beta\gamma_1 + \delta_1 &= \beta(\gamma_1 + v) + \delta_2 = \beta\gamma_1 + \beta v + \delta_2 \\ \delta_1 &= \beta v + \delta_2. \end{aligned}$$

But $\beta v + \delta_2 < \beta$. Therefore $v = 0$ and hence $\delta_1 = \delta_2 \wedge \gamma_1 = \gamma_2$.

Corollary 8.28. $n \neq 0 \rightarrow (\exists! q) (\exists! r) [m = nq + r \wedge r < n]$.

Proof. By Theorem 8.27 $(\exists! \gamma) (\exists! \delta) [m = n\gamma + \delta \wedge \delta < n]$. But $n\gamma + \delta \in \omega \rightarrow n\gamma \in \omega$ and $\delta \in \omega$. Furthermore $1 \leq n \rightarrow \gamma \leq n\gamma$. Therefore $\gamma \in \omega$.

Theorem 8.29. $\gamma \in K_{\text{II}} \wedge m \neq 0 \rightarrow m(\gamma + n) = \gamma + mn$.

Proof (by induction on $\gamma + n$). $m\omega = \bigcup_{n < \omega} mn$. Since $mn < \omega$ we have

$$\bigcup_{n < \omega} mn \leq \omega.$$

Furthermore $p < \omega \rightarrow (\exists q) (\exists r) [p = mq + r \wedge r < m]$. But $p = mq + r \leq mq + m = m(q + 1)$. Therefore $p \in \bigcup_{n < \omega} mn$; hence

$$\bigcup_{n < \omega} mn = \omega.$$

If $m(\gamma + n) = \gamma + mn$ then $m(\gamma + n + 1) = m(\gamma + n) + m = (\gamma + mn) + m = \gamma + m(n + 1)$. If $\gamma > \omega$ then

$$m\gamma = \bigcup_{\delta < \gamma} m\delta.$$

If $\delta < \gamma$ then $\delta < \omega \vee \omega \leq \delta$. If $\delta < \omega$ then $m\delta < \omega < \gamma$.

If $\omega \leq \delta$ then $(\exists \beta) (\exists n) [\beta \in K_{\text{II}} \wedge \delta = \beta + n]$. Then from the induction hypothesis $m\delta = \beta + mn$. But $\beta \leq \delta < \gamma$ and $\gamma \in K_{\text{II}}$. Therefore $\beta + mn < \gamma$.

Since $(\forall \delta) [\delta < \gamma \rightarrow m\delta < \gamma]$ we conclude that

$$m\gamma \leq \gamma.$$

But $1 \leq m$ and hence $\gamma \leq m\gamma$. Therefore $m\gamma = \gamma$.

Exercises. Prove the following.

- 1) $\beta \in K_{II} \rightarrow [\gamma < \alpha\beta \leftrightarrow (\exists \delta) [\delta < \beta \wedge \gamma < \alpha\delta]].$
- 2) $\alpha\gamma \leq \beta\gamma \rightarrow \gamma = 0 \vee \alpha \leq \beta.$
- 3) $mn = nm \wedge (m + n)k = mk + nk.$
- 4) In Theorem 8.29 can m be replaced by α with the restriction that $\alpha < \gamma$?
- 5) $\alpha\beta \in K_{II} \leftrightarrow \alpha\beta \neq 0 \wedge [\beta \in K_{II} \vee \alpha \in K_{II}].$
- 6) $\alpha\beta$ is order isomorphic to $\alpha \times \beta$ well ordered lexicographically i.e. $(\exists f) [f \text{ Isom}_{E, L_e}(\alpha\beta, \alpha \times \beta)].$

Remark. When restricted to ω ordinal multiplication has the properties expected. On the class of all ordinals however multiplication is not commutative.

$$2 \cdot \omega = \omega \quad \text{and} \quad \omega \cdot 2 = \omega + \omega.$$

We do not have a right hand cancellation law:

$$1 \cdot \omega = 2 \cdot \omega \quad \text{but} \quad 1 \neq 2.$$

Having defined multiplication as repeated addition we next define exponentiation as repeated multiplication.

Definition 8.30.

$$\begin{aligned} \alpha^0 &= 1. \\ \alpha^{\beta+1} &= \alpha^\beta \cdot \alpha. \\ \alpha^\beta &= \bigcup_{\gamma < \beta} \alpha^\gamma, \beta \in K_{II} \wedge \alpha \neq 0. \\ \alpha^\beta &= 0, \beta \in K_{II} \wedge \alpha = 0. \end{aligned}$$

Theorem 8.31.

- 1) $0^0 = 1.$
- 2) $0^\beta = 0, \beta \geq 1.$
- 3) $1^\beta = 1.$

Proof.

- 1) From Definition 8.30, $0^0 = 1.$
- 2) $\beta \geq 1 \rightarrow \beta \in K_{II} \vee (\exists \delta) [\beta = \delta + 1].$ If $\beta \in K_{II}$ then by Definition 8.30, $0^\beta = 0.$ If $\beta = \delta + 1$ then $0^\beta = 0^{\delta+1} = 0^\delta \cdot 0 = 0.$

3) (By transfinite induction) $1^0 = 1$. If $1^\beta = 1$ then $1^{\beta+1} = 1^\beta \cdot 1 = 1^\beta = 1$. If $\beta \in K_{II}$ then $1^\beta = \bigcup_{\alpha < \beta} 1^\alpha = 1$.

Theorem 8.32. $1 \leq \alpha \rightarrow 1 \leq \alpha^\beta$.

Proof (by transfinite induction on β). $\alpha^0 = 1$. If $1 \leq \alpha^\beta$ then since $1 \leq \alpha$ we have $\alpha^\beta \leq \alpha^\beta \cdot \alpha$ i.e. $1 \leq \alpha^{\beta+1}$. If $\beta \in K_{II}$ then since $\alpha \neq 0$

$$\alpha^\beta = \bigcup_{\gamma < \beta} \alpha^\gamma.$$

Since $0 < \beta \wedge \alpha^0 = 1$ we have $1 \leq \bigcup_{\gamma < \beta} \alpha^\gamma = \alpha^\beta$.

Theorem 8.33. $\alpha < \beta \wedge 1 < \gamma \rightarrow \gamma^\alpha < \gamma^\beta$.

Proof (by transfinite induction on β). $1 < \gamma \rightarrow \gamma^\alpha < \gamma^{\alpha+1}$. If $\alpha < \beta \wedge 1 < \gamma \rightarrow \gamma^\alpha < \gamma^\beta$ then since $\alpha < \beta + 1 \rightarrow \alpha < \beta \vee \alpha = \beta$ we have that $\gamma^\alpha \leq \gamma^\beta < \gamma^{\beta+1}$. If $\beta \in K_{II}$ then since $\gamma \neq 0$

$$\gamma^\beta = \bigcup_{\delta < \beta} \gamma^\delta.$$

Then $\alpha < \beta \rightarrow \alpha + 1 < \beta$ and hence $\gamma^\alpha < \bigcup_{\delta < \beta} \gamma^\delta = \gamma^\beta$.

Corollary 8.34. $1 < \gamma \wedge \gamma^\alpha < \gamma^\beta \rightarrow \alpha < \beta$.

Proof. $\beta \leq \alpha \wedge 1 < \gamma \rightarrow \gamma^\beta \leq \gamma^\alpha$.

Theorem 8.35. $\alpha < \beta \rightarrow \alpha^\gamma \leq \beta^\gamma$.

Proof (by transfinite induction on γ). $\alpha^0 = 1 = \beta^0$. If $\alpha^\gamma \leq \beta^\gamma$ then $\alpha^{\gamma+1} = \alpha^\gamma \cdot \alpha \leq \beta^\gamma \cdot \alpha < \beta^\gamma \cdot \beta = \beta^{\gamma+1}$. If $\gamma \in K_{II}$ and if $\alpha^\delta \leq \beta^\delta$ for $\delta < \gamma$ then

$$\alpha^\gamma = \bigcup_{\delta < \gamma} \alpha^\delta \leq \bigcup_{\delta < \gamma} \beta^\delta = \beta^\gamma.$$

Corollary 8.36. $\alpha < \beta \wedge \gamma \in K_I \wedge \gamma \neq 0 \rightarrow \alpha^\gamma < \beta^\gamma$.

Proof. $\gamma \in K_I \wedge \gamma \neq 0 \rightarrow (\exists \delta) [\gamma = \delta + 1]$. Then $\alpha^\delta \leq \beta^\delta$. But $\alpha^\gamma = \alpha^\delta \cdot \alpha \leq \beta^\delta \cdot \alpha < \beta^\delta \cdot \beta = \beta^\gamma$.

Remark. That $\alpha < \beta \wedge \gamma \in K_{II}$ does not imply $\alpha^\gamma < \beta^\gamma$ follows from the observation that $2 < 3$ but

$$2^\omega = 3^\omega = \omega.$$

The proof is left to the reader.

Theorem 8.37. $\alpha > 1 \rightarrow \beta \leq \alpha^\beta$.

Proof (by transfinite induction on β). $0 \leq \alpha^0 = 1$. If $\beta \leq \alpha^\beta$ then $\beta + 1 \leq \alpha^\beta + 1$. But since $\beta < \beta + 1$ we have from Theorem 8.33 that

$\alpha^\beta < \alpha^{\beta+1}$ and hence $\alpha^\beta + 1 \leq \alpha^{\beta+1}$ i.e. $\beta + 1 \leq \alpha^{\beta+1}$. If $\beta \in K_{II}$ and $\gamma < \beta \rightarrow \gamma \leq \alpha^\gamma$ then

$$\beta \leq \bigcup_{\gamma < \beta} \alpha^\gamma = \alpha^\beta.$$

Theorem 8.38. $\alpha > 1 \wedge \beta > 0 \rightarrow (\exists! \delta) [\alpha^\delta \leq \beta < \alpha^{\delta+1}]$.

Proof. Since by Theorem 8.37 $\beta \leq \alpha^\beta$ and since $\alpha^\beta < \alpha^{\beta+1}$ there exists a smallest ordinal γ such that $\beta < \alpha^\gamma$. From Definition 8.30 it follows that $\gamma \in K_I$. Since $\alpha^0 = 1 \wedge \beta \geq 1$ it follows that $\gamma \neq 0$; therefore $(\exists \delta) [\gamma = \delta + 1]$. But $\delta < \delta + 1$ hence $\alpha^\delta \leq \beta < \alpha^{\delta+1}$.

If $\alpha^\delta \leq \beta < \alpha^{\delta+1}$ and $\alpha^\gamma \leq \beta < \alpha^{\gamma+1}$ then $\delta < \gamma \rightarrow \delta + 1 \leq \gamma$. Hence

$$\beta < \alpha^{\delta+1} \leq \alpha^\gamma \leq \beta.$$

Similarly $\gamma < \delta \rightarrow \beta < \beta$. Therefore $\delta = \gamma$.

Theorem 8.39.

$$1) \quad \alpha > 1 \wedge \beta \in K_{II} \rightarrow \alpha^\beta \in K_{II}.$$

$$2) \quad \alpha \in K_{II} \wedge \beta > 0 \rightarrow \alpha^\beta \in K_{II}.$$

Proof.

1) $\alpha > 1 \rightarrow \alpha^\beta \geq 1$ i.e. $\alpha^\beta \neq 0$. Therefore $\alpha^\beta \in K_{II}$ or $(\exists \delta) [\delta + 1 = \alpha^\beta]$. But since $\beta \in K_{II} \wedge \alpha \neq 0$

$$\alpha^\beta = \bigcup_{\gamma < \beta} \alpha^\gamma.$$

Therefore since $\delta \in \delta + 1 = \alpha^\beta$ it follows that $(\exists \gamma) [\delta \in \alpha^\gamma \wedge \gamma < \beta]$. Since $1 < \alpha, \alpha^\gamma < \alpha^{\gamma+1}$. Then $\delta < \alpha^\gamma$ hence $\delta + 1 \leq \alpha^\gamma < \alpha^{\gamma+1}$ and $\gamma + 1 < \beta$ i.e. $\delta + 1 \in \alpha^\beta = \delta + 1$. From this contradiction we conclude that

$$\alpha^\beta \in K_{II}.$$

2) If $\beta \in K_{II}$ then $\alpha^\beta \in K_{II}$ by 1) above. If $\beta \in K_I$ then since $\beta \neq 0$, $(\exists \delta) [\beta = \delta + 1]$. Then $\alpha^\beta = \alpha^{\delta+1} = \alpha^\delta \cdot \alpha$. Since $\alpha \in K_{II}$, $\alpha^\delta \neq 0$, therefore $\alpha^\delta \alpha \in K_{II}$.

Theorem 8.40. $\beta \in K_{II} \wedge \gamma < \alpha^\beta \rightarrow (\exists \delta) [\delta < \beta \wedge \gamma < \alpha^\delta]$.

Proof. Since $\gamma < \alpha^\beta$ we have that $\alpha \neq 0$. Therefore $\alpha^\beta = \bigcup_{\delta < \beta} \alpha^\delta$. Then $\gamma < \alpha^\beta \rightarrow (\exists \delta) [\delta < \beta \wedge \gamma < \alpha^\delta]$.

Theorem 8.41. $\alpha^\beta \cdot \alpha^\gamma = \alpha^{\beta+\gamma}$.

Proof (by transfinite induction on γ). $\alpha^\beta \cdot \alpha^0 = \alpha^\beta \cdot 1 = \alpha^\beta = \alpha^{\beta+0}$. If $\alpha^\beta \cdot \alpha^\gamma = \alpha^{\beta+\gamma}$ then $\alpha^\beta \cdot \alpha^{\gamma+1} = \alpha^\beta \cdot \alpha^\gamma \alpha = \alpha^{\beta+\gamma} \alpha = \alpha^{\beta+(\gamma+1)}$. If $\gamma \in K_{II}$ then $\beta + \gamma \in K_{II}$. If $\alpha = 0$ then $\alpha^\gamma = 0 \wedge \alpha^{\beta+\gamma} = 0$. Thus $\alpha^\beta \cdot \alpha^\gamma = 0 = \alpha^{\beta+\gamma}$. If

$\alpha = 1$ then $\alpha^\beta \cdot \alpha^\gamma = 1 \cdot 1 = 1 = \alpha^{\beta+\gamma}$. If $\alpha > 1$ then $\alpha^\gamma \in K_{II}$ and

$$\alpha^\beta \cdot \alpha^\gamma = \bigcup_{\delta < \alpha^\gamma} \alpha^\beta \delta,$$

$$\alpha^{\beta+\gamma} = \bigcup_{\eta < \beta+\gamma} \alpha^\eta.$$

If $\delta < \alpha^\gamma$ then by Theorem 8.40 $(\exists \tau) [\tau < \gamma \wedge \delta < \alpha^\tau]$. Since by the induction hypothesis $\tau < \gamma \rightarrow \alpha^\beta \alpha^\tau = \alpha^{\beta+\tau}$, and since $\tau < \gamma \rightarrow \beta + \tau < \beta + \gamma$

$$\alpha^\beta \delta \leq \alpha^\beta \alpha^\tau \leq \alpha^{\beta+\tau}$$

i.e. $\alpha^\beta \cdot \alpha^\gamma \leq \alpha^{\beta+\gamma}$. Also $\eta < \beta + \gamma \rightarrow \eta \leq \beta \vee (\exists \tau) [\eta = \beta + \tau]$. If $\eta \leq \beta$ then

$$\alpha^\eta \leq \alpha^\beta \cdot 1 \wedge 1 < \alpha^\gamma$$

If $\eta = \beta + \tau$ then $\tau < \gamma$. Hence $\alpha^{\beta+\tau} = \alpha^\beta \cdot \alpha^\tau$ and

$$\alpha^\eta = \alpha^{\beta+\tau} = \alpha^\beta \cdot \alpha^\tau \wedge \alpha^\tau < \alpha^\gamma.$$

Thus $\alpha^\beta \cdot \alpha^\gamma = \alpha^{\beta+\gamma}$.

Theorem 8.42. $(\alpha^\beta)^\gamma = \alpha^{\beta\gamma}$.

Proof (by transfinite induction on γ). $(\alpha^\beta)^0 = 1 = \alpha^{\beta+0}$. If $(\alpha^\beta)^\gamma = \alpha^{\beta\gamma}$ then $(\alpha^\beta)^{\gamma+1} = (\alpha^\beta)^\gamma \alpha^\beta = \alpha^{\beta\gamma} \alpha^\beta = \alpha^{\beta\gamma+\beta} = \alpha^{\beta(\gamma+1)}$. If $\gamma \in K_{II}$ then $\beta = 0 \vee \beta\gamma \in K_{II}$. If $\beta = 0$ then $(\alpha^\beta)^\gamma = 1^\gamma = 1 = \alpha^{\beta\gamma}$. If $\beta\gamma \in K_{II}$ then $\alpha = 0 \vee \alpha \neq 0$. If $\alpha = 0$ then $\alpha^\beta = 0$ and hence $(\alpha^\beta)^\gamma = 0 = \alpha^{\beta\gamma}$. If $\alpha \neq 0$ then $\alpha^\beta \neq 0$ and

$$(\alpha^\beta)^\gamma = \bigcup_{\delta < \gamma} (\alpha^\beta)^\delta,$$

$$\alpha^{\beta\gamma} = \bigcup_{\eta < \beta\gamma} \alpha^\eta.$$

If $\delta < \gamma$ then by the induction hypothesis $(\alpha^\beta)^\delta = \alpha^{\beta\delta}$. Since $\delta < \gamma \rightarrow \beta\delta < \beta\gamma$ we have that $(\alpha^\beta)^\delta \leq \alpha^{\beta\gamma}$. If $\eta < \beta\gamma$ then $(\exists \delta) [\delta < \gamma \wedge \eta < \beta\delta]$. Hence

$$\alpha^\eta < \alpha^{\beta\delta} \quad \text{and} \quad \delta < \gamma.$$

Therefore $\alpha^{\beta\gamma} = (\alpha^\beta)^\gamma$.

Theorem 8.43. $\alpha > 1 \wedge \gamma_n < \alpha \wedge \dots \wedge \gamma_0 < \alpha \wedge 0 \leq \beta_0 < \dots < \beta_n < \beta$
 $\rightarrow \alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0 < \alpha^\beta$.

Proof (by induction on n). If $n=0$ then since $\gamma_0 < \alpha$ we have that $\alpha^{\beta_0} \gamma_0 < \alpha^{\beta_0+1} \leq \alpha^\beta$. If $n > 0$ then since $\beta_{n-1} < \beta_n < \beta$ we have as our induction hypothesis

$$\alpha^{\beta_{n-1}} \gamma_{n-1} + \dots + \alpha^{\beta_0} \gamma_0 < \alpha^{\beta_n}.$$

Therefore

$$\alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0 < \alpha^{\beta_n} \gamma_n + \alpha^{\beta_n} = \alpha^{\beta_n} (\gamma_n + 1).$$

Since $\gamma_n < \alpha$ we have $\gamma_n + 1 \leq \alpha$ and hence

$$\alpha^{\beta_n}(\gamma_n + 1) \leq \alpha^{\beta_n+1} \leq \alpha^\beta.$$

Theorem 8.44. $\beta > 0 \wedge \alpha > 1 \rightarrow (\exists! n) (\exists! \beta_0) \dots (\exists! \beta_n) (\exists! \gamma_0) \dots (\exists! \gamma_n)$
 $[\beta = \alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0 \wedge 0 \leq \beta_0 < \beta_1 < \dots < \beta_n \wedge 0 < \gamma_0 < \alpha \wedge \dots$
 $\dots \wedge 0 < \gamma_n < \alpha].$

Proof (by transfinite induction on β). By Theorem 8.38 $(\exists! \delta) [\alpha^\delta \leq \beta < \alpha^{\delta+1}]$. Then $(\exists! \tau) (\exists! \nu) [\beta = \alpha^\delta \tau + \nu \wedge \nu < \alpha^\delta]$. From $\alpha^\delta \tau \leq \beta$ it follows that $\tau < \alpha$. Since $\nu < \alpha^\delta$ it follows from the induction hypothesis that $(\exists! n) (\exists! \beta_0) \dots (\exists! \beta_n) (\exists! \gamma_0) \dots (\exists! \gamma_n) [\nu = \alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0 \wedge 0 \leq \beta_0 < \beta_1 < \dots < \beta_n \wedge 0 < \gamma_0 < \alpha \wedge \dots \wedge 0 < \gamma_n < \alpha]$. From $\gamma_n \geq 1$, $\alpha^{\beta_n} \leq \alpha^{\beta_n} \gamma_n$. But $\alpha^{\beta_n} \gamma_n \leq \nu < \alpha^\delta$ i.e. $\alpha^{\beta_n} < \alpha^\delta$. Since $\alpha > 1$ it then follows from Corollary 8.34 that $\beta_n < \delta$, thus

$$\beta = \alpha^\delta \tau + \alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0,$$

$$0 \leq \beta_0 < \beta_1 < \dots < \beta_n < \delta \wedge 0 < \gamma_0 < \alpha \wedge \dots \wedge 0 < \gamma_n < \alpha \wedge 0 < \tau < \alpha.$$

Furthermore if

$$\beta = \alpha^{\beta'_{m+1}} \gamma'_{m+1} + \dots + \alpha^{\beta'_0} \gamma'_0,$$

$0 \leq \beta'_0 < \dots < \beta'_{m+1} \wedge 0 < \gamma'_0 < \alpha \wedge \dots \wedge 0 < \gamma'_{m+1} < \alpha$ then by Theorem 8.43

$$\alpha^{\beta'_{m+1}} \leq \beta < \alpha^{\beta'_{m+1}+1}.$$

From Theorem 8.38, $\beta'_{m+1} = \delta$. Since by Theorem 8.43

$$\alpha^{\beta'_m} \gamma'_m + \dots + \alpha^{\beta'_0} \gamma'_0 < \alpha^{\beta'_{m+1}} = \alpha^\delta$$

it follows from the division algorithm (Theorem 8.27) that $\gamma'_m = \tau$ and $\nu = \alpha^{\beta'_m} \gamma'_m + \dots + \alpha^{\beta'_0} \gamma'_0$. Therefore from the induction hypothesis $m = n \wedge \beta'_m = \beta_m \wedge \dots \wedge \beta'_0 = \beta_0 \wedge \gamma'_m = \gamma_m \wedge \dots \wedge \gamma'_0 = \gamma_0$.

Theorem 8.45. $\alpha > 1 \wedge \beta_0 < \beta_1 < \dots < \beta_n \wedge 0 < \gamma_0 < \alpha \wedge \dots \wedge 0 < \gamma_n < \alpha \wedge \delta \geq \omega \rightarrow (\alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0) \alpha^\delta = \alpha^{\beta_n+\delta}$.

Proof. From Theorem 8.43.

$$\alpha^{\beta_n} \leq \alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0 < \alpha^{\beta_n+1}.$$

Therefore

$$\alpha^{\beta_n+\delta} = \alpha^{\beta_n} \alpha^\delta \leq (\alpha^{\beta_n} \gamma_n + \dots + \alpha^{\beta_0} \gamma_0) \alpha^\delta \leq \alpha^{\beta_n+1} \alpha^\delta = \alpha^{\beta_n+1+\delta} = \alpha^{\beta_n+\delta}.$$

Theorem 8.46. $\alpha \in K_{II} \wedge \beta_0 < \beta_1 < \dots < \beta_n \wedge 0 < m_n \wedge \delta > 0 \rightarrow (\alpha^{\beta_n} m_n + \dots + \alpha^{\beta_0} m_0) \alpha^\delta = \alpha^{\beta_n+\delta}$.

Proof. From Theorem 8.43.

$$\alpha^{\beta_{n-1}} m_{n-1} + \dots + \alpha^{\beta_0} m_0 < \alpha^{\beta_n}.$$

Therefore

$$\alpha^{\beta_n} \leq \alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0 < \alpha^{\beta_n} m_n + \alpha^{\beta_n} = \alpha^{\beta_n} (m_n + 1).$$

Then

$$\alpha^{\beta_n} \alpha^\delta \leq (\alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0) \alpha^\delta \leq \alpha^{\beta_n} (m_n + 1) \alpha^\delta = \alpha^{\beta_n} \alpha^\delta = \alpha^{\beta_n + \delta}.$$

Theorem 8.47. *If $\alpha \in K_{II} \wedge \beta > 0 \wedge m > 0$ then*

- 1) $(\alpha^\beta m)^\gamma = \alpha^{\beta \gamma} m, \gamma \in K_I \wedge \gamma \neq 0.$
- 2) $(\alpha^\beta m)^\gamma = \alpha^{\beta \gamma}, \gamma \in K_{II}.$

Proof (by transfinite induction on γ). $(\alpha^\beta m)^1 = \alpha^{\beta \cdot 1} m$. If $(\alpha^\beta m)^\gamma = \alpha^{\beta \gamma} m$ then $(\alpha^\beta m)^{\gamma+1} = (\alpha^\beta m)^\gamma \alpha^\beta m = \alpha^{\beta \gamma} m \alpha^\beta m = \alpha^{\beta \gamma} \alpha^\beta m = \alpha^{\beta(\gamma+1)} m$. If $(\alpha^\beta m)^\gamma = \alpha^{\beta \gamma}$ then $(\alpha^\beta m)^{\gamma+1} = (\alpha^\beta m)^\gamma \alpha^\beta m = \alpha^{\beta \gamma} \alpha^\beta m = \alpha^{\beta(\gamma+1)} m$. If $\gamma \in K_{II}$ then $(\alpha^\beta)^\gamma \leq (\alpha^\beta m)^\gamma = \bigcup_{\delta < \gamma} (\alpha^\beta m)^\delta \leq \bigcup_{\delta < \gamma} \alpha^{\beta \delta} m \leq \bigcup_{\delta < \gamma} \alpha^{\beta(\delta+1)} = \alpha^{\beta \gamma}.$

Theorem 8.48.

$$\alpha \in K_{II} \wedge \beta_0 < \beta_1 < \cdots < \beta_n \rightarrow (\alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0)^\gamma \leq \alpha^{\beta_n \gamma} (m_n + 1).$$

Proof. $\alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0 \leq \alpha^{\beta_n} (m_n + 1)$. Therefore by Theorem 8.47 $(\alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0)^\gamma \leq [\alpha^{\beta_n} (m_n + 1)]^\gamma \leq \alpha^{\beta_n \gamma} (m_n + 1)$.

Theorem 8.49.

$$\alpha \in K_{II} \wedge \beta_0 < \beta_1 < \cdots < \beta_n \wedge \gamma \in K_{II} \rightarrow (\alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0)^\gamma = \alpha^{\beta_n \gamma}.$$

Proof. $\alpha^{\beta_n} \leq \alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0 \leq \alpha^{\beta_n} (m_n + 1)$. Therefore by Theorem 8.47

$$(\alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0)^\gamma = \alpha^{\beta_n \gamma}.$$

Corollary 8.50.

$$\alpha \in K_{II} \wedge \beta_0 < \beta_1 < \cdots < \beta_n \wedge \gamma > 0 \rightarrow (\alpha^{\beta_n} m_n + \cdots + \alpha^{\beta_0} m_0)^{\alpha^\gamma} = \alpha^{\beta_n \alpha^\gamma}.$$

Proof. $\gamma > 0 \wedge \alpha \in K_{II} \rightarrow \alpha^\gamma \in K_{II}.$

9 Relational Closure and the Rank Function

In this section we introduce two ideas important for the work to follow. The first of these is relational closure. In later sections we will be especially interested in sets that are transitive. While there exist sets that are not transitive every set has a transitive extension. Indeed, every set has a smallest transitive extension which we call its transitive closure.

Theorem 9.1. $(\forall a) (\exists b) [a \subseteq b \wedge \text{Tr}(b) \wedge (\forall y) [a \subseteq y \wedge \text{Tr}(y) \rightarrow b \subseteq y]$.

Proof. If $G'x = x \cup (\cup(x))$, if $f \mathcal{F}^n \omega \wedge f'0 = a \wedge f(n+1) = G'f'n$ and if

$$b = \bigcup_{n < \omega} f'n$$

then $a = f'0 \subseteq b$. From the definition of G

$$f(n) \subseteq f(n+1) \wedge \cup(f'n) \subseteq f(n+1).$$

If $z \in y \wedge y \in b$ then $(\exists n) [y \in f'n]$ and hence

$$z \in \cup(f'n) \subseteq f(n+1)$$

i.e., $z \in f(n+1)$. Then $z \in b$ and hence b is transitive.

If $a \subseteq y \wedge \text{Tr}(y)$ then we prove by induction that

$$f'n \subseteq y$$

$f'0 = a \subseteq y$. If $f'n \subseteq y$ then since y is transitive $\cup(f'n) \subseteq y$ i.e.,

$$z \in f'n \wedge f'n \subseteq y \rightarrow z \in y.$$

Thus $f(n+1) = f(n) \cup (\cup(f'n)) \subseteq y$.

Consequently $b = \bigcup_{n < \omega} f'n \subseteq y$.

Definition 9.2. $\text{Tr Cl}(a) = \cap \{y \mid a \subseteq y \wedge \text{Tr}(y)\}$.

Remark. Theorem 9.1 has a natural and useful generalization to well founded relations.

Theorem 9.3. *If R Wfr $A \wedge a \subseteq A$ then $(\exists b) [[a \subseteq b \subseteq A]$ and*

- 1) $(\forall x \in A) (\forall y) [x R y \wedge y \in b \rightarrow x \in b]$.
- 2) $(\forall x \in b) [x \in a \vee (\exists n) (\exists g) [g: n+1 \rightarrow b \wedge g(0) \in a \wedge g(n) = x \wedge (\forall i < n) [g(i+1) R g(i)]]]$.
- 3) $(\forall c) [[a \subseteq c \subseteq A \wedge (\forall x \in A) (\forall y) [x R y \wedge y \in c \rightarrow x \in c]] \rightarrow b \subseteq c]$.

Proof.

- 1) Since R is well founded on A , $x \in A \rightarrow \mathcal{M}(A \cap R^{-1}\{x\})$. Therefore if

$$B = \{\langle x, A \cap R^{-1}\{x\} \rangle \mid x \in A\}$$

then B is a function. If $y \subseteq A$ then $B''y$ is a set hence so is $\cup(B''y)$. But

$$\begin{aligned} \cup(B''y) &= \cup\{z \mid (\exists x \in y) [\langle x, z \rangle \in B]\} \\ &= \cup\{z \mid (\exists x \in y) [z = A \cap R^{-1}\{x\}]\} \\ &= \cup\{A \cap R^{-1}\{x\} \mid x \in y\} \\ &= A \cap R^{-1}y. \end{aligned}$$

Thus $A \cap R^{-1}y$ is a set.

If $G'x = x \cup (A \cap R^{-1}x) \wedge f \mathcal{F}_n \omega \wedge f'0 = a \wedge f(n+1) = G'f'n$. Then $f'0 = a \subseteq A$. If $f(n)$ is a subset of A then since

$$f(n+1) = f(n) \cup (A \cap R^{-1}f'n)$$

$f(n+1)$ is a subset of A . Thus $\cup(f''\omega)$ is a subset of A . If $b = \cup(f''\omega)$ then $a = f'0 \subseteq b$. From the definition of G

$$f'n \subseteq f(n+1) \wedge A \cap R^{-1}f'n \subseteq f(n+1).$$

If $x \in A \wedge x R y \wedge y \in b$ then $(\exists n) [x R y \wedge y \in f'n]$ i.e., $x \in R^{-1}f'n \subseteq f(n+1)$. Thus $x \in b$.

- 2) $(\forall x \in b) (\exists n) [x \in f'n]$. If $n=0$ then $f'n = a \wedge x \in a$. If the result holds for each element in $f'n$ and $x \in f(n+1)$ then since

$$f(n+1) = f(n) \cup (A \cap R^{-1}f'n)$$

it follows that either $x \in f(n)$, hence the conclusion follows from the induction hypothesis, or $(\exists y \in f'n) [x R y]$ in which case the conclusion again follows from the induction hypothesis.

- 3) The proof is left to the reader.

Theorem 9.4. R Wfr $A \wedge B \subseteq A \wedge B \neq 0 \rightarrow (\exists x \in B) [B \cap R^{-1}\{x\} = 0]$.

Proof. Since $B \neq 0$, $\exists a \in B$ and by Theorem 9.3,

$$(\exists b) [\{a\} \subseteq b \subseteq A \wedge (\forall x) (\forall y) [x R y \wedge y \in b \rightarrow x \in b]].$$

Then $b \cap B$ is a nonempty subset of A . Therefore

$$(\exists x \in b \cap B) [(b \cap B) \cap R^{-1}\{x\} = 0].$$

If $y \in B \cap R^{-1}\{x\}$ then $y \in B \wedge y R x$. But $x \in b$ and hence $y \in b$ i.e.,

$$y \in [b \cap B \cap R^{-1}\{x\}].$$

Therefore $B \cap R^{-1}\{x\} = 0$.

Remark. Note in Theorem 9.1 that the set b has the property $E^{-1}b \subseteq b$ and in Theorem 9.3, $R^{-1}b \subseteq b$. We say that b is closed with respect to (w.r.t.) the binary relations E^{-1} and R^{-1} respectively.

Definition 9.5.

$$1) \quad \text{Cl}(R, A) \xleftrightarrow{\Delta} R^{\Delta} A \subseteq A.$$

$$2) \quad \text{Cl}_2(R, A) \xleftrightarrow{\Delta} R^{\Delta} A^2 \subseteq A.$$

Theorem 9.6. If $\text{Cl}(R_1, A) \wedge \dots \wedge \text{Cl}(R_p, A) \wedge \text{Cl}_2(S_1, A) \wedge \dots$
 $\dots \wedge \text{Cl}_2(S_q, A)$ and if $\forall x \subseteq A, \mathcal{M}(R_1^{\Delta} x) \wedge \dots \wedge \mathcal{M}(R_p^{\Delta} x) \wedge \mathcal{M}(S_1^{\Delta} x^2) \wedge \dots$
 $\dots \wedge \mathcal{M}(S_q^{\Delta} x^2)$ then

$$(\forall a \subseteq A) (\exists b) [a \subseteq b \subseteq A \wedge \text{Cl}(R_1, b) \wedge \dots \wedge \text{Cl}(R_p, b) \wedge \text{Cl}_2(S_1, b) \wedge \dots$$

$$\dots \wedge \text{Cl}_2(S_q, b)].$$

Proof. If $G^{\Delta} x = x \cup R_1^{\Delta} x \cup \dots \cup R_p^{\Delta} x \cup S_1^{\Delta} x^2 \cup \dots \cup S_q^{\Delta} x^2, x \subseteq A$ and if

$$f \mathcal{F}_n \omega \wedge f^{\Delta} 0 = a \wedge f(n+1) = G^{\Delta} f^{\Delta} n \wedge b = \bigcup_{n < \omega} f^{\Delta} n$$

then from the definition of G

$$f(n+1) = f^{\Delta} n \cup R_1^{\Delta} f^{\Delta} n \cup \dots \cup R_p^{\Delta} f^{\Delta} n \cup S_1^{\Delta} (f^{\Delta} n)^2 \cup \dots \cup S_q^{\Delta} (f^{\Delta} n)^2.$$

$f^{\Delta} 0 = a \subseteq A$. If $f^{\Delta} n \subseteq A$ then since each R_i and each S_i is closed on A it follows that $f(n+1) \subseteq A$. Therefore $b \subseteq A$. Furthermore $f^{\Delta} 0 = a \subseteq b$.

If $y \in R_i^{\Delta} b$ then $(\exists x \in b) [\langle x, y \rangle \in R_i]$. But

$$x \in b \rightarrow (\exists n) [x \in f^{\Delta} n].$$

Thus

$$y \in R_i^{\Delta} f^{\Delta} n \subseteq f(n+1).$$

Therefore $y \in b$, i.e., $R_i^{\Delta} b \subseteq b$.

If $z \in S_i^{\Delta} b^2$ then $\exists x, y \in b, \langle x, y, z \rangle \in S_i$. But

$$x, y \in b \rightarrow (\exists m, n) [x \in f^{\Delta} m \wedge y \in f^{\Delta} n].$$

If $r = \max(m, n)$ then $x, y \in f^{\Delta} r$, i.e.,

$$z \in S_i^{\Delta} (f^{\Delta} r)^2 \subseteq f(r+1).$$

Therefore $z \in b$, i.e., $S_i^{\Delta} b^2 \subseteq b$.

Theorem 9.7. If $R \text{ Wfr } A$, if $K = \{f \mid (\exists a \subseteq A) [\text{Cl}(R^{-1}, a) \wedge f \mathcal{F}_n a \wedge (\forall x \in a) [f'x = G(f \upharpoonright R^{-1} \{x\})]]\}$ and if $F = \cup(K)$, then

- 1) $F \mathcal{F}_n A$.
- 2) $(\forall x \in A) [F'x = G(F \upharpoonright R^{-1} \{x\})]$.
- 3) F is unique.

The proof is left to the reader.

Remark. Theorem 9.1 assures us that every set a has a smallest transitive extension. This extension of a we call the transitive closure of a . In order to define "rank" we are interested in sets that are not only transitive but supertransitive in the sense of

Definition 9.8. $\text{St}(A) \stackrel{\Delta}{\leftrightarrow} \text{Tr}(A) \wedge (\forall x) [x \in A \rightarrow P(x) \subseteq A]$.

Definition 9.9.

$$\begin{aligned} R(0) &= 0 \\ R(\alpha + 1) &= \mathcal{P}(R'\alpha). \\ R(\alpha) &= \bigcup_{\beta < \alpha} R'\beta, \alpha \in K_{\text{II}}. \end{aligned}$$

Theorem 9.10.

- 1) $\mathcal{M}(R'\alpha) \wedge \text{St}(R'\alpha)$.
- 2) $\alpha < \beta \rightarrow R'\alpha \in R'\beta \wedge R'\alpha \subseteq R'\beta$.

Proof.

1) (By transfinite induction on α .) If $\alpha = 0$ then $R'\alpha = 0$ and hence $R'\alpha$ is a supertransitive set. If $R'\alpha$ is a supertransitive set then, since $R(\alpha + 1) = \mathcal{P}(R'\alpha)$, $R(\alpha + 1)$ is a set and

$$\begin{aligned} x \in z \wedge y \subseteq z \wedge z \in R(\alpha + 1) &\rightarrow x \in z \wedge y \subseteq z \wedge z \subseteq R'\alpha \\ &\rightarrow x \in R'\alpha \wedge y \subseteq R'\alpha \\ &\rightarrow x \subseteq R'\alpha \wedge y \subseteq R'\alpha \\ &\rightarrow x \in \mathcal{P}(R'\alpha) \wedge y \in \mathcal{P}(R'\alpha), \end{aligned}$$

i.e., $z \in R(\alpha + 1) \rightarrow z \subseteq R(\alpha + 1) \wedge \mathcal{P}(z) \subseteq R(\alpha + 1)$.

If $\alpha \in K_{\text{II}}$ then $z \in R'\alpha \rightarrow (\exists \beta < \alpha) [z \in R'\beta]$

$$\begin{aligned} &\rightarrow (\exists \beta < \alpha) [z \subseteq R'\beta \wedge \mathcal{P}(z) \subseteq R'\beta] \\ &\rightarrow z \subseteq R'\alpha \wedge \mathcal{P}(z) \subseteq R'\alpha. \end{aligned}$$

Since $R'\alpha$ is the union of a set it is a set.

2) Since $R'\beta$ is transitive it is sufficient to prove that $R'\alpha \in R'\beta$. This we do by induction on β . Since $R'\beta \subseteq R'\beta$ we have $R'\beta \in \mathcal{P}(R'\beta) = R(\beta + 1)$. In particular $R'\alpha \in R(\alpha + 1)$.

If $\alpha < \beta \rightarrow R'\alpha \in R'\beta$ then since $\alpha < \beta + 1 \rightarrow \alpha < \beta \vee \alpha = \beta$ we have

$$\alpha = \beta \rightarrow R'\alpha \in R(\beta + 1)$$

and, from the induction hypothesis

$$\begin{aligned} \alpha < \beta &\rightarrow R'\alpha \in R'\beta \wedge R'\alpha \subseteq R(\beta + 1) \\ &\rightarrow R'\alpha \in R(\beta + 1). \end{aligned}$$

If $\beta \in K_{II}$ then

$$R'\beta = \bigcup_{\gamma < \beta} R'\gamma.$$

Since by the induction hypothesis $\alpha < \beta \rightarrow R'\alpha \in R(\alpha + 1)$ it follows that

$$\alpha < \beta \rightarrow R'\alpha \in R'\beta.$$

Definition 9.11. $Wf(a) \stackrel{\Delta}{\leftrightarrow} (\exists \alpha) [a \in R'\alpha].$

Remark. $Wf(a)$ is read “ a is well founded”. With the aid of the following theorem and the Axiom of Regularity we can prove that every set is well founded.

Theorem 9.12. $(\forall x \in a) [Wf(x)] \rightarrow Wf(a).$

Proof. Since $x \in a \rightarrow (\exists \alpha) [x \in R'\alpha]$ there is a smallest such α . If

$$F'x = \mu_{\alpha}(x \in R'\alpha)$$

then since F is a function $F'a$ is a set and indeed a subset of On . Therefore $\cup(F'a)$ is an ordinal. If $\beta = \cup(F'a) + 1$ then $F'a \subseteq \beta$ i.e., $x \in a \rightarrow F'x < \beta$. By Theorem 9.10

$$R'F'x \subseteq R'\beta$$

Also $x \in R'F'x$ i.e.,

$$\begin{aligned} x \in a &\rightarrow x \in R'F'x \\ &\rightarrow x \in R'\beta. \end{aligned}$$

Thus $a \subseteq R'\beta$ and hence $a \in \mathcal{P}(R'\beta) = R(\beta + 1)$.

Theorem 9.13. $(\forall a) [Wf(a)].$

Proof. From the Axiom of Regularity E is a well founded relation on V . Then from Theorem 9.12 if

$$A = \{x \mid Wf(x)\}$$

if $a \subseteq A$ then $a \in A$. Then by \in -induction (Theorem 5.24) $A = V$.

Remark. From Theorem 9.13 we see that in the presence of the Axiom of Regularity the function R determines a class of sets $\{R'\alpha \mid \alpha \in On\}$

whose union is the entire universe. Furthermore, from Theorem 9.10, these sets are nested i.e., $\alpha < \beta \rightarrow R'\alpha \subset R'\beta$.

We offer the following pictorial representation of this nesting of sets. The universe is represented as the points in a V-shaped wedge.




Fig. 2.

If $\alpha \in K_{II}$ then $R'\alpha = \bigcup_{\beta < \alpha} R'\beta$. Thus any set in $R'\alpha$ is also in some $R'\beta$ with $\beta < \alpha$. Then for each a the smallest ordinal β for which $a \in R'\beta$ is a nonlimit ordinal i.e.,

$$(\exists \alpha) [a \notin R'\alpha \wedge a \in R(\alpha + 1)].$$

This particular ordinal α we call the rank of a .

Definition 9.14. $\text{rank}(a) = \mu_{\alpha}(a \in R(\alpha + 1))$.

Theorem 9.15.

- 1) $\text{rank}(a) \in \text{On}$.
- 2) $\alpha = \text{rank}(a) \leftrightarrow a \notin R'\alpha \wedge a \in R(\alpha + 1)$.
- 3) $\beta \leq \text{rank}(a) \leftrightarrow a \notin R'\beta$.

Proof.

1) Definition 9.14.

2) From Definition 9.14 if $\alpha = \text{rank}(a)$ then $a \in R(\alpha + 1)$. If $\alpha = 0$ then $a \in R'\alpha \rightarrow \alpha \in 0$. If $(\exists \gamma) [\alpha = \gamma + 1]$ then $a \in R'\alpha \rightarrow \gamma \geq \alpha$. If $\alpha \in K_{II}$ then $a \in R'\alpha \rightarrow (\exists \beta < \alpha) [a \in R'\beta]$. But $R'\beta \subseteq R(\beta + 1)$, hence $a \in R(\beta + 1)$. But then $\beta \geq \alpha$. Since in each case we have a contradiction we conclude that $\alpha = \text{rank}(a) \rightarrow a \notin R'\alpha$.

Conversely $a \in R(\alpha + 1) \rightarrow \alpha \geq \text{rank}(a)$. If $a \notin R'\alpha$ then since $\beta \leq \alpha \rightarrow R'\beta \subseteq R'\alpha$ we have that $\beta \leq \alpha \rightarrow a \notin R'\beta$. But $x \in R(\text{rank}(x) + 1)$. Therefore $\alpha < \text{rank}(x) + 1$ i.e., $\alpha \leq \text{rank}(x)$. Thus $\alpha = \text{rank}(a)$.

3) If $\alpha = \text{rank}(a)$ then by 2) $a \notin R'\alpha$. Furthermore $\beta \leq \alpha \rightarrow R'\beta \subseteq R'\alpha$. Therefore $a \notin R'\beta$. Also $\alpha < \beta \rightarrow R(\alpha + 1) \subseteq R'\beta$. Since $a \in R(\alpha + 1)$, $a < \beta \rightarrow a \in R'\beta$.

Theorem 9.16. $a \in b \rightarrow \text{rank}(a) < \text{rank}(b)$.

Proof. By Theorem 9.15, $\alpha = \text{rank}(a) \rightarrow a \notin R'\alpha$. Then, since $a \in b$, $b \notin R'\alpha$ and hence $b \notin R(\alpha + 1)$. Thus $\alpha < \text{rank}(b)$.

Theorem 9.17. $\text{rank}(a) = \mu_\beta((\forall x \in a) [\text{rank}(x) < \beta])$.

Proof. From Theorem 9.16, $x \in a \rightarrow \text{rank}(x) < \text{rank}(a)$. Furthermore, if

$$x \in a \rightarrow \text{rank}(x) < \beta$$

then since $x \in R(\text{rank}(x) + 1) \subseteq R'\beta$ we have $a \subseteq R'\beta$ and hence $a \in R(\beta + 1)$ i.e.,

$$\text{rank}(a) \leq \beta.$$

Theorem 9.18. $\text{rank}(\alpha) = \alpha$.

Proof (by transfinite induction on α). If as our induction hypothesis we have

$$\gamma < \alpha \rightarrow \text{rank}(\gamma) = \gamma.$$

Then from Theorem 9.17

$$\text{rank}(\alpha) = \mu_\beta((\forall \gamma < \alpha) [\text{rank}(\gamma) < \beta]) = \mu_\beta(\beta \geq \alpha) = \alpha.$$

Theorem 9.19. $(\exists \alpha) (\forall x \in A) [\text{rank}(x) \leq \alpha] \rightarrow \mathcal{M}(A)$.

Proof. $\text{rank}(x) \leq \alpha \rightarrow x \in R(\alpha + 1)$. Hence $A \subseteq R(\alpha + 1)$.

Exercises.

1) $a \subseteq b \rightarrow \text{rank}(a) \leq \text{rank}(b)$.

2) $\mathcal{P}_r(A) \rightarrow (\forall \alpha) (\exists a \in A) [\text{rank}(a) > \alpha]$.

Remark. Earlier we promised to prove the equivalence of the weak and strong forms of the Axiom of Regularity. We redeemed that promise with Theorem 9.4. We now give a second proof using the rank function.

Theorem 9.20. (Axiom 6'). $A \neq 0 \rightarrow (\exists x \in A) [A \cap x = 0]$.

Proof. If $B = \{\text{rank}(x) \mid x \in A\}$ then $A \neq 0 \rightarrow B \neq 0$. Thus B is a nonempty class of ordinals, hence, by Theorem 6.26, which was proved using only the weak form of the Axiom of Regularity, B has an E-minimal element α . Since $\alpha \in B$ it follows that

$$(\exists x \in A) [\alpha = \text{rank}(x)].$$

Furthermore since α is an E-minimal element of B it follows from Theorem 9.16 that

$$A \cap x = 0.$$

Remark. The simplicity of the proof of Theorem 9.20 illustrates the power of the rank function. Indeed with the aid of the rank function we can prove the following generalization of Theorem 9.4.

Theorem 9.21. $R \text{ Fr } A \wedge B \subseteq A \wedge B \neq 0 \rightarrow (\exists x \in B) [B \cap R^{-1}\{x\} = 0].$

Proof (by contradiction). If

$$B_0 = \{x \in B \mid (\forall y \in B) [\text{rank}(x) \leq \text{rank}(y)]\},$$

$$B_{n+1} = \{x \in B \mid (\exists y \in B_n) [x R y \wedge (\forall z \in B) [z R y \rightarrow \text{rank}(x) \leq \text{rank}(z)]]\},$$

then all elements of B_0 have the same rank hence B_0 is a set. If B_n is a set then since

$$B_{n+1} = \bigcup_{y \in B_n} \{x \in B \mid x R y \wedge (\forall z \in B) [z R y \rightarrow \text{rank}(x) \leq \text{rank}(z)]\} \quad B_{n+1} \text{ is a}$$

set. Thus $(\forall n \geq 0) [\mathcal{M}(B_n)]$. If

$$b = \bigcup_{n < \omega} B_n$$

then $b \subseteq B \subseteq A \wedge b \neq 0$.

If $(\forall x \in B) [B \cap R^{-1}\{x\} \neq 0]$ then in particular

$$x \in b \rightarrow B \cap R^{-1}\{x\} \neq 0.$$

Furthermore $(\forall x \in b) (\exists n) [x \in B_n]$. Since $B \cap R^{-1}\{x\}$ is not empty it contains an element of minimal rank i.e.,

$$(\exists y \in B) [y R x \wedge (\forall z \in B) [z R x \rightarrow \text{rank}(y) \leq \text{rank}(z)]].$$

Since $x \in B_n, y \in B_{n+1}$ and hence $y \in b$. But $y R x$, that is

$$(\forall x \in b) [b \cap R^{-1}\{x\} \neq 0].$$

This contradicts the fact that $R \text{ Fr } A$.

Theorem 9.22. *If $R \text{ Fr } A \wedge B \subseteq A \wedge (\forall x \in A) [A \cap R^{-1}\{x\} \subseteq B \rightarrow x \in B]$ then $A = B$.*

The proof is left to the reader.

10 Cardinal Numbers

The equivalence of sets is basic to the theory of cardinal numbers. Two sets are equivalent, or equipollent, provided there exists a one-to-one correspondence between them.

Definition 10.1. $a \simeq b \stackrel{\Delta}{\leftrightarrow} (\exists f) [f : a \xrightarrow{\text{onto}} b]$.

Theorem 10.2.

- 1) $a \simeq a$.
- 2) $a \simeq b \rightarrow b \simeq a$.
- 3) $a \simeq b \wedge b \simeq c \rightarrow a \simeq c$.

The proof is left to the reader.

Remark. A basic result for the theory of set equivalence is the following theorem first proved by Cantor. Cantor's proof however tacitly assumed the Axiom of Choice. Ernst Schröder in 1896 and independently Felix Bernstein in 1898 gave proofs that did not require the Axiom of Choice. The theorem states that if a set a is equivalent to a subset of b and if b is equivalent to a subset of a then $a \simeq b$.

Theorem 10.3 (Cantor-Schröder-Bernstein).

$$a \simeq c \wedge c \subseteq b \wedge b \simeq d \wedge d \subseteq a \rightarrow a \simeq b .$$

Proof. $a \simeq c \rightarrow (\exists f) [f : a \xrightarrow{\text{onto}} c]$,

$$b \simeq d \rightarrow (\exists g) [g : b \xrightarrow{\text{onto}} d] .$$

If $H'x = (g \circ f)''x$ then there exists a function h defined on ω such that

$$h(0) = a - d ,$$

$$h(n + 1) = H'h'n = (g \circ f)''h'n .$$

Since $h'0 \subseteq a$ and since $g \circ f$ maps a into a it follows by induction that $(\forall n) [h'n \subseteq a]$. Consequently $(\forall n) [f''h'n \subseteq b]$.

We next define a function F on a in the following way

$$\begin{aligned} F'x &= f(x) \quad \text{if } x \in a \wedge (\exists n) [x \in h'n] \\ &= g^{-1}(x) \quad \text{if } x \in a \wedge (\forall n) [x \notin h'n]. \end{aligned}$$

Then $F : a \rightarrow b$. To prove that F is onto we note that

$$y \in b \rightarrow (\exists n) [y \in f'h'n] \vee (\forall n) [y \notin f'h'n].$$

If $(\exists n) [y \in f'h'n]$ then $(\exists x \in h'n) [y = f'x]$. But $x \in h'n \rightarrow x \in a$ and $x \in a \wedge x \in h'n \rightarrow F'x = f(x) = y$.

If $(\forall n) [y \notin f'h'n]$ then it follows that $(\forall n) [g'y \notin h'n]$ for if $(\exists n) [g'y \in h'n]$ then since $h'0 = a - d$ and since $g'y \in d$ it follows that $n \neq 0$. Therefore $(\exists m) [n = m + 1]$. But $h(m + 1) = g'f'h'm$. Consequently, since g is one-to-one, $g'y \in h(m + 1) \rightarrow y \in f'h'm$. This is a contradiction from which we conclude that $(\forall n) [g'y \notin h'n]$. On the other hand since $y \in b$ it follows that $g'y \in a$. Therefore $F'g'y = g^{-1}(g'y) = y$ and hence $\mathcal{W}(F) = b$.

To prove that F is one-to-one we note that if $x \in a \wedge y \in a \wedge F'x = F'y$ then $(\exists m) [x \in h'm] \leftrightarrow (\exists m) [y \in h'm]$ for if $x \in h'm \wedge (\forall n) [y \notin h'n]$ then $F'x = F'y \rightarrow f(x) = g^{-1}(y)$ i.e. $y = (g \circ f)'x$. But $x \in h'm$; therefore

$$y \in (g \circ f)'h'm = h(m + 1).$$

Similarly we can prove that $y \in h'm \rightarrow x \in h(m + 1)$.

Thus $F'x = F'y \rightarrow [f(x) = f(y) \vee g^{-1}(x) = g^{-1}(y)]$. Since both f and g are one-to-one it follows that $x = y$. Since F is a function with domain a , F is a set.

Theorem 10.4 (Cantor). $a \not\cong \mathcal{P}(a)$.

Proof. If $(\exists f) [f : a \xrightarrow{\text{onto}} \mathcal{P}(a)]$ and if

$$B = \{b \in a \mid b \notin f'b\}$$

then $B \subseteq a$. Therefore B is a set and hence $B \in \mathcal{P}(a)$. Consequently $(\exists b) [b \in a \wedge B = f'b]$. From the definition of B it then follows that

$$b \in B \leftrightarrow b \notin f'b.$$

But $f'b = B$ i.e. $b \in B \leftrightarrow b \notin B$.

Corollary 10.5. $\neg (\exists f) [f : \mathcal{P}(a) \xrightarrow{1-1} a]$.

Proof. If

$$g'b = \{b\}, \quad b \in a$$

then $g : a \xrightarrow{1-1} \mathcal{P}(a)$. If $(\exists f) [f : \mathcal{P}(a) \xrightarrow{1-1} a]$ then by the Cantor-Bernstein-Schröder Theorem $a \simeq \mathcal{P}(a)$. This contradicts Theorem 10.4.

Theorem 10.6. $a \simeq b \rightarrow \mathcal{P}(a) \simeq \mathcal{P}(b)$.

Proof. $a \simeq b \rightarrow (\exists f) [f : a \xrightarrow[\text{onto}]{1-1} b]$. If $F = \{\langle x, f''x \rangle \mid x \in \mathcal{P}(a)\}$ then $F : \mathcal{P}(a) \rightarrow \mathcal{P}(b)$. Furthermore since $f''x = \{f'z \mid z \in x\}$ it follows that if $f''x = f''y$ then $z \in x \rightarrow f'z \in f''x = f''y$. That is $(\exists w) [w \in y \wedge f'z = f'w]$. But f is one-to-one; therefore $z = w$ and hence $z \in y$. Similarly $z \in y \rightarrow z \in x$. Therefore $x = y$ i.e. F is one-to-one.

Finally if $y \in \mathcal{P}(b)$ and if $x = (f^{-1})''y$ then $x \in \mathcal{P}(a)$ and

$$F'x = f''(f^{-1})''y = y.$$

$$\text{Thus } \mathcal{W}(F) = \mathcal{P}(b).$$

Exercises. Prove the following.

- 1) $a \simeq 0 \leftrightarrow a = 0$.
- 2) $(\forall b) [\{b\} \times a \simeq a \times \{b\} \simeq a]$.
- 3) $(\forall b) [[a \cup \{b\} \simeq a] \vee [a \cup \{b\} \simeq a \cup \{a\}]]$.
- 4) $a_1 \simeq a_2 \wedge b_1 \simeq b_2 \rightarrow a_1 \times b_1 \simeq a_2 \times b_2$.
- 5) $a \simeq b \leftrightarrow a \cup \{a\} \simeq b \cup \{b\}$.
- 6) $a \times b \simeq b \times a$.
- 7) $\alpha \geq \omega \rightarrow \alpha \simeq \alpha + 1$.
- 8) $\alpha \geq \omega \rightarrow \alpha \simeq \alpha + n$.

Remark. Cantor defined the cardinal number of a set M to be “the general concept which, by means of our active faculty of thought, arises from the aggregate M when we make abstraction of the nature of its various elements m and of the order in which they are given.” Cantor’s choice of words suggests that for him a cardinal number is not a mathematical object but a psychological entity.

Frege, in 1884, and Russell, independently in 1903, removed cardinal numbers from this psychic realm by defining the cardinal number of a set a to be the class \bar{a} of all sets that are equivalent to the given set a . We choose instead to define \bar{a} to be the smallest ordinal number that is equivalent to a . If there is an equivalence class that contains no ordinal number then that equivalent class will not be represented among our cardinal numbers. From Cohen’s work we know that the question of whether or not such an equivalence class exists is undecidable in ZF. This question is, however, decided by the Axiom of Choice to be introduced in § 11. To help clarify the role of the Axiom of Choice we will next develop properties of cardinals that do not require the Axiom of Choice.

Definition 10.7. $\bar{a} = \mu_\alpha(\alpha \simeq a)$.

Remark. Note, in Definition 10.7, that if there is no α equivalent to a then $\bar{a} = 0$.

Theorem 10.8. $\bar{a} \in \text{On}$.

Theorem 10.9.

- 1) $a \simeq \bar{a} \leftrightarrow (\exists \alpha) [\alpha \simeq a]$.
- 2) $a \simeq \bar{a} \leftrightarrow (\exists b) [b \subseteq \text{On} \wedge b \simeq a]$.

Theorem 10.10. $\bar{a} = 0 \vee \bar{a} \simeq a$.

Theorem 10.11.

- 1) $\alpha \simeq \bar{\alpha}$.
- 2) $\bar{\alpha} = 0 \leftrightarrow \alpha = 0$.

Theorem 10.12. $(\forall \alpha) [a \simeq \alpha \rightarrow \bar{a} \leq \alpha]$.

Corollary 10.13. $\bar{\alpha} \leq \alpha$.

Theorem 10.14. $a \simeq b \rightarrow \bar{a} = \bar{b}$.

The foregoing results follow easily from Definition 10.7. Their proofs are left to the reader.

Theorem 10.15.

- 1) $a \cup b \simeq \overline{\overline{a \cup b}} \leftrightarrow a \simeq \bar{a} \wedge b \simeq \bar{b}$.
- 2) $a \neq 0 \wedge b \neq 0 \rightarrow [a \times b \simeq \overline{\overline{a \times b}} \leftrightarrow a \simeq \bar{a} \wedge b \simeq \bar{b}]$.

Proof.

- 1) $a \cup b \simeq \overline{\overline{a \cup b}} \rightarrow (\exists f) [f : a \cup b \xrightarrow[1-1]{\text{onto}} \overline{\overline{a \cup b}}]$.

Then $a \simeq f^{-1} \overline{\overline{a \cup b}} \subseteq \overline{\overline{a \cup b}} \wedge b \simeq f^{-1} \overline{\overline{a \cup b}} \subseteq \overline{\overline{a \cup b}}$. From Theorem 10.9, $a \simeq \bar{a} \wedge b \simeq \bar{b}$.

Conversely $a \simeq \bar{a} \wedge b \simeq \bar{b} \rightarrow (\exists f) [f : a \xrightarrow[1-1]{\text{onto}} \bar{a}] \wedge (\exists g) [g : b \xrightarrow[1-1]{\text{onto}} \bar{b}]$. If

$$\begin{aligned} F^{-1}x &= f^{-1}x \quad \text{for } x \in a \\ &= \bar{a} + g^{-1}x \quad \text{for } x \in b - a, \end{aligned}$$

then $F : a \cup b \xrightarrow[1-1]{} \bar{a} + \bar{b}$. Then $a \cup b \simeq F^{-1}(\bar{a} + \bar{b}) \subseteq \bar{a} + \bar{b}$. From Theorem 10.9, $a \cup b \simeq \overline{\overline{a \cup b}}$.

- 2) $a \times b \simeq \overline{\overline{a \times b}} \rightarrow (\exists f) [f : a \times b \xrightarrow[1-1]{\text{onto}} \overline{\overline{a \times b}}]$.

Since $a \neq 0 \wedge b \neq 0$, $(\exists x \in a) \wedge (\exists y \in b)$. Then

$$a \simeq f^{-1}(\overline{\overline{a \times \{y\}}}) \subseteq \overline{\overline{a \times b}} \wedge b \simeq f^{-1}(\overline{\overline{\{x\} \times b}}) \subseteq \overline{\overline{a \times b}}.$$

From Theorem 10.9, $a \simeq \bar{a} \wedge b \simeq \bar{b}$.

Conversely if $a \simeq \bar{a} \wedge b \simeq \bar{b}$ then $(\exists f) [f : a \xrightarrow[1-1]{\text{onto}} \bar{a}] \wedge (\exists g) [g : b \xrightarrow[1-1]{\text{onto}} \bar{b}]$. If

$$F(x, y) = \bar{a}g^{-1}y + f^{-1}x \quad \text{for } x \in a \wedge y \in b$$

then $F : a \times b \xrightarrow{1-1} \bar{a}\bar{b}$. Consequently $a \times b \simeq F''(a \times b) \subseteq \bar{a}\bar{b}$ and hence $a \times b \simeq \overline{a \times b}$.

Theorem 10.16. $\overline{\bar{a}} = \bar{a}$.

Proof. If $\bar{a} = 0$ then $\overline{\bar{a}} = \bar{0} = 0 = \bar{a}$. If $\bar{a} \neq 0$ then, by Theorem 10.10, $a \simeq \bar{a}$. Therefore $\alpha \simeq a \iff \alpha \simeq \bar{a}$ i.e. $\{\alpha \mid \alpha \simeq a\} = \{\alpha \mid \alpha \simeq \bar{a}\}$. Hence $\overline{\bar{a}} = \bar{a}$.

Theorem 10.17. $m \simeq n \rightarrow m = n$.

Proof (by induction on n). If $m \simeq 0$ then $m = 0$. If $(\forall m) [m \simeq n \rightarrow m = n]$ then, since $m \simeq (n + 1) \rightarrow m \neq 0$, $(\exists p) [m = p + 1]$. But $p + 1 \simeq n + 1 \rightarrow p \simeq n$ (Exercise 5, above). Then, from the induction hypothesis, $p = n$ and hence, $m = p + 1 = n + 1$.

Corollary 10.18.

- 1) $n \not\simeq n + 1$.
- 2) $\neg(\exists f) [f : (n + 1) \xrightarrow{1-1} n]$.

The proofs are left to the reader.

Theorem 10.19. $\bar{n} = n$.

Proof. Since $\bar{n} \leq n$ it follows that $\bar{n} \in \omega$. Since $\bar{n} \simeq n$ we have from Theorem 10.17 that $\bar{n} = n$.

Theorem 10.20. $\alpha \simeq n \rightarrow \alpha = n$.

Proof. If $\alpha \geq \omega$ then $n < \alpha$; hence $n + 1 \leq \alpha$. Since $\alpha \simeq n \subset n + 1$ it follows from the Cantor-Bernstein-Schröder Theorem that $\alpha \simeq n + 1$. Then $n \simeq n + 1$ and hence $n = n + 1$. From this contradiction we conclude that $\alpha \simeq n \rightarrow \alpha < \omega$. Then, by Theorem 10.17, $\alpha \simeq n \rightarrow \alpha = n$.

Remark. We next introduce predicates “Fin” and “Inf” for finite and infinite sets. Note that Inf is different from inf.

Definition 10.21.

- 1) $\text{Fin}(a) \xleftrightarrow{\Delta} (\exists n) [a \simeq n]$.
- 2) $\text{Inf}(a) \xleftrightarrow{\Delta} \neg \text{Fin}(a)$.

Exercises. Prove the following.

- 1) $\text{Fin}(a) \rightarrow \text{Fin}(a \cup \{b\})$.
- 2) $\text{Fin}(a) \rightarrow \text{Fin}(a - \{b\})$.
- 3) $\text{Fin}(a) \rightarrow \text{Fin}(\{b\} \times a)$.
- 4) $\text{Fin}(n)$.

- 5) $\text{Inf}(a) \rightarrow \bar{a} = a \cup \{a\}.$
- 6) $\text{Inf}(a) \wedge a \simeq b \rightarrow \text{Inf}(b).$
- 7) $a \cap b = 0 \wedge a \simeq \alpha \wedge b \simeq \beta \rightarrow a \cup b \simeq \alpha + \beta.$
- 8) $\alpha \times \beta \simeq \alpha\beta.$
- 9) $\text{Fin}(a) \rightarrow (\forall x) [x \subset a \rightarrow x \not\subset a].$
- 10) $a \simeq a \cup \{a\} \leftrightarrow (\exists x) [x \subset a \wedge x \simeq \omega].$

Theorem 10.22. $a \subseteq b \wedge b \simeq \bar{b} \rightarrow \bar{a} \leq \bar{b}.$

Proof. $b \simeq \bar{b} \rightarrow (\exists f) [f : b \xrightarrow[\text{onto}]{1-1} \bar{b}].$ Since $a \subseteq b, f''a \subseteq \bar{b}.$ Then $(\exists \beta) (\exists h) [h \text{ Isom}_{E,E}(\beta, f''a)].$ Consequently $a \simeq \beta$ and hence $\bar{a} \leq \beta.$ But since h is a strictly monotone ordinal function

$$\alpha < \beta \rightarrow \alpha \leq h'\alpha \wedge h'\alpha \in f''b.$$

But $f''b = \bar{b}$ and \bar{b} is transitive hence $\beta \subseteq \bar{b}$ i.e. $\bar{a} \leq \bar{b}.$

Corollary 10.23. $a \subseteq \alpha \rightarrow \bar{a} \leq \bar{\alpha}.$

Proof. Theorems 10.11 and 10.22.

Theorem 10.24. $\text{Fin}(a) \wedge b \subseteq a \rightarrow \text{Fin}(b).$

Proof. $\text{Fin}(a) \rightarrow (\exists n) [n \simeq a].$ Then $a \simeq \bar{a} \in \omega.$ By Theorem 10.22, $b \subseteq a \rightarrow \bar{b} \leq \bar{a}$ i.e. $\bar{b} \in \omega.$ Furthermore by Theorem 10.9, $b \simeq \bar{b}$ hence b is finite.

Theorem 10.25. $(a \times b) \simeq \overline{\overline{a \times b}} \wedge b \neq 0 \rightarrow \bar{a} \leq \overline{\overline{a \times b}}.$

Proof. $b \neq 0 \rightarrow \exists y \in b.$ If

$$f'x = \langle x, y \rangle, \quad x \in a$$

then $f : a \xrightarrow{1-1} a \times b.$ Then $a \simeq f''a \subseteq a \times b.$ Since $a \times b \simeq \overline{\overline{a \times b}}$ it follows from Theorem 10.22 that

$$\bar{a} = \overline{\overline{f''a}} \leq \overline{\overline{a \times b}}.$$

Theorem 10.26. $a \simeq \bar{a} \wedge \mathcal{U}_n(A) \rightarrow \overline{\overline{A''a}} \leq \bar{a}.$

Proof. Since $a \simeq \bar{a}, (\exists h) [h : \bar{a} \xrightarrow[\text{onto}]{1-1} a].$ Then $A \circ h : \bar{a} \rightarrow A''a.$ But $x \in A''a$ implies that $(\exists b \in a) [x = A'b]$ and $b \in a$ implies $(\exists \beta \in \bar{a}) [b = h'\beta].$ Thus $x \in A''a \rightarrow (\exists \beta \in \bar{a}) [x = (A \circ h)'\beta]$ i.e. $A \circ h : \bar{a} \xrightarrow[\text{onto}]{} A''a.$ Furthermore if

$$B = \{\beta \in \bar{a} \mid (\forall \alpha < \beta) [A'h'\alpha \neq A'h'\beta]\}$$

then $B \subseteq \bar{a}$ i.e. B is a set and $\bar{B} \leq \bar{a}.$ Also $\beta \in B \wedge \gamma \in B \wedge A'h'\beta = A'h'\gamma \rightarrow \beta = \gamma.$ Finally if $(\forall x \in A''a) (\exists \beta \in \bar{a}) [x = A'h'\beta].$ There is then a smallest such $\beta.$ For this β we have $\alpha < \beta \rightarrow A'h'\alpha \neq A'h'\beta.$ Therefore $\beta \in B$ i.e.

(A \circ h) $\Gamma B : B \xrightarrow{1-1}_{\text{onto}} A \text{ " } a$. Then

$$\overline{\overline{A \text{ " } a}} = \overline{\overline{B}} \leq \overline{\overline{a}}.$$

Theorem 10.27. $\overline{\overline{a}} > 0 \rightarrow [(\exists f) [f : a \xrightarrow{\text{onto}} b] \leftrightarrow 0 < \overline{\overline{b}} \leq \overline{\overline{a}}]$.

Proof. From Theorem 10.26, $a \simeq \overline{\overline{a}} \wedge [f : a \xrightarrow{\text{onto}} b] \rightarrow \overline{\overline{b}} = \overline{\overline{f \text{ " } a}} \leq \overline{\overline{a}}$. Also $\overline{\overline{a}} > 0 \rightarrow a \neq 0$. Hence $f \text{ " } a \neq 0$. Therefore $\overline{\overline{b}} > 0$.

Conversely if $0 < \overline{\overline{b}} \leq \overline{\overline{a}}$ then $b \simeq \overline{\overline{b}} \wedge a \simeq \overline{\overline{a}}$. Furthermore

$$(\exists g) [g : b \xrightarrow{1-1} a]$$

and $\exists y \in b$. If

$$f \text{ " } x = g^{-1}(x) \quad x \in g \text{ " } b,$$

$$f \text{ " } x = y \quad x \in a - g \text{ " } b.$$

Then $f : a \xrightarrow{\text{onto}} b$.

Theorem 10.28. $\overline{\overline{a}} > 1 \wedge \overline{\overline{b}} > 1 \rightarrow \overline{\overline{a \cup b}} \leq \overline{\overline{a \times b}}$.

Proof. Since $\overline{\overline{a}} > 0 \wedge \overline{\overline{b}} > 0$ it follows from Theorem 10.15 that $a \times b \simeq \overline{\overline{a \times b}}$. Since $\overline{\overline{a}} > 1 \wedge \overline{\overline{b}} > 1$ it follows that

$$(\exists x_1 \in a) (\exists x_2 \in a) [x_1 \neq x_2] \wedge (\exists y_1 \in b) (\exists y_2 \in b) [y_1 \neq y_2].$$

If

$$F \text{ " } \langle x_2, y_1 \rangle = y_1,$$

$$F \text{ " } \langle x_2, y_2 \rangle = x_2,$$

$$F \text{ " } \langle x, y_1 \rangle = x \text{ if } x \neq x_2,$$

$$F \text{ " } \langle x, y \rangle = y \text{ otherwise.}$$

then $F : a \times b \rightarrow a \cup b$; hence by Theorem 10.26

$$\overline{\overline{a \cup b}} = \overline{\overline{F \text{ " } (a \times b)}} \leq \overline{\overline{a \times b}}.$$

Theorem 10.29. $\text{Fin}(a) \wedge \text{Fin}(b) \rightarrow \text{Fin}(a \cup b) \wedge \text{Fin}(a \times b)$.

Proof. $\text{Fin}(a) \wedge \text{Fin}(b) \rightarrow (\exists m) [m \simeq a] \wedge (\exists n) [n \simeq b]$. Then

$$a \times b \simeq m \times n \simeq mn$$

(Exercise 8, above). Therefore $a \times b$ is finite.

If $m > 1 \wedge n > 1$ then by Theorem 10.28, $\overline{\overline{a \cup b}} \leq mn$ i.e. $\overline{\overline{a \cup b}} \in \omega$. Furthermore since $a \simeq \overline{\overline{a}}$ and $b \simeq \overline{\overline{b}}$ it follows from Theorem 10.15 that $a \cup b \simeq \overline{\overline{a \cup b}}$, hence $a \cup b$ is finite. That $a \cup b$ is finite for $m \leq 1 \vee n \leq 1$ we leave to the reader.

Theorem 10.30. $\text{Fin}(\alpha) \leftrightarrow \alpha \in \omega$.

Proof. $\text{Fin}(\alpha) \rightarrow (\exists n) [n \simeq \alpha]$. From Theorem 10.20, $\alpha \simeq n \rightarrow \alpha = n$ i.e. $\alpha \in \omega$. Since $\alpha \simeq \alpha$, $\alpha \in \omega \rightarrow \text{Fin}(\alpha)$.

Theorem 10.31. $\bar{\alpha} < \bar{\beta} \leftrightarrow \alpha < \beta$.

Proof. Since $\bar{\alpha} \leq \alpha$ it follows that $\alpha < \bar{\beta} \rightarrow \bar{\alpha} < \bar{\beta}$. Also $\bar{\beta} \leq \alpha \rightarrow \bar{\beta} = \overline{(\bar{\beta})} \leq \bar{\alpha}$.

Theorem 10.32. $\alpha > 1 \rightarrow \overline{\alpha + 1} \leq \overline{\alpha \times \alpha}$.

Proof. Since $\alpha \times \alpha$ is well ordered by R_0 there is an ordinal number equivalent to $\alpha \times \alpha$. Hence $\alpha \times \alpha \simeq \overline{\alpha \times \alpha}$. If

$$\begin{aligned} g'\beta &= \langle 0, \beta \rangle \quad \text{for } \beta < \alpha, \\ g'\alpha &= \langle 1, 0 \rangle \end{aligned}$$

then $g: (\alpha + 1) \xrightarrow{1-1} \alpha \times \alpha$ and hence $(\alpha + 1) \simeq g''(\alpha + 1) \subseteq \alpha \times \alpha$. From Theorem 10.22

$$\overline{\alpha + 1} \leq \overline{\alpha \times \alpha}.$$

Theorem 10.33. $\alpha \geq \omega \rightarrow \overline{\alpha \times \alpha} = \bar{\alpha}$.

Proof (by transfinite induction on α). As our induction hypothesis we have

$$\mu < \alpha \rightarrow [\mu < \omega \vee \overline{\mu \times \mu} = \bar{\mu}].$$

By Corollary 10.23, $\mu < \alpha \rightarrow \bar{\mu} \leq \bar{\alpha}$. If $(\exists \mu < \alpha) [\bar{\mu} = \bar{\alpha}]$ then $\mu \simeq \alpha$ and

$$\overline{\alpha \times \alpha} = \overline{\mu \times \mu} = \bar{\mu} = \bar{\alpha}.$$

If $(\forall \mu < \alpha) [\bar{\mu} < \bar{\alpha}]$ then since $\mu < \omega \vee \overline{\mu + 1} = \bar{\mu}$ it follows from Theorem 10.31 that $\mu + 1 < \bar{\alpha} \leq \alpha$. From our induction hypothesis it then follows that

$$\alpha > \mu \geq \omega \rightarrow \overline{(\mu + 1) \times (\mu + 1)} = \overline{\mu + 1}.$$

Recall the relation R_0 of Definition 7.55. By Theorem 7.56 R_0 well orders On^2 . Consequently there is an order isomorphism J_0 such that $J_0 \text{ Isom}_{R_0, E}(\text{On}^2, \text{On})$. We wish to show that $J_0''(\alpha \times \alpha) \subseteq \bar{\alpha}$. First we recall that an order isomorphism maps initial segments into initial segments:

$$J_0'' R_0^{-1} \{ \langle \beta, \gamma \rangle \} = E^{-1} \{ J_0' \langle \beta, \gamma \rangle \} = J_0' \langle \beta, \gamma \rangle.$$

Then since J_0 is one-to-one

$$\overline{J_0' \langle \beta, \gamma \rangle} = \overline{J_0'' R_0^{-1} \{ \langle \beta, \gamma \rangle \}} = \overline{R_0^{-1} \{ \langle \beta, \gamma \rangle \}}.$$

But if $\langle \beta, \gamma \rangle \in \alpha \times \alpha$ and if $\mu = \max(\beta, \gamma)$ then $\mu < \alpha$ and

$$\begin{aligned} \langle \eta, \theta \rangle \in R_0^{-1} \{ \langle \beta, \gamma \rangle \} &\rightarrow \langle \eta, \theta \rangle R_0 \langle \beta, \gamma \rangle \\ &\rightarrow \max(\eta, \theta) \leq \mu \\ &\rightarrow \eta \leq \mu \wedge \theta \leq \mu \\ &\rightarrow \langle \eta, \theta \rangle \in (\mu + 1) \times (\mu + 1) \end{aligned}$$

i.e. $R_0^{-1} \{ \langle \beta, \gamma \rangle \} \subseteq (\mu + 1) \times (\mu + 1)$.

Since $\mu < \alpha$ it follows from the induction hypothesis and Theorem 10.22 that if $\mu \geq \omega$ then

$$\overline{\overline{R_0^{-1} \{ \langle \beta, \gamma \rangle \}}} \leq \overline{(\mu + 1) \times (\mu + 1)} = \overline{\mu + 1} < \overline{\alpha}.$$

Therefore $\overline{J_0' \langle \beta, \gamma \rangle} < \overline{\alpha}$ and hence $J_0' \langle \beta, \gamma \rangle < \overline{\alpha}$.

If $\mu < \omega$ then $(\mu + 1) \times (\mu + 1)$ is finite. Hence $R_0^{-1} \{ \langle \beta, \gamma \rangle \}$ is finite and finally $J_0' \langle \beta, \gamma \rangle$ is finite. Since $\alpha \geq \omega$, $J_0' \langle \beta, \gamma \rangle < \overline{\alpha}$.

Thus

$$J_0''(\alpha \times \alpha) \subseteq \overline{\alpha}.$$

Since J_0 is one-to-one

$$\overline{\alpha \times \alpha} = \overline{J_0''(\alpha \times \alpha)} \subseteq \overline{\alpha}.$$

But from Theorem 10.32

$$\overline{\alpha} = \overline{\alpha + 1} \leq \overline{\alpha \times \alpha}.$$

Therefore $\overline{\alpha \times \alpha} = \overline{\alpha}$.

Theorem 10.34. $\overline{a} \geq \omega \rightarrow \overline{a \times a} = \overline{a}$.

Proof. Since $a \simeq \overline{a}$ we have $a \times a \simeq \overline{a} \times \overline{a}$. Then from Theorem 10.33

$$\overline{a \times a} = \overline{\overline{a} \times \overline{a}} = \overline{a}.$$

Theorem 10.35. $\overline{a} \geq \omega \wedge \overline{b} > 0 \rightarrow \overline{a \times b} = \overline{a \cup b} = \max(\overline{a}, \overline{b})$.

Proof. If $\alpha = \max(\overline{a}, \overline{b})$ then since $a \simeq \overline{a} \wedge b \simeq \overline{b}$

$$a \times b \simeq \overline{a} \times \overline{b} \subseteq \alpha \times \alpha.$$

Then

$$\overline{a \times b} \leq \overline{\alpha \times \alpha} = \alpha.$$

If $\overline{b} = 1$ then $\overline{a} \geq \omega \rightarrow a \times b \simeq a \cup b$ hence $\overline{a \times b} = \overline{a \cup b} = \overline{a} = \max(\overline{a}, \overline{b})$.

If $\overline{b} > 1$ then by Theorem 10.28, $\overline{a \cup b} \leq \overline{a \times b}$. Also since $\overline{a} > 0 \wedge \overline{b} > 0$, $a \cup b \simeq \overline{a \cup b}$ and hence $\overline{a} \leq \overline{a \cup b} \wedge \overline{b} \leq \overline{a \cup b}$ i.e. $\max(\overline{a}, \overline{b}) \leq \overline{a \cup b} \leq \overline{a \times b} \leq \max(\overline{a}, \overline{b})$.

Remark. We next introduce the class of cardinal numbers.

Definition 10.36. $N = \{ \alpha \mid (\exists x) [\alpha = \overline{x}] \}$.

Theorem 10.37. $N \subseteq \text{On}$.

Proof. Definition 10.36.

Theorem 10.38. $\alpha \in N \iff \alpha = \bar{\alpha}$.

Proof. If $\alpha = \bar{\alpha}$ then by Definition 10.36, $\alpha \in N$. If $\alpha \in N$ then $(\exists x) [\alpha = \bar{x}]$. From Theorem 10.11 it then follows that $\alpha = 0 \vee x \simeq \bar{x}$. If $\alpha = 0$ then $\bar{\alpha} = \bar{0} = 0 = \alpha$. If $x \simeq \bar{x}$ then since $\alpha \simeq \bar{\alpha}$ we have that $x \simeq \bar{\alpha}$. Therefore $\bar{x} \leq \bar{\alpha}$. But $\bar{x} = \alpha$. Hence $\alpha \leq \bar{\alpha}$. From Corollary 10.13 $\bar{\alpha} \leq \alpha$. Consequently $\alpha = \bar{\alpha}$.

Theorem 10.39. $\omega \subseteq N$.

Proof. Theorems 10.38 and 10.19.

Remark. Our next objective is to prove that N is a proper class. This is easily proved from the following theorem.

Theorem 10.40.

$$(\forall a) (\exists \beta \in N) [\beta = \{\alpha \mid (\exists f) [f : \alpha \xrightarrow{1-1} a]\} \wedge \neg (\exists f) [f : \beta \xrightarrow{1-1} a]].$$

Proof. For each $r \subseteq a \times a$ and for each $x \subseteq a$ if $r \text{ We } x$ then

$$(\exists ! \beta) (\exists ! f_{r,x}) [f_{r,x} \text{ Isom}_{r,E}(x, \beta)].$$

Let

$$\begin{aligned} F' \langle r, x \rangle &= f_{r,x}'' x \quad \text{if } r \text{ We } x \\ &= 0 \quad \text{otherwise,} \end{aligned}$$

then F' is a function on $\mathcal{P}(a) \times \mathcal{P}(a) \times \mathcal{P}(a) \triangleq \mathcal{P}^3(a)$. Therefore $F'' \mathcal{P}^3(a)$ is a set, and indeed a set of ordinals.

If $b = F'' \mathcal{P}^3(a)$ then

$$\begin{aligned} y \in b &\iff (\exists r \subseteq a \times a) (\exists x \subseteq a) [y = F' \langle r, x \rangle] \\ &\iff y = 0 \vee (\exists r \subseteq a \times a) (\exists x \subseteq a) [r \text{ We } x \wedge y = f_{r,x}'' x] \\ &\iff y \in \text{On} \wedge (\exists f) [f : y \xrightarrow{1-1} a], \end{aligned}$$

i.e. $b = \{\alpha \mid (\exists f) [f : \alpha \xrightarrow{1-1} a]\}$.

If $\gamma < \beta \wedge \beta \in b$ then $(\exists f) [f : \beta \xrightarrow{1-1} a]$. Then $f \upharpoonright \gamma : \gamma \xrightarrow{1-1} a$; hence $\gamma \in b$. Thus b is a transitive set of ordinals. Therefore b is an ordinal i.e. $(\exists \beta) [\beta = \{\alpha \mid (\exists f) [f : \alpha \xrightarrow{1-1} a]\}]$. Furthermore if $\gamma \simeq \beta$ and $\gamma < \beta$ then $(\exists g) [g : \beta \xrightarrow{1-1} \gamma]$ and $(\exists f) [f : \gamma \xrightarrow{1-1} a]$. Then $f \circ g : \beta \xrightarrow{1-1} a$ i.e., $\beta \in \beta$. From this contradiction we conclude that

$$\gamma \simeq \beta \rightarrow \beta \leq \gamma$$

hence $\bar{\beta} = \beta$ i.e., $\beta \in N$.

Furthermore if $(\exists f) [f : \beta \xrightarrow{1-1} a]$ then again $\beta \in \beta$, hence

$$\neg (\exists f) [f : \beta \xrightarrow{1-1} a].$$

Theorem 10.41. $\mathcal{P}_r(N)$.

Proof. If $(\exists a) [a = N]$ then since $N \subseteq \text{On}$, $\cup(a) \in \text{On}$. If $\alpha = \cup(a)$ then by Theorem 10.40

$$(\exists \beta \in a) [\beta = \{\delta \mid (\exists f) [f : \delta \xrightarrow{1-1} \alpha]\} \wedge \neg (\exists f) [f : \beta \xrightarrow{1-1} \alpha]].$$

But $\beta \in a \rightarrow \beta \subseteq \alpha$ hence $(\exists f) [f : \beta \xrightarrow{1-1} \alpha]$.

From this contradiction we conclude that N is a proper class.

Definition 10.42. $N' \triangleq N - \omega$.

Theorem 10.43. $\mathcal{P}_r(N')$.

Proof. $N = N' \cup \omega$. Therefore $\mathcal{M}(N') \rightarrow \mathcal{M}(N)$.

Remark. Since N' is a proper class of ordinals it is order isomorphic to On (Corollary 7.50).

Definition 10.44. $\aleph \text{ Isom}_{E,E}(\text{On}, N')$.

Definition 10.45. $\aleph_\alpha = \aleph^{\alpha}$.

Theorem 10.46.

- 1) $\alpha < \aleph_\beta \iff \bar{\alpha} < \aleph_\beta$.
- 2) $\overline{\aleph_\alpha \times \aleph_\alpha} = \aleph_\alpha$.
- 3) $0 < \gamma < \aleph_{\alpha+1} \rightarrow (\exists f) [f : \aleph_\alpha \xrightarrow{\text{onto}} \gamma]$.

Proof. Theorems 10.31, 10.33, and 10.27 respectively.

Exercises. *Prove the following.*

- 1) $\omega \in N$.
- 2) $\alpha \leq \aleph_\alpha$.
- 3) $\aleph_0 = \omega$.
- 4) $N' \subseteq K_{\text{II}}$.

Definition 10.47. $a^b \triangleq \{f \mid f : b \rightarrow a\}$.

Theorem 10.48. $\mathcal{M}(a^b)$.

Proof. $a^b \subseteq \mathcal{P}(b \times a)$.

Theorem 10.49. $2^a \simeq \mathcal{P}(a)$.

Proof. We define a function h on 2^a in the following way

$$(\forall f \in 2^a) [h^{\ast} f = \{x \in a \mid f^{\ast} x = 1\}].$$

Then $h : 2^a \rightarrow \mathcal{P}(a)$. If $b \in \mathcal{P}(a)$ and if

$$\begin{aligned} f'x &= 1 & \text{if } x \in b \\ &= 0 & \text{if } x \in a - b \end{aligned}$$

then $f \in 2^a$ and $h'f = b$. Thus h is an onto mapping.

Furthermore if $h'f = h'g$, $f \in 2^a \wedge g \in 2^a$, then

$$\{x \in a \mid f'x = 1\} = \{x \in a \mid g'x = 1\}.$$

Thus if $b = \{x \in a \mid f'x = 1\}$ and $x \in b$ then $f'x = g'x = 1$. If $x \in a - b$ then $f'x = g'x = 0$, i.e., $f = g$. Then $h : 2^a \xrightarrow[\text{onto}]{1-1} \mathcal{P}(a)$, that is, $2^a \simeq \mathcal{P}(a)$.

Theorem 10.50. $(a^b)^c \simeq a^{b \times c}$.

Proof. If

$$F = \{\langle f, g \rangle \mid f \in (a^b)^c \wedge g \in a^{b \times c} \wedge (\forall y \in b) (\forall x \in a) [g(y, x) = (f'x)'y]\}$$

then $F : (a^b)^c \rightarrow a^{b \times c}$. If $g \in a^{b \times c}$ and if $\forall x \in c$

$$\tilde{f}'_x y = g(y, x), \quad y \in b$$

then $\tilde{f}'_x \in a^b$. If

$$f'x = \tilde{f}'_x, \quad x \in c$$

then $f \in (a^b)^c$ and $g(y, x) = (f'x)'y$. Therefore

$$F'f = g.$$

Thus F is an onto mapping. If $F'f_1 = F'f_2$, $f_1 \in (a^b)^c$, and $f_2 \in (a^b)^c$ then

$$(\forall x \in c) (\forall y \in b) [(f_1'x)'y = (f_2'x)'y],$$

$$(\forall x \in c) [f_1'x = f_2'x],$$

$$f_1 = f_2.$$

Then $F : (a^b)^c \xrightarrow[\text{onto}]{1-1} a^{b \times c}$ i.e., $(a^b)^c \simeq a^{b \times c}$.

Remark. From the one-to-one correspondences we have discovered many interesting properties of sets. The strictly monotone ordinal functions provide another interesting avenue of investigation. We wish to consider strictly monotone ordinal functions defined on an ordinal β and whose range is a subset of a larger ordinal α . We require that every element of α either be in the range of the function or be smaller than some ordinal that is in the range. Under these circumstances we say that α is cofinal with β .

Definition 10.51.

$$\text{cof}(\alpha, \beta) \xleftarrow{\Delta} \beta \leq \alpha \wedge (\exists f) [\text{Smo}(f) \wedge f : \beta \rightarrow \alpha \wedge (\forall \gamma < \alpha) (\exists \delta < \beta) [f'\delta \geq \gamma]].$$

Theorem 10.52.

- 1) $\text{cof}(\alpha, \alpha)$.
- 2) $\text{cof}(\alpha, \beta) \wedge \text{cof}(\beta, \gamma) \rightarrow \text{cof}(\alpha, \gamma)$.

Theorem 10.53.

- 1) $\text{cof}(\alpha, 0) \leftrightarrow \alpha = 0$.
- 2) $\alpha \in K_I \wedge \beta \in K_I \wedge 0 < \beta \leq \alpha \rightarrow \text{cof}(\alpha, \beta)$.

Corollary 10.54. $1 \leq \alpha \in K_I \rightarrow \text{cof}(\alpha, 1)$.

The proofs are left to the reader.

Theorem 10.55. $\text{cof}(\alpha, \beta) \rightarrow [\alpha \in K_{II} \leftrightarrow \beta \in K_{II}]$.

Proof. By hypothesis $\beta \leq \alpha$. Therefore $\alpha = 0 \rightarrow \beta = 0$. Furthermore

$$(\exists f) [\text{Smo}(f) \wedge [f: \beta \rightarrow \alpha] \wedge (\forall \gamma < \alpha) (\exists \delta < \beta) [f'\delta \geq \gamma]].$$

If $\alpha \neq 0 \wedge \alpha \in K_I$ then $(\exists \gamma) [\alpha = \gamma + 1]$. Since $\gamma < \alpha$, $(\exists \delta < \beta) [f'\delta \geq \gamma]$. But f is strictly monotone and $\alpha - (\gamma + 1) = 0$. Therefore $\beta - (\delta + 1) = 0$ i.e., $\beta = \delta + 1$.

If $\alpha \in K_{II}$ then $\beta \neq 0$. If $(\exists \delta) [\beta = \delta + 1]$ then since f is strictly monotone $\tau < \delta$ implies that $f'\tau < f'\delta$, i.e., $\neg (\exists \tau < \beta) [f'\tau \geq f'\delta + 1]$. Since $f'\delta + 1 < \alpha$ this is a contradiction. Thus $\beta \in K_{II}$.

Theorem 10.56. $\beta < \alpha \wedge (\exists f) [f: \beta \rightarrow \alpha \wedge (\forall \gamma < \alpha) (\exists \delta < \beta) [f'\delta \geq \gamma]] \rightarrow (\exists \eta \leq \beta) [\text{cof}(\alpha, \eta)]$.

Proof. If $a = \{\delta < \beta \mid (\forall \gamma < \delta) [f'\gamma < f'\delta]\}$ then $a \subseteq \beta$. Therefore

$$(\exists \eta \leq \beta) (\exists h) [h \text{ Isom}_{E,E}(\eta, a)].$$

If $g = f \circ h$ then $g: \eta \rightarrow \alpha$. If $\delta < \gamma < \eta$ then $h'\delta < h'\gamma \wedge h'\delta \in a \wedge h'\gamma \in a$. Then

$$f'h'\delta < f'h'\gamma$$

i.e., g is strictly monotone.

Since $(\forall \gamma < \alpha) (\exists \delta < \beta) [f'\delta \geq \gamma]$ it follows that there is a smallest such δ . This δ is in a and hence $(\exists v < \eta) [h'v = \delta]$. Then

$$\gamma \leq f'\delta = f'h'v = g'v.$$

Consequently $\text{cof}(\alpha, \eta)$.

Corollary 10.57. $\beta \leq \alpha \wedge \beta \simeq \alpha \rightarrow (\exists \eta \leq \beta) [\text{cof}(\alpha, \eta)]$.

Proof. $\beta \simeq \alpha \rightarrow (\exists f) [f: \beta \xrightarrow[\text{onto}]{1-1} \alpha]$. Then $(\forall \gamma < \alpha) (\exists \delta < \beta) [f'\delta = \gamma]$. By Theorem 10.56 $(\exists \eta \leq \beta) [\text{cof}(\alpha, \eta)]$.

Theorem 10.58.

- 1) $\alpha \in K_{\text{II}} \wedge \text{cof}(\alpha, \beta) \rightarrow (\exists f) [f \mathcal{F}_n \beta \wedge \alpha = \cup(f''\beta)].$
- 2) $\beta \in K_{\text{II}} \wedge f \mathcal{F}_n \beta \wedge \alpha = \cup(f''\beta) \wedge \text{Smo}(f) \rightarrow \text{cof}(\alpha, \beta).$

Theorem 10.59. $\alpha \in K_{\text{II}} \rightarrow \text{cof}(\aleph_\alpha, \alpha).$

The proofs are left to the readers.

Theorem 10.60. $\text{cof}(\alpha, \beta) \wedge \text{cof}(\alpha, \gamma) \wedge \gamma \leq \beta \rightarrow (\exists \eta \leq \gamma) [\text{cof}(\beta, \eta)].$

Proof. Since α is cofinal with β and with γ

$$(\exists f) [f: \beta \rightarrow \alpha \wedge (\forall \tau < \alpha) (\exists \delta < \beta) [f'\delta \geq \tau] \wedge \text{Smo}(f)],$$

$$(\exists g) [g: \gamma \rightarrow \alpha \wedge (\forall \tau < \alpha) (\exists \delta < \gamma) [g'\delta \geq \tau]].$$

In particular, since $\delta < \gamma$ implies $g'\delta < \alpha$, $(\exists \tau < \beta) [f'\tau \geq g'\delta]$. There is then a smallest such τ . If

$$F'\delta = \mu_\tau(f'\tau \geq g'\delta), \quad \delta < \gamma$$

then $F: \gamma \rightarrow \beta$. Furthermore $\delta < \beta \rightarrow \delta \leq f'\delta < \alpha$. Then $(\exists \tau < \gamma) [g'\tau \geq f'\delta]$. Since f is strictly monotone $v < \delta \rightarrow f'v < f'\delta \leq g'\tau$ i.e., the smallest ordinal for which $f'v \geq g'\tau$ is greater than or equal to δ . Thus $F'\tau \geq \delta$. Then from Theorem 10.56 $(\exists \eta \leq \gamma) [\text{cof}(\beta, \eta)]$.

Exercises. *Prove the following.*

- 1) $\text{cof}(\aleph_\omega, \omega).$
- 2) $\text{cof}(\aleph_{\aleph_\omega}, \omega).$
- 3) If $r \text{ We } a \wedge s \text{ We } b \wedge R = \{ \langle f, g \rangle \in b^a \times b^a \mid (\exists x \in a) [f \upharpoonright r^{-1}\{x\} = g \upharpoonright r^{-1}\{x\}] \wedge (f'x) s (g'x) \}$ then $R \text{ We } b^a$.

Remark. If α is cofinal with β and $\beta < \alpha$ then α can be “reached” by a mapping from “below” in a sense made clear by the definition of cofinality. In particular note that ω cannot be reached by a strictly monotone ordinal function from below. That is ω is not cofinal with any finite ordinal. Thus cofinality describes a sense in which ω is much, much larger than any finite ordinal. Are there other large ordinals such as ω , large in the sense that they are not cofinal with any smaller ordinal. Corollary 10.54 assures us that such an ordinal must be a limit ordinal. To assist us in our efforts to answer this question we introduce a symbol for the character of cofinality of α by which we mean the smallest ordinal cofinal with α .

Definition 10.61. $\text{cf}(\alpha) = \mu_\beta(\text{cof}(\alpha, \beta)).$

Exercises. Prove the following.

- 1) $\text{cf}(\alpha) \leq \alpha$.
- 2) $\text{cf}(0) = 0$.
- 3) $\text{cf}(\alpha + 1) = 1$.
- 4) $\text{cf}(\omega) = \omega$.
- 5) $\text{cof}(\alpha, \text{cf}(\alpha))$.

Theorem 10.62. $\text{cf}(\alpha) \in N$.

Proof. If $\beta = \text{cf}(\alpha)$ and if $\gamma \simeq \beta$ then by Theorem 10.52 and Corollary 10.57

$$(\exists \eta \leq \gamma) \text{cof}(\alpha, \eta).$$

Therefore $\gamma \simeq \beta \rightarrow \beta \leq \gamma$. But this means that $\beta = \bar{\beta}$ i.e., $\text{cf}(\alpha) \in N$.

Remark. Theorem 10.62 assures us that if there is an ordinal that is large, in the sense that it is not cofinal with any smaller ordinal, that ordinal is a cardinal number.

Theorem 10.63. $\alpha \in N' \rightarrow \text{cf}(\alpha) \in N'$.

Proof. $\alpha \in N' \rightarrow \alpha \in K_{II}$. But since $\text{cof}(\alpha, \text{cf}(\alpha))$ it follows from Theorem 10.55 that $\text{cf}(\alpha) \in K_{II}$ i.e., $\text{cf}(\alpha) \notin \omega$. From Theorem 10.62, $\text{cf}(\alpha) \in N'$.

Theorem 10.64. $\alpha \in K_{II} \rightarrow \text{cf}(\alpha) = \text{cf}(\aleph_\alpha)$.

Proof. $\alpha \in K_{II} \rightarrow \text{cof}(\aleph_\alpha, \alpha)$. Since $\text{cof}(\alpha, \text{cf}(\alpha))$ we have $\text{cof}(\aleph_\alpha, \text{cf}(\alpha))$.

Therefore

$$\text{cf}(\aleph_\alpha) \leq \text{cf}(\alpha).$$

Furthermore $\text{cof}(\aleph_\alpha, \text{cf}(\aleph_\alpha))$. Then by Theorem 10.60

$$(\exists \eta \leq \text{cf}(\aleph_\alpha)) [\text{cof}(\text{cf}(\alpha), \eta)].$$

But $\text{cof}(\alpha, \text{cf}(\alpha)) \wedge \text{cof}(\text{cf}(\alpha), \eta) \rightarrow \text{cof}(\alpha, \eta)$. Therefore

$$\text{cf}(\alpha) \leq \eta \leq \text{cf}(\aleph_\alpha),$$

hence $\text{cf}(\alpha) = \text{cf}(\aleph_\alpha)$.

Definition 10.65.

- 1) $\text{Reg}(\alpha) \stackrel{\Delta}{\leftrightarrow} \alpha \in N' \wedge \text{cf}(\alpha) = \alpha$.
- 2) $\text{Sing}(\alpha) \stackrel{\Delta}{\leftrightarrow} \alpha \in N' \wedge \text{cf}(\alpha) < \alpha$.

Remark. Since $\omega < \aleph_\omega$ and $\text{cf}(\aleph_\omega) = \omega$ it follows that \aleph_ω is singular. From this example it is clear that $\aleph_{\omega 2}$, $\aleph_{\omega 3}$, etc. are all singular. Also $\text{cf}(\aleph_0) = \aleph_0$ hence \aleph_0 is regular. In the next section we will prove that \aleph_α is always regular if $\alpha \in K_I$. Does there exist a regular cardinal \aleph_α with $\alpha \in K_{II}$? Such a cardinal we call weakly inaccessible. The question of

whether or not there exists an inaccessible cardinal is as yet not completely resolved. $\text{Reg}(\alpha)$ and $\text{Sing}(\alpha)$ are read respectively “ α is a regular cardinal” and “ α is a singular cardinal”.

Definition 10.66.

- 1) $\text{Inacc}_w(\aleph_\alpha) \stackrel{\Delta}{\longleftrightarrow} \alpha \in K_{II} \wedge \text{Reg}(\aleph_\alpha)$.
- 2) $\text{Inacc}(\aleph_\alpha) \stackrel{\Delta}{\longleftrightarrow} \text{Inacc}_w(\aleph_\alpha) \wedge (\forall x) [\bar{x} < \aleph_\alpha \rightarrow \overline{\overline{\mathcal{P}(x)}} < \aleph_\alpha]$.

Theorem 10.67. $\text{Inacc}_w(\aleph_\alpha) \rightarrow \aleph_\alpha = \alpha$.

Proof. $\alpha \leq \aleph_\alpha$. On the other hand \aleph_α weakly inaccessible implies that $\text{cf}(\aleph_\alpha) = \aleph_\alpha \wedge \alpha \in K_{II}$. Since by Theorem 10.59, $\text{cof}(\aleph_\alpha, \alpha)$,

$$\aleph_\alpha = \text{cf}(\aleph_\alpha) \leq \alpha.$$

Then $\alpha = \aleph_\alpha$.

Remark. That there do exist cardinals for which $\aleph_\alpha = \alpha$ is easily established.

Theorem 10.68. $[f : a \rightarrow N] \rightarrow \cup(f''a) \in N$.

Proof. Since $\mathcal{D}(f)$ is a set $f''a$ is a set and indeed a set of ordinals. If $\beta = \cup(f''a)$ then $\bar{\beta} \leq \beta$. If $\bar{\beta} < \beta$ then $(\exists x \in a) [\beta < f'x \wedge f'x \in f''a]$. Then $f'x \subseteq \beta$ and since $f'x \in N$

$$f'x = \overline{\overline{f'x}} \leq \bar{\beta}.$$

This is a contradiction.

Corollary 10.69. $[f : a \rightarrow N] \wedge (\exists x \in a) [f'x \in N'] \rightarrow \cup(f''a) \in N'$.

Proof. By Theorem 10.68, $\cup(f''a) \in N$. Since $(\exists x \in a) [f'x \in N']$ and since $f'x \leq \cup(f''a)$ it follows that $\cup(f''a) \in N'$.

Theorem 10.70. $(\exists \alpha) [\alpha = \aleph_\alpha]$.

Proof. If $h(0) = \aleph_0$,

$$h(n+1) = \aleph_{h(n)}$$

then $h : \omega \rightarrow N'$. By Corollary 10.69, $\cup(h''\omega) \in N'$. Thus $(\exists \alpha) [\aleph_\alpha = \cup(h''\omega) \wedge \alpha \leq \aleph_\alpha]$. If $\alpha < \aleph_\alpha$ then $(\exists n) [\alpha < h'n]$. Then $\aleph_\alpha < \aleph_{h'n} = h(n+1) \leq \aleph_\alpha$. This is a contradiction. Hence $\alpha = \aleph_\alpha$.

11 The Axiom of Choice, the Generalized Continuum Hypothesis and Cardinal Arithmetic

In Section 10 we defined the cardinal number of a set, \bar{a} , to be the smallest ordinal that is equivalent to a . If no such ordinal exists then $\bar{a} = 0$. This definition has the advantage of connecting the theory of cardinal numbers to the properties of ordinals. A more traditional view is that \bar{a} is the equivalence class of sets equipollant to a .

To establish the equivalence of the two views of cardinal numbers we should prove that every set is equivalent to an ordinal. If this were the case then \bar{a} would be an element of the equivalence class determined by a . From the work of Cohen, to be studied later, we know that such a proof is not possible. We are then compelled to accept the possibility that a set exists that is not equivalent to an ordinal. Earlier we proved that a set is equivalent to an ordinal if and only if that set can be well ordered.

Thus the question of whether or not all equivalence classes are represented by the ordinals in N reduces to the question of whether or not every set can be well ordered. That every set can be well ordered is deducible from the following axiom.

Axiom of Choice (weak form)

$$(\forall a) (\exists f) (\forall x \in a) [x \neq 0 \rightarrow f'x \in x].$$

Axiom of Choice (strong form)

$$(\exists F) (\forall x) [x \neq 0 \rightarrow F'x \in x].$$

Remark. Since we cannot quantify on class symbols the strong form of the Axiom of Choice could only be expressed in ZF by adding a constant f_0 to the language together with the additional axioms

$$\mathcal{Fnc}(f_0)$$

$$(\forall x) [x \neq 0 \rightarrow f_0'x \in x].$$

The weak and strong forms of the Axiom of Choice are not equivalent for it can be shown that there are wffs in ZF that are provable with the strong form that are not provable with the weak form. Hereafter when we refer to the Axiom of Choice (AC) we will mean the weak form.

When we prove the relative consistence of ZF and ZF + AC we will in fact prove it for the strong form. Our method will be to construct a particular model of ZF in which we will be able to define a universal choice function f_0 . For the success of this project it is essential that all results depend only on the axioms of ZF and that no appeal be made to the AC or to any result that requires the AC. Certain theorems of this section must therefore be avoided. As a further reminder we will mark such theorems with an asterisk.

We will prove that the AC implies that every set can be well ordered and conversely that if every set can be well ordered then the AC holds. The assumption that every set can be well ordered we will call the Well Ordering Principle (WOP).

Well Ordering Principle. $(\forall a) (\exists r) [r \text{ We } a]$.

There are other principles equivalent to the AC. We present two others. Cantor assumed that every pair of sets a and b are comparable in the sense that either a is equivalent to a subset of b or b is equivalent to a subset of a :

Cantor's Law of Trichotomy.

$$(\forall a) (\forall b) [(\exists f) [f : a \xrightarrow{1-1} b] \vee (\exists f) [f : b \xrightarrow{1-1} a]].$$

The second principle that we will show to be equivalent to the AC is known as Zorn's Lemma. For its statement we must define "B is a chain in A" and "a is a maximal element of A."

Definition 11.1.

$$1) \quad \text{ch}(B, A) \xleftrightarrow{\Delta} B \subseteq A \wedge (\forall x \in B) (\forall y \in B) [x \subseteq y \vee y \subseteq x].$$

$$2) \quad \max_A(a) \xleftrightarrow{\Delta} a \in A \wedge (\forall x \in A) [a \subseteq x \rightarrow a = x].$$

$$\text{Zorn's Lemma: } a \neq 0 \wedge (\forall b) [\text{ch}(b, a) \rightarrow \cup(b) \in a] \rightarrow (\exists x) [\max_a(x)].$$

Theorem 11.2. *The following are equivalent*

- 1) AC,
- 2) WOP,
- 3) Zorn's Lemma,
- 4) Cantor's Law of Trichotomy.

Proof. 1) \rightarrow 2). From the AC

$$(\forall a) (\exists f) (\forall x \in \mathcal{P}(a)) [x \neq 0 \rightarrow f'x \in x].$$

If $G'x = f(a - \mathcal{W}(x))$ and $F \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [F'\alpha = G(F \uparrow \alpha)]$ then since $a - \mathcal{W}(x) \subseteq a$ it follows that if $a - F'\alpha \neq 0$ then $F'\alpha \in a - F'\alpha$. Hence by

Theorem 7.47

$$(\exists \alpha) [F \uparrow \alpha : \alpha \xrightarrow{\text{onto}} a].$$

Since $\alpha \simeq a$ the natural ordering of α induces a well ordering of a .

2) \rightarrow 3). If $(\forall a) (\exists r) [r \text{ We } a]$ and if $a \neq 0$ then $\exists x \in a$. We then define F by transfinite recursion such that $F(0) = x$, $F(\alpha + 1)$ is the r -minimal element of $\{y \in a \mid F' \alpha \subset y\}$ if such exists, $F(\alpha + 1) = F' \alpha$ otherwise and $F' \alpha = \bigcup_{\beta < \alpha} F' \beta$ if $\alpha \in K_{\parallel}$. By induction it then follows that $F' \alpha \subseteq F' \beta$ if

$\alpha < \beta$. Also by induction it follows that $F'' \text{On} \subseteq a$. Then $F'' \text{On}$ is a chain in a . Therefore if $b = \cup(F'' \text{On})$ then $b \in a$. Furthermore

$$(\exists \alpha) [F \uparrow \alpha : \alpha \xrightarrow{\text{onto}} F'' \text{On}] \quad \text{i.e.,} \quad b = \bigcup_{\beta < \alpha} F' \beta.$$

If $(\exists x \in a) [b \subset x]$ then there is an r -minimal such x . If $\alpha = \delta + 1$ then $b = F' \delta$ and $x = F' \alpha$. If $\alpha \in K_{\parallel}$ then $b = F' \alpha$ and $x = F(\alpha + 1)$. In either case $x \subseteq b$ which is a contradiction. Therefore b is a maximal element of a .

3) \rightarrow 4). If

$$A = \{f \subseteq a \times b \mid \langle x_1, y_1 \rangle \in f \wedge \langle x_2, y_2 \rangle \in f \rightarrow [x_1 = x_2 \leftrightarrow y_1 = y_2]\}$$

then A is a set. Furthermore if c is a chain in A then $\cup(c) \in A$. Therefore by 3) A has a maximal element, f . If $a - \mathcal{D}(f) \neq 0 \wedge b - \mathcal{W}(f) \neq 0$ then $(\exists x \in a - \mathcal{D}(f)) (\exists y \in a - \mathcal{W}(f)) [f \subset (f \cup \{\langle x, y \rangle\}) \in A]$. This is a contradiction. Therefore $[a - \mathcal{D}(f) = 0] \vee [b - \mathcal{W}(f) = 0]$ i.e., $[f : a \xrightarrow{1-1} b] \vee [f^{-1} : b \xrightarrow{1-1} a]$.

4) \rightarrow 2). If $b = \{\alpha \mid (\exists f) [f : \alpha \xrightarrow{1-1} a]\}$ then by Theorem 10.40 and 4)

$$(\exists h) [h : a \xrightarrow{1-1} b].$$

Since b is a set of ordinals $h'' a$ is a set of ordinals hence $(\exists \alpha) [h'' a \simeq \alpha]$. But $h : a \xrightarrow{\text{onto}} h'' a$. Therefore $a \simeq \alpha$. Then $(\exists r) [r \text{ We } a]$.

2) \rightarrow 1). If every set can be well ordered then in particular if $b = \cup(a)$, $(\exists r) [r \text{ We } b]$. Since $x \in a \rightarrow x \subseteq b$ it follows that if $x \neq 0$, x has an r -minimal element. If we define f on a in such a way that $f' x$ is the r -minimal element of x for $x \neq 0$ and $f' x = 0$ otherwise then $x \in a \wedge x \neq 0 \rightarrow f' x \in x$.

***Theorem 11.3.** $(\forall a) (\exists \alpha) [\alpha \simeq a]$.

Proof. From the AC it follows that every set can be well ordered. But every well ordered set is equivalent to an ordinal.

***Theorem 11.4.**

$$1) \quad (\forall a) [a \simeq \bar{a}].$$

$$2) \quad \bar{a} = 0 \leftrightarrow a = 0.$$

Proof. Theorems 11.3 and 10.9.

Remark. With the AC we can improve certain theorems of Section 10. For example, from Theorem 11.4 and Theorems 10.22, 10.25, and 10.26 we obtain

***Theorem 11.5.** $a \subseteq b \rightarrow \bar{a} \leq \bar{b}$.

***Theorem 11.6.** $b \neq 0 \rightarrow \bar{a} \leq \overline{a \times b}$.

***Theorem 11.7.** $\mathcal{U}^n(A) \rightarrow \overline{A^c a} \leq \bar{a}$.

Remark. Using the AC we can also provide simpler proofs and stronger statements of certain theorems. For example

Theorem 11.8 (Cantor-Bernstein-Schröder).

$$a \simeq c \subseteq b \wedge b \simeq d \subseteq a \rightarrow a \simeq b.$$

**Proof.* From *Theorem 11.5, $a \simeq c \subseteq b \rightarrow \bar{a} \leq \bar{b}$ and $b \simeq d \subseteq a \rightarrow \bar{b} \leq \bar{a}$. Therefore $\bar{a} = \bar{b}$.

***Theorem 11.9** (Cantor). $\bar{a} < \overline{\mathcal{P}(a)}$.

Proof. $\overline{\mathcal{P}(a)} \leq \bar{a}$ implies that $\mathcal{P}(a)$ is equivalent to a subset of a . This contradicts Corollary 10.5.

Exercises. Prove the following.

- 1) $\bar{a} = \bar{b} \rightarrow \overline{2^a} = \overline{2^b}$.
- 2) $\bar{a} = \bar{b} \rightarrow \bar{a^c} = \bar{b^c}$.
- 3) $\text{Inf}(a) \rightarrow \overline{2^{a \times a}} = \overline{2^a}$.
- 4) $\bar{a} < \bar{b} \rightarrow \overline{2^a} < \overline{2^b}$.
- 5) $\bar{a} < \bar{b} \rightarrow \bar{a^c} \leq \bar{b^c}$.
- 6) $b \neq 0 \rightarrow \bar{a} \leq \overline{a^b}$.
- 7) $\beta < \aleph_\alpha \wedge \gamma < \aleph_\alpha \rightarrow \beta + \gamma < \aleph_\alpha$.
- 8) $\beta < \aleph_\alpha \rightarrow \beta + \aleph_\alpha = \aleph_\alpha$.
- 9) $\overline{\aleph_\alpha + \aleph_\alpha} = \aleph_\alpha$.
- 10) $\neg(\exists f)[f : \aleph_\alpha \xrightarrow{\text{onto}} \aleph_{\alpha+1}]$.
- 11) $\alpha \in K_{II} \rightarrow \bigcup_{\beta < \alpha} \aleph_\beta \in N'$.
- 12) $\alpha \in K_{II} \wedge \beta < \aleph_\alpha \rightarrow (\exists \gamma < \alpha)[\beta < \aleph_\gamma]$.
- 13) $\alpha \in K_{II} \rightarrow \bigcup_{\beta < \alpha} \aleph_\beta = \aleph_\alpha$.
- 14) $a \neq 0 \wedge (\exists f)[f : a \xrightarrow{\text{onto}} b] \leftrightarrow \bar{b} \leq \bar{a} \wedge b \neq 0$.

***Theorem 11.10.** $(\forall x \in a) (\exists y) \varphi(x, y) \rightarrow (\exists f) (\forall x \in a) \varphi(x, f'x)$.

Proof. If $G'x = \mu_\alpha((\exists y) [\alpha = \text{rank}(y) \wedge \varphi(x, y)])$ then $G''a$ is a set of ordinals. If $\alpha = \cup(G''a) + 1$ then

$$(\forall x \in a) (\exists y) [\varphi(x, y)] \rightarrow (\exists y) [\varphi(x, y) \wedge \text{rank}(y) = G'x].$$

Furthermore $\text{rank}(y) = G'x \rightarrow \text{rank}(y) < \alpha$ i.e., $y \in R'\alpha$.

From the AC we can well order $R'\alpha$ and define $f'x$ to be the first element in $R'\alpha$ for which $\varphi(x, f'x)$. More precisely from the AC

$$(\exists \beta) (\exists h) [h : \beta \xrightarrow[1-1]{\text{onto}} R'\alpha].$$

Since $(\forall x \in a) (\exists y \in R'\alpha) [\varphi(x, y)]$ and since h is onto

$$(\forall x \in a) (\exists \gamma \in \beta) [\varphi(x, h'\gamma)].$$

There is then a smallest such γ . Therefore if

$$f = \{ \langle x, h'\gamma \rangle \in a \times h''\beta \mid \varphi(x, h'\gamma) \wedge (\forall \delta < \gamma) [\neg \varphi(x, h'\delta)] \}$$

then $f : a \rightarrow R'\alpha$ and $(\forall x \in a) [\varphi(x, f'x)]$.

Remark. *Theorem 11.10 is a generalization of the AC to classes: Given a collection of nonempty classes

$$A_x, x \in a$$

there is a choice function f such that $x \in a \rightarrow f'x \in A_x$.

***Theorem 11.11.** $(\forall x \in a) [\bar{x} \leq \bar{b}] \rightarrow \overline{\cup(a)} \leq \overline{a \times b}$.

Proof. $x \in a \rightarrow \bar{x} \leq \bar{b}$ it follows that

$$(\forall x \in a) (\exists \tilde{f}_x) [\tilde{f}_x : x \xrightarrow[1-1]{} b].$$

Then by Theorem 11.10

$$(\exists f) (\forall x \in a) [f'x : x \xrightarrow[1-1]{} b].$$

Also $x \in \cup(a) \rightarrow (\exists y) [x \in y \wedge y \in a]$. Again from Theorem 11.10

$$(\exists h) (\forall x \in \cup(a)) [x \in h'x \wedge h'x \in a].$$

We then define a mapping F on $\cup(a)$ thus

$$F'x = \langle h'x, (f'h'x)(x) \rangle, x \in \cup(a).$$

Since $x \in \cup(a) \rightarrow x \in h'x \wedge h'x \in a$ and $x \in h'x \wedge h'x \in a \rightarrow (f'h'x)(x) \in b$,

$$F : \cup(a) \rightarrow a \times b.$$

Furthermore if $F(x) = F(y)$ for $x \in \cup(a) \wedge y \in \cup(a)$ then

$$h'x = h'y \wedge (f'h'x)(x) = (f'h'y)(y).$$

But $h'x = h'y \rightarrow f'h'x = f'h'y$. Since $f'h'x$ is one-to-one

$$(f'h'x)(x) = (f'h'y)(y) \rightarrow x = y.$$

Thus $F : \cup(a) \xrightarrow{1-1} a \times b$ and hence

$$\overline{\cup(a)} \leq \overline{a \times b}.$$

***Theorem 11.12.** $\mathcal{U}_n(F) \wedge (\forall y \in b) [\overline{F'y} \leq \bar{a}] \rightarrow \overline{\cup(F''b)} \leq \overline{a \times b}$.

Proof. $\mathcal{U}_n(F) \rightarrow \overline{F''b} \leq \bar{b}$. Furthermore $x \in F''b \rightarrow (\exists y \in b) [x = F'y]$. Since $y \in b \rightarrow \overline{F'y} \leq \bar{a}$ we have from Theorem 11.11, $\overline{\cup(F''b)} \leq \overline{a \times F''b} \leq \overline{a \times b}$.

***Theorem 11.13.** $\aleph_{\alpha+1}$ is regular.

Proof. If $\aleph_\beta = \text{cf}(\aleph_{\alpha+1}) < \aleph_{\alpha+1}$ then $(\exists h) [[h : \aleph_\beta \rightarrow \aleph_{\alpha+1}] \wedge \aleph_{\alpha+1} = \cup(h''\aleph_\beta)]$. Since $(\forall \delta \in \aleph_\beta) [h'\delta < \aleph_{\alpha+1}]$ we have $\overline{h'\delta} \leq \aleph_\alpha$. Then from Theorem 11.12

$$\aleph_{\alpha+1} = \overline{\cup(h''\aleph_\beta)} \leq \overline{\aleph_\alpha \times \aleph_\beta} \leq \overline{\aleph_\alpha \times \aleph_\alpha} = \aleph_\alpha.$$

From this contradiction we conclude that $\text{cf}(\aleph_{\alpha+1}) = \aleph_{\alpha+1}$.

Remark. Every set a is equivalent to a subset of its power set $\mathcal{P}(a)$. From Cantor's Theorem we know that no set is equivalent to its power set. This we interpret to mean that the cardinality of $\mathcal{P}(a)$ is greater than the cardinality of a . Are there cardinalities intermediate between that of a and of $\mathcal{P}(a)$?

With the aid of the AC this question has a very simple formulation: Does there exist a set b such that

$$\bar{a} < \bar{b} < \overline{\mathcal{P}(a)}?$$

Using set equivalence the question can be formulated without the AC. Does there exist a set b such that a is equivalent to a subset of b , b is equivalent to a subset of $\mathcal{P}(a)$, yet b is neither equivalent to a nor to $\mathcal{P}(a)$.

If a is a finite set containing more than one element then such a set b will exist. If a is infinite it is known from the work of Cohen that the question is undecidable in ZF. The assertion that for every infinite set a there are no intermediate cardinalities between that of a and $\mathcal{P}(a)$ is called the Generalized Continuum Hypothesis (GCH), the Continuum Hypothesis (CH) being the special case $a = \omega$. In stating the GCH and the CH there is no loss in generality if we require that b be a subset of $\mathcal{P}(a)$.

Generalized Continuum Hypothesis:

$$(\forall a) (\forall b \subseteq \mathcal{P}(a)) [\text{Inf}(a) \wedge (\exists x) [a \subseteq x \simeq b] \rightarrow b \simeq a \vee b \simeq \mathcal{P}(a)].$$

Continuum Hypothesis:

$$(\forall b \subseteq \mathcal{P}(\omega)) (\exists x) [\omega \subseteq x \simeq b] \rightarrow b \simeq \omega \vee b \simeq \mathcal{P}(\omega).$$

A weaker form of the GCH is the following:

Aleph Hypothesis: $(\forall \alpha) [2^{\aleph_\alpha} = \aleph_{\alpha+1}]$.

It is easily established that there is a one-to-one correspondence between the points on a Euclidean line and the subsets of ω . Thus in the presence of the AC the continuum hypothesis is the conjecture that the number of points on a line, i.e. the cardinality of the continuum, is \aleph_1 .

In 1926 Lindenbaum and Tarski conjectured that the GCH implies the AC. This conjecture was proved by Sierpinski in 1947. For the proof of Sierpinski's Theorem we need the following lemmas.

Lemma 1. $a \cap b = 0 \rightarrow \mathcal{P}(a \cup b) \simeq \mathcal{P}(a) \times \mathcal{P}(b)$.

Proof. If

$$f'c = \langle a \cap c, b \cap c \rangle, \text{ for } c \subseteq a \cup b$$

then $f : \mathcal{P}(a \cup b) \rightarrow \mathcal{P}(a) \times \mathcal{P}(b)$. If $\langle x, y \rangle \in \mathcal{P}(a) \times \mathcal{P}(b)$ then $x \cup y \subseteq a \cup b$. Since $a \cap b = 0$ it follows that $a \cap y = 0$ and $b \cap x = 0$. Therefore

$$a \cap (x \cup y) = (a \cap x) \cup (a \cap y) = a \cap x = x$$

$$b \cap (x \cup y) = (b \cap x) \cup (b \cap y) = b \cap y = y$$

i.e. $f(x \cup y) = \langle x, y \rangle$.

If $f'c = f'd$ for $c \subseteq a \cup b \wedge d \subseteq a \cup b$ then

$$a \cap c = a \cap d \wedge b \cap c = b \cap d.$$

Therefore

$$c = (a \cap c) \cup (b \cap c) = (a \cap d) \cup (b \cap d) = d.$$

Thus $f : \mathcal{P}(a \cup b) \xrightarrow[\text{onto}]{1-1} \mathcal{P}(a) \times \mathcal{P}(b)$ i.e. $\mathcal{P}(a \cup b) \simeq \mathcal{P}(a) \times \mathcal{P}(b)$.

Lemma 2. If $a \simeq a \cup \{a\} \wedge b = \mathcal{P}(a)$ then $\mathcal{P}(b) \simeq \mathcal{P}(b) \times \mathcal{P}(b)$.

Proof. Since $a \simeq a \cup \{a\}$, $\mathcal{P}(a) \simeq \mathcal{P}(a \cup \{a\})$. If

$$f'c = \langle 0, c \rangle, \text{ for } c \subseteq a \cup \{a\} \wedge a \notin c$$

$$= \langle 1, c \rangle, \text{ for } c \subseteq a \cup \{a\} \wedge a \in c$$

then $f : \mathcal{P}(a \cup \{a\}) \rightarrow (\{0\} \times \mathcal{P}(a)) \cup (\{1\} \times \mathcal{P}(a \cup \{a\}))$. Furthermore if $c \subseteq a$ then

$$f(c) = \langle 0, c \rangle \wedge f(c \cup \{a\}) = \langle 1, c \rangle.$$

Therefore f is onto. Clearly $f'c = f'd$ implies $c = d$. Thus

$$\begin{aligned} b = \mathcal{P}(a) &\simeq \mathcal{P}(a \cup \{a\}) \simeq (\{0\} \times \mathcal{P}(a)) \cup (\{1\} \times \mathcal{P}(a)) \\ &= (\{0\} \times b) \cup (\{1\} \times b). \end{aligned}$$

Since $(\{0\} \times b) \cap (\{1\} \times b) = 0$ we have from Lemma 1

$$\mathcal{P}(b) \simeq \mathcal{P}(\{0\} \times b) \times \mathcal{P}(\{1\} \times b) \simeq \mathcal{P}(b) \times \mathcal{P}(b).$$

Lemma 3. If $a \simeq a \cup \{a\} \wedge b = \mathcal{P}(a) \wedge b \cup c \simeq \mathcal{P}(b)$ then $c \simeq \mathcal{P}(b)$.

Proof. If $b \cup c \simeq \mathcal{P}(b)$ then $(\exists f) [f : b \cup c \xrightarrow{\text{onto}} \mathcal{P}(b)]$. Then $f \upharpoonright c$ maps c one-to-one onto a subset of $\mathcal{P}(b)$.

From Lemma 2, $\mathcal{P}(b) \simeq \mathcal{P}(b) \times \mathcal{P}(b)$. Therefore

$$(\exists h) [h : b \cup c \xrightarrow{\text{onto}} \mathcal{P}(b) \times \mathcal{P}(b)].$$

If

$$\mathcal{P}(b) = \mathcal{D}(h''b)$$

and if $k'\langle x, y \rangle \triangleq x$ for $\langle x, y \rangle \in \mathcal{P}(b) \times \mathcal{P}(b)$ then $k \circ h : b \xrightarrow{\text{onto}} \mathcal{P}(b)$. Since this is not possible it follows that $(\exists x \in \mathcal{P}(b)) [x \notin \mathcal{D}(h''b)]$. Therefore $\{x\} \times \mathcal{P}(b) \subseteq h''c$, that is, h^{-1} maps $\{x\} \times \mathcal{P}(b)$ one-to-one onto a subset of c . But $\{x\} \times \mathcal{P}(b) \simeq \mathcal{P}(b)$.

Since c is equivalent to a subset of $\mathcal{P}(b)$ and $\mathcal{P}(b)$ is equivalent to a subset of c

$$c \simeq \mathcal{P}(b).$$

Lemma 4. If $a \simeq a \cup \{a\}$ then $\mathcal{P}(a) \simeq \mathcal{P}(a) \cup \{\mathcal{P}(a)\}$.

The proof is left to the reader.

Definition 5.

- 1) $\mathcal{P}^2(a) = \mathcal{P}(\mathcal{P}(a))$.
- 2) $\mathcal{P}^3(a) = \mathcal{P}(\mathcal{P}^2(a))$.
- 3) $\mathcal{P}^4(a) = \mathcal{P}(\mathcal{P}^3(a))$.

Definition 6. $\text{Fld}(a) = \mathcal{D}(a) \cup \mathcal{W}(a)$.

Theorem 11.14 (Sierpinski). $\text{GCH} \rightarrow \text{AC}$.

Proof. Since the AC is equivalent to the WOP it is sufficient to prove that every set c can be well ordered. If $a = c \cup \omega$ then c is equivalent to a subset of $\mathcal{P}(a)$. It is therefore sufficient to prove that if $b = \mathcal{P}(a)$ then b can be well ordered.

Since $\omega \subseteq a$, $a \simeq a \cup \{a\}$. By Lemma 4

$$\begin{aligned} b &\simeq b \cup \{b\}, \\ \mathcal{P}(b) &\simeq \mathcal{P}(b) \cup \{\mathcal{P}(b)\}, \\ \mathcal{P}^2(b) &\simeq \mathcal{P}^2(b) \cup \{\mathcal{P}^2(b)\}, \\ \mathcal{P}^3(b) &\simeq \mathcal{P}^3(b) \cup \{\mathcal{P}^3(b)\}. \end{aligned}$$

From Theorem 10.40

$$(\exists \beta) [\beta = \{\alpha \mid (\exists f) [f : \alpha \xrightarrow{1-1} b]\} \wedge \neg (\exists f) [f : \beta \xrightarrow{1-1} b]].$$

If $\forall \alpha < \beta$, $S_\alpha = \{r \subseteq b \times b \mid r \text{ We Fld}(r) \wedge (\exists f) [f \text{ Isom}_{E,r}(\alpha, \text{Fld}(r))]\}$ then since

$$\langle x, y \rangle \in b \times b \rightarrow \{\{x\}, \{x, y\}\} \in \mathcal{P}^2(b)$$

we have that $b \times b \subseteq \mathcal{P}^2(b)$ and hence $S_\alpha \subseteq \mathcal{P}^3(b)$. If

$$A = \{S_\alpha \mid \alpha < \beta\}$$

then $A \subseteq \mathcal{P}^4(b)$. Furthermore $\alpha < \beta \rightarrow (\exists f) [f : \alpha \xrightarrow{1-1} b]$. Then f induces a well ordering of $f''\alpha$ i.e. $(\exists r \subseteq b \times b) [\text{Fld}(r) = f''\alpha \wedge r \text{ We Fld}(r)]$. Therefore if $S_\alpha = S_\delta$ then

$$\begin{aligned} (\exists r \subseteq b \times b) [r \text{ We Fld}(r) \wedge (\exists f) [f \text{ Isom}_{E,r}(\alpha, \text{Fld}(r))] \\ \wedge (\exists g) [g \text{ Isom}_{E,r}(\delta, \text{Fld}(r))]] . \end{aligned}$$

Thus $\alpha = \delta$ and hence $\beta \simeq A$. Therefore A can be well ordered. Furthermore $\mathcal{P}^3(b)$ is equivalent to a subset of $\mathcal{P}^3(b) \cup A$ and since $A \subseteq \mathcal{P}^4(b)$ and since $\mathcal{P}^3(b)$ is equivalent to a subset of $\mathcal{P}^4(b)$ it follows that $\mathcal{P}^3(b) \cup A$ is equivalent to a subset of $\mathcal{P}^4(b)$. Then by the GCH

$$\mathcal{P}^3(b) \cup A \simeq \mathcal{P}^3(b) \vee \mathcal{P}^3(b) \cup A \simeq \mathcal{P}^4(b).$$

Since $\mathcal{P}^2(b) \simeq \mathcal{P}^2(b) \cup \{\mathcal{P}^2(b)\}$ it follows from Lemma 3 that if $\mathcal{P}^3(b) \cup A \simeq \mathcal{P}^4(b)$ then $A \simeq \mathcal{P}^4(b)$. From this it follows that $\mathcal{P}^4(b)$ can be well ordered and hence $\mathcal{P}^3(b)$ can be well ordered.

If $\mathcal{P}^3(b) \cup A \simeq \mathcal{P}^3(b)$ then A is equivalent to a subset of $\mathcal{P}^3(b)$. Then as before $\mathcal{P}^2(b)$ is equivalent to a subset of $\mathcal{P}^2(b) \cup A$ and $\mathcal{P}^2(b) \cup A$ is equivalent to a subset of $\mathcal{P}^3(b)$. Then by the GCH

$$\mathcal{P}^2(b) \cup A \simeq \mathcal{P}^2(A) \vee \mathcal{P}^2(b) \cup A \simeq \mathcal{P}^3(b).$$

Since $\mathcal{P}(b) \simeq \mathcal{P}(b) \cup \{\mathcal{P}(b)\}$ it follows from Lemma 3 that if $\mathcal{P}^2(b) \cup A \simeq \mathcal{P}^3(b)$ then $A \simeq \mathcal{P}^3(b)$. It then follows that $\mathcal{P}^3(b)$ can be well ordered and hence $\mathcal{P}^2(b)$ can be well ordered.

If $\mathcal{P}^2(b) \cup A \simeq \mathcal{P}^2(A)$ then A is equivalent to a subset of $\mathcal{P}^2(b)$. Then $\mathcal{P}(b)$ is equivalent to a subset of $\mathcal{P}(b) \cup A$ and $\mathcal{P}(b) \cup A$ is equivalent to a subset of $\mathcal{P}^2(b)$. From the GCH,

$$\mathcal{P}(b) \cup A \simeq \mathcal{P}(b) \vee \mathcal{P}(b) \cup A \simeq \mathcal{P}^2(b).$$

Since $b \simeq b \cup \{b\}$ we have from Lemma 3 that if $\mathcal{P}(b) \cup A \simeq \mathcal{P}^2(b)$ then $A \simeq \mathcal{P}^2(b)$. Thus $\mathcal{P}^2(b)$ can be well ordered and hence $\mathcal{P}(b)$ can be well ordered. If $\mathcal{P}(b) \cup A \simeq \mathcal{P}(b)$ then A is equivalent to a subset of $\mathcal{P}(b)$.

Finally, b is equivalent to a subset of $b \cup A$ and $b \cup A$ is equivalent to a subset of $\mathcal{P}(b)$. By the GCH,

$$b \cup A \simeq b \vee b \cup A \simeq \mathcal{P}(b).$$

Since $a \simeq a \cup \{a\}$ it follows from Lemma 3 that if $b \cup A \simeq \mathcal{P}(b)$ then $A \simeq \mathcal{P}(b)$. Therefore $\mathcal{P}(b)$ can be well ordered hence b can be well ordered.

If $b \cup A \simeq b$ it would follow that A is equivalent to a subset of b . But since $\beta \simeq A$ and β is not equivalent to a subset of b this is not possible.

We have thus established that b can be well ordered. But $b = \mathcal{P}(a)$; therefore a can be well ordered. Since $c \subseteq a$, c can be well ordered.

Remark. In 1960 H. Rubin proved that the Aleph Hypothesis (AH) also implies the AC. As we will see Rubin's proof does not require the full strength of the AH but only the hypothesis that the power set of each cardinal number can be well ordered.

Theorem 11.15 (Rubin). $AH \rightarrow AC$.

Proof. Among other things the AH assures us that the power set of each cardinal number can be well ordered. From this we will prove that for each α a function F_α can be defined on On by transfinite recursion in such a way that if $\beta < \alpha$ then $F_\alpha(\beta)$ is a well ordering of the set of sets of rank β .

It is sufficient to show that $F_\alpha \beta$ can be defined from $F_\alpha \upharpoonright \beta$ when $\beta < \alpha$. If $\beta = 0$ the set of sets of rank β is $\{0\}$. We define $F_\alpha \beta$ to be 0. If $0 < \beta < \alpha$ then $(\exists \gamma) [\beta = \gamma + 1]$. We then define a relation S_β on $R(\beta)$ by

$$x S_\beta y \iff \text{rank}(x) < \text{rank}(y) \vee [\text{rank}(x) = \text{rank}(y) \\ \wedge \langle x, y \rangle \in (F_\alpha \upharpoonright \beta)' \text{rank}' x].$$

Then S_β We $R(\beta)$. Therefore

$$(\exists! \delta) (\exists! f_\beta) [f_\beta \text{ Isom}_{S_\beta, E}(R(\beta), \delta)].$$

By Theorem 10.40

$$(\exists \tau) [\aleph_\tau = \{v \mid (\exists f) f : v \xrightarrow{1-1} R(\alpha)\}].$$

By hypothesis there exists a relation T that well orders $\mathcal{P}(\aleph_\tau)$. If

$$(\forall x \subseteq R(\beta)) [g_\beta' x = f_\beta'' x]$$

then $g_\beta : \mathcal{P}(R(\beta)) \xrightarrow{1-1} \mathcal{P}(\aleph_\tau)$.

If $\text{rank}(x) = \text{rank}(y) = \beta$, $x \subseteq R(\beta) \wedge y \subseteq R(\beta)$. We then define

$$F_\alpha \beta = \{\langle x, y \rangle \mid \text{rank}(x) = \text{rank}(y) = \beta \wedge g_\beta(x) T g_\beta(y)\}.$$

It follows that $F_\alpha \beta$ well orders the set of sets of rank β since T well orders $\mathcal{P}(\aleph_\tau)$.

Remark. Clearly the GCH implies the AH. From Theorem 11.15 we see that the AH implies the AC. In 1963 Cohen proved that the AC does not imply the AH and hence does not imply GCH.

Since the AC implies that all sets are comparable we have from Theorems 11.14 and 11.15

Theorem 11.16. $\text{GCH} \leftrightarrow \text{AH}$.

Remark. It should be noted that we have not proved that GCH and AH are logically equivalent. That $\text{GCH} \rightarrow \text{AH}$ is a logical implication is clear. Theorem 11.15 however was proved using the Axiom of Regularity.

In proving that the GCH is consistent with ZF we will provide a model of ZF in which the GCH holds. We will prove this by showing simply that the AH holds in this model.

Theorem 11.17. $\text{GCH} \rightarrow [\text{Inacc}_w(\aleph_\alpha) \leftrightarrow \text{Inacc}(\aleph_\alpha)]$.

Proof. By definition \aleph_α inaccessible implies \aleph_α weakly inaccessible. Conversely

$$\bar{x} < \aleph_\alpha \rightarrow (\exists \beta) [\bar{x} = \aleph_\beta \wedge \beta < \alpha].$$

Since $\alpha \in K_{\text{II}}$, $\beta + 1 < \alpha$ hence $\aleph_{\beta+1} < \aleph_\alpha$. But by the GCH

$$\overline{\overline{\mathcal{P}(\bar{x})}} = \aleph_{\beta+1}.$$

***Theorem 11.18.** $\text{Inf}(b) \wedge 2 \leq \bar{a} \leq \overline{\overline{\mathcal{P}(b)}} \rightarrow \bar{a}^b = \overline{\overline{\mathcal{P}(b)}}$.

Proof. By Theorem 10.49, $\overline{\overline{\mathcal{P}(b)}} = \overline{2^b}$. Therefore $\bar{a} \leq \overline{\overline{\mathcal{P}(b)}} \rightarrow \bar{a} \leq \overline{2^b}$. Then

$$\bar{a}^b \leq \overline{\overline{(2^b)^b}} = \overline{\overline{2^{b \times b}}} = \overline{2^b} = \overline{\overline{\mathcal{P}(b)}}$$

i.e. $\bar{a}^b \leq \overline{\overline{\mathcal{P}(b)}}$.

On the other hand $2 \leq \bar{a} \rightarrow \overline{2^b} \leq \bar{a}^b$. Therefore

$$\bar{a}^b = \overline{\overline{\mathcal{P}(b)}}.$$

***Corollary 11.19.**

- 1) $\aleph_\alpha \leq \aleph_\beta \rightarrow \overline{\overline{\aleph_\alpha^{\aleph_\beta}}} = \overline{\overline{2^{\aleph_\beta}}}$.
- 2) $\overline{\overline{\aleph_\alpha^{\aleph_\alpha}}} = \overline{\overline{2^{\aleph_\alpha}}}$.
- 3) $\aleph_\alpha \leq \aleph_\beta \rightarrow \overline{\overline{\aleph_\beta^{\aleph_\alpha}}} \leq \overline{\overline{2^{\aleph_\beta}}}$.

Proof.

1) Since \aleph_β is infinite and $2 \leq \aleph_\alpha \leq \aleph_\beta < \overline{\overline{\mathcal{P}(\aleph_\beta)}}$ we have from Theorem 11.18

$$\overline{\overline{\aleph_\alpha^{\aleph_\beta}}} = \overline{\overline{\mathcal{P}(\aleph_\beta)}} = \overline{\overline{2^{\aleph_\beta}}}.$$

2) Obvious from 1) with $\alpha = \beta$.

3) $\overline{\overline{\aleph_\beta^{\aleph_\alpha}}} \leq \overline{\overline{\aleph_\beta^{\aleph_\beta}}} = \overline{\overline{2^{\aleph_\beta}}}$.

***Theorem 11.20.**

$\alpha \in K_{\text{II}} \wedge (\forall \gamma < \alpha) [\overline{2^{\aleph_\gamma}} < \aleph_\alpha] \wedge [\aleph_\beta < \text{cf}(\aleph_\alpha)] \rightarrow \overline{\overline{\aleph_\alpha^{\aleph_\beta}}} = \aleph_\alpha$.

Proof. If $a = \{\aleph_\gamma^{\aleph_\beta} \mid \gamma < \alpha\}$ and if $f \in \cup(a)$ then

$$(\exists \gamma < \alpha) [f : \aleph_\beta \rightarrow \aleph_\gamma].$$

Since $\aleph_\gamma \subset \aleph_\alpha$ it follows that $f \in \aleph_\alpha^{\aleph_\beta}$ i.e.

$$\cup(a) \subseteq \aleph_\alpha^{\aleph_\beta}.$$

If $f \in \aleph_\alpha^{\aleph_\beta}$ then $f: \aleph_\beta \rightarrow \aleph_\alpha$. Since $\aleph_\beta < \text{cf}(\aleph_\alpha)$ it follows that

$$(\exists \delta < \aleph_\alpha) [\mathcal{W}(f) \subseteq \delta].$$

Furthermore since $\alpha \in K_{\text{II}}$

$$(\exists \gamma < \alpha) [\delta \leq \aleph_\gamma < \aleph_\alpha].$$

Therefore $f \in \aleph_\gamma^{\aleph_\beta}$ i.e. $f \in \cup(a)$.

Thus $\aleph_\alpha^{\aleph_\beta} = \cup(a)$ and hence

$$\overline{\aleph_\alpha^{\aleph_\beta}} = \overline{\cup(a)}.$$

Further $x \in a \rightarrow (\exists \gamma < \alpha) [x = \aleph_\gamma^{\aleph_\beta}]$. If $\gamma < \beta$

$$\overline{\aleph_\gamma^{\aleph_\beta}} \leq \overline{\aleph_\beta^{\aleph_\beta}} = \overline{2^{\aleph_\beta}} < \overline{\aleph_\alpha}.$$

If $\beta \leq \gamma$ then by *Corollary 11.19

$$\overline{\aleph_\gamma^{\aleph_\beta}} \leq \overline{2^{\aleph_\gamma}} < \overline{\aleph_\alpha}.$$

Finally since $\bar{a} \leq \bar{\alpha} \leq \aleph_\alpha$ we have from *Theorem 11.11

$$\overline{\cup(a)} \leq \overline{\aleph_\alpha \times \aleph_\alpha} = \overline{\aleph_\alpha}.$$

Therefore

$$\overline{\aleph_\alpha^{\aleph_\beta}} = \overline{\aleph_\alpha}.$$

Definition 11.21. $\prod_{x \in a} c'x \triangleq \{g \mid g \mathcal{F}n a \wedge (\forall x \in a) [g'x \in c'x]\}$.

Remark. $\prod_{x \in a} c'x$ is called the cross product of the collection of sets $\{c'x \mid x \in a\}$. To see that this is a reasonable generalization of the cross product of two sets note that if $a = 2$ and if $g \in \prod_{x \in 2} c'x$ then $g \mathcal{F}n 2 \wedge g'0 \in c'0 \wedge g'1 \in c'1$ i.e., $\langle g'0, g'1 \rangle \in c'0 \times c'1$. Conversely if $\langle x, y \rangle \in c'0 \times c'1$ and if we define g on 2 by $g'0 = x \wedge g'1 = y$ then $g \in \prod_{x \in a} c'x$. Clearly there is a natural one-to-one correspondence between $\prod_{x \in a} c'x$ and $c'0 \times c'1$.

Theorem 11.22. $\mathcal{M}\left(\prod_{x \in a} c'x\right)$.

Proof. $\prod_{x \in a} c'x \subseteq [\cup(c''a)]^a$.

***Theorem 11.23 (Zermelo).** $(\forall x \in a) [\overline{b'x} < \overline{c'x}] \rightarrow \overline{\cup(b''a)} < \overline{\prod_{x \in a} c'x}$.

Proof. (By contradiction.) Otherwise

$$(\exists f) \left[f: \cup(b''a) \xrightarrow{\text{onto}} \prod_{x \in a} c'x \right].$$

If we define d on a by

$$d'x = \{(f'z)'x \mid z \in b'x\}.$$

Then $\overline{d'x} \leq \overline{b'x}$ and $d'x \subseteq c'x$. Since by hypothesis $\overline{b'x} < \overline{c'x}$ it follows that $\overline{d'x} < \overline{c'x}$ and hence $c'x - d'x \neq 0$. Therefore by the AC

$$(\exists e) (\forall x \in a) [e'x \in c'x - d'x].$$

Since $e'x \in c'x$ and since $e \mathcal{F}_n a$, $e \in \prod_{x \in a} c'x$. Then

$$(\exists z \in \cup(b''a)) [f'z = e].$$

But $z \in \cup(b''a) \rightarrow (\exists x \in a) [z \in b'x]$. Consequently

$$(f'z)'x \in d'x \wedge (f'z)'x \in c'x - d'x.$$

Definition 11.24. $a^+ \triangleq \mu_\alpha(\bar{\alpha} > \bar{a})$.

Remark. Note that a^+ is a cardinal number.

***Theorem 11.25.** $\aleph_\alpha < \overline{\aleph_\alpha^{\text{cf}(\aleph_\alpha)}}$.

Proof. If $\alpha \in K_1$ then $\text{cf}(\aleph_\alpha) = \aleph_\alpha$. Therefore

$$\aleph_\alpha < 2^{\aleph_\alpha} = \overline{\aleph_\alpha^{\aleph_\alpha}} = \overline{\aleph_\alpha^{\text{cf}(\aleph_\alpha)}}.$$

If $\alpha \in K_{II}$ and if $\beta = \text{cf}(\aleph_\alpha)$ then

$$(\exists f) [\text{Smo}(f) \wedge [f: \beta \rightarrow \aleph_\alpha] \wedge \aleph_\alpha = \cup(f''\beta)].$$

If

$$c'\gamma = (f'\gamma)^+, \quad \gamma < \beta$$

then $(\forall \gamma < \beta) [\overline{f'\gamma} < \overline{c'\gamma}]$. Therefore by *Theorem 11.23

$$\aleph_\alpha = \overline{\cup(f''\beta)} < \prod_{\gamma < \beta} c'\gamma.$$

Since

$$\prod_{\gamma < \beta} c'\gamma \subseteq (\cup(c''\beta))^\beta,$$

$$\overline{\prod_{\gamma < \beta} c'\gamma} \leq \overline{(\cup(c''\beta))^\beta}.$$

Furthermore $\gamma < \beta \rightarrow c'\gamma < \aleph_\alpha$ and $\beta \leq \aleph_\alpha$. Therefore

$$\overline{(\cup(c''\beta))^\beta} \leq \overline{\aleph_\alpha^\beta} = \overline{\aleph_\alpha \times \aleph_\alpha} = \aleph_\alpha$$

and hence

$$\overline{(\cup(c''\beta))^\beta} \leq \overline{\aleph_\alpha^\beta} = \overline{\aleph_\alpha^{\text{cf}(\aleph_\alpha)}}$$

i.e.

$$\aleph_\alpha < \overline{\aleph_\alpha^{\text{cf}(\aleph_\alpha)}}.$$

***Theorem 11.26.** $\aleph_\beta < \text{cf}(\aleph_\alpha^{\aleph_\beta})$.

Proof (by contradiction). If $\text{cf}(\aleph_\alpha^{\aleph_\beta}) = \aleph_\gamma \leq \aleph_\beta$ then by Theorems 11.25, 10.50, and 10.35

$$\aleph_\alpha^{\aleph_\beta} < (\aleph_\alpha^{\aleph_\beta})^{\aleph_\gamma} = \aleph_\alpha^{\aleph_\beta \times \aleph_\gamma} = \aleph_\alpha^{\aleph_\beta}.$$

***Corollary 11.27** (König). $\aleph_\alpha < \text{cf}(2^{\aleph_\alpha})$.

Proof. $\aleph_\alpha < \text{cf}(\aleph_0^{\aleph_\alpha}) = \text{cf}(2^{\aleph_\alpha})$.

Theorem 11.28. $\text{GCH} \rightarrow \aleph_\alpha^{\aleph_\beta} = \aleph_\alpha$ if $\aleph_\beta < \text{cf}(\aleph_\alpha)$
 $= \aleph_{\alpha+1}$ if $\text{cf}(\aleph_\alpha) \leq \aleph_\beta \leq \aleph_\alpha$
 $= \aleph_{\beta+1}$ if $\aleph_\alpha \leq \aleph_\beta$.

Proof. If $\aleph_\beta < \text{cf}(\aleph_\alpha)$ then $\alpha \neq 0$. If $(\exists \gamma) [\alpha = \gamma + 1]$ then \aleph_α is regular and hence

$$\aleph_\beta < \text{cf}(\aleph_\alpha) = \aleph_\alpha = \aleph_{\gamma+1}.$$

Therefore $\aleph_\beta \leq \aleph_\gamma$. Since by the GCH $\aleph_{\gamma+1} = 2^{\aleph_\gamma}$

$$\aleph_\alpha^{\aleph_\beta} = \aleph_{\gamma+1}^{\aleph_\beta} = (2^{\aleph_\gamma})^{\aleph_\beta} = 2^{\aleph_\gamma \times \aleph_\beta} = 2^{\aleph_\gamma} = \aleph_{\gamma+1} = \aleph_\alpha.$$

If $\alpha \in K_{\text{II}}$ then since $\gamma < \alpha \rightarrow 2^{\aleph_\gamma} = \aleph_{\gamma+1} < \aleph_\alpha$ we have from *Theorem 11.20

$$\aleph_\alpha^{\aleph_\beta} = \aleph_\alpha.$$

If $\text{cf}(\aleph_\alpha) \leq \aleph_\beta \leq \aleph_\alpha$ then from *Theorem 11.26 and *Corollary 11.19

$$\aleph_\alpha < \aleph_\alpha^{\text{cf}(\aleph_\alpha)} \leq \aleph_\alpha^{\aleph_\beta} \leq 2^{\aleph_\alpha} = \aleph_{\alpha+1}.$$

That is $\aleph_\alpha < \aleph_\alpha^{\aleph_\beta} \leq \aleph_{\alpha+1}$. Therefore

$$\aleph_\alpha^{\aleph_\beta} = \aleph_{\alpha+1}.$$

If $\aleph_\alpha \leq \aleph_\beta$ then since $\aleph_\beta < \aleph_{\beta+1} = 2^{\aleph_\beta}$

$$\aleph_\alpha^{\aleph_\beta} \leq (2^{\aleph_\beta})^{\aleph_\beta} = 2^{\aleph_\beta \times \aleph_\beta} = 2^{\aleph_\beta} = \aleph_{\beta+1}.$$

By Theorem 11.26

$$\aleph_\beta < \text{cf}(\aleph_\alpha^{\aleph_\beta}) \leq \aleph_\alpha^{\aleph_\beta}.$$

Therefore

$$\aleph_\alpha^{\aleph_\beta} = \aleph_{\beta+1}.$$

Remark. With the aid of the AC we can also improve on Theorem 9.16.

***Theorem 11.29.** $\text{Inf}(a) \wedge \text{Cl}(R_1, A) \wedge \cdots \wedge \text{Cl}(R_m, A) \wedge \text{Cl}_2(S_1, A) \wedge \cdots$
 $\cdots \wedge \text{Cl}_2(S_n, A) \wedge \mathcal{U}n(R_1) \wedge \cdots \wedge \mathcal{U}n(R_m) \wedge \mathcal{U}n(S_1) \wedge \cdots \wedge \mathcal{U}n(S_n) \wedge a \subset A$
 $\rightarrow (\exists b \subset A) [\text{Cl}(R_1, b) \wedge \cdots \wedge \text{Cl}(R_m, b) \wedge \text{Cl}_2(S_1, b) \wedge \cdots \wedge \text{Cl}_2(S_n, b)$
 $\wedge a \subseteq b \wedge \bar{a} = \bar{b}]$.

Proof. The proof proceeds as for Theorem 9.16. We then note that since R_i and S_i are single valued

$$\begin{aligned} \overline{\overline{R_i f'k}} &\leq \overline{f'k} \quad i = 1, \dots, m \\ \overline{\overline{S_i (f'k)^2}} &\leq \overline{(f'k)^2} = \overline{f'k}, \quad i = 1, \dots, n. \end{aligned}$$

Therefore $f(k+1)$ is the union of a finite number of sets each of cardinality not greater than $\overline{f'k}$. Then by Theorem 10.35

$$\overline{\overline{f(k+1)}} = \overline{\overline{f(k)}} = \dots = \overline{\overline{f(0)}} = \overline{a}.$$

Then from Theorem 11.11 and the fact that a is infinite

$$\overline{b} \leq \overline{a \times \omega} = \overline{a}.$$

Furthermore since $a \subseteq b$, $\overline{a} \leq \overline{b}$. Therefore $\overline{a} = \overline{b}$.

12 Models

We turn now to the very interesting subject of models of set theory. Intuitively by a model of set theory we mean a system in which the axioms and theorems of ZF are true. Such a system must consist of a domain of objects that we interpret as the universe V of our theory and a binary relation that we interpret as the \in -relation of our theory.

Assuming consistency there is a model of ZF consisting of a universe of “sets” V on which there is defined an “ \in -relation”. Given such a universe V it is possible that some subclass A of V together with some relation R on A is also a model of ZF. With $A \subseteq V$ and $R \subseteq A \times A$ the language of ZF is adequate for the development of a theory of such internal models. Our next objective is to make the foregoing ideas precise and thereby compel ZF to tell us about some of its models.

In order to define model we first introduce the idea of a structure or relational system. For each class A and each relation $R \subseteq A \times A$ we introduce the term

$$[A, R]$$

which we call a structure (or relational system); A is the universe of this structure and the elements of A we call individuals.

We next define “the structure $[A, R]$ satisfies the wff φ .” We introduce the predicate symbol “ \models ” which is read “satisfies”. Our definition is by induction on the number of logical symbols in φ . For this purpose we assume \neg , \wedge , and \forall as primitive.

Definition 12.1.

- 1) $[A, R] \models a \in b \iff a \in A \wedge b \in A \wedge aRb.$
- 2) $[A, R] \models \neg \psi \xleftrightarrow{\Delta} \neg [[A, R] \models \psi].$
- 3) $[A, R] \models \psi \wedge \eta \xleftrightarrow{\Delta} [[A, R] \models \psi] \wedge [[A, R] \models \eta].$
- 4) $[A, R] \models (\forall x) \psi(x) \xleftrightarrow{\Delta} (\forall x \in A) [[A, R] \models \psi(x)].$

Remark. With the understanding that the terms $[A, R]$ and the satisfaction symbol \models occur only in contexts covered by Definition 12.1 it is clear that these symbols can be eliminated from our language,

that is,

$$[A, R] \models \varphi$$

is an abbreviation for a wff of our language.

If $A \neq 0 \wedge [A, R] \models \varphi$ we say that $[A, R]$ is a model of φ . $[A, R]$ is a model of a collection of wffs provided $[A, R] \models \varphi$ for each φ in the collection. In order to prove that a certain structure $[A, R]$ is a model of a given wff we must prove a certain wff in ZF namely the well formed formula $[A, R] \models \varphi$. We will be particularly interested in structures for which R is the usual \in -relation. Such a structure we call a standard structure.

Definition 12.2. $[A, R]$ is a standard structure iff $R = E \cap A^2$.

Definition 12.3. $A \models \varphi \stackrel{\Delta}{\longleftrightarrow} [A, E \cap A^2] \models \varphi$.

Definition 12.4.

- 1) $[a \in b]^A \stackrel{\Delta}{\longleftrightarrow} a \in b$.
- 2) $[\neg \psi]^A \stackrel{\Delta}{\longleftrightarrow} \neg \psi^A$.
- 3) $[\psi \wedge \eta]^A \stackrel{\Delta}{\longleftrightarrow} \psi^A \wedge \eta^A$.
- 4) $[(\forall x) \psi(x)]^A \longleftrightarrow (\forall x \in A) [\psi^A(x)]$.

Remark. From Definition 12.4 we see that φ^A is simply the wff obtained from φ by replacing each occurrence of a quantified variable $(\forall x)$ by $(\forall x \in A)$.

Theorem 12.5.

- 1) If φ is closed then

$$A \models \varphi \longleftrightarrow \varphi^A.$$

- 2) If all free variables occurring in φ are among a_1, \dots, a_n then

$$a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [A \models \varphi \longleftrightarrow \varphi^A].$$

Proof. We consider 1) to be the special case of 2) with $n = 0$. We prove 2) by induction on the number of logical symbols in φ .

If φ is of the form $a \in b$ and if a and b are among a_1, \dots, a_n then

$$\begin{aligned} a_1 \in A \wedge \dots \wedge a_n \in A &\rightarrow [a \in b \wedge a \in A \wedge b \in A \longleftrightarrow a \in b] \\ &\rightarrow [A \models \varphi \longleftrightarrow \varphi^A]. \end{aligned}$$

If φ is of the form $\neg \psi$ and if all of the free variables of φ are among a_1, \dots, a_n , then all of the free variables of ψ are among a_1, \dots, a_n . Then

from the induction hypothesis

$$\begin{aligned} a_1 \in A \wedge \cdots \wedge a_n \in A &\rightarrow [A \models \psi \leftrightarrow \psi^A] \\ &\rightarrow [\neg [A \models \psi] \leftrightarrow \neg \psi^A] \\ &\rightarrow [A \models \varphi \leftrightarrow \varphi^A]. \end{aligned}$$

If φ is of the form $\psi \wedge \eta$ and if all of the free variables of φ are among a_1, \dots, a_n then all of the free variables of ψ and of η are among a_1, \dots, a_n . Then

$$\begin{aligned} a_1 \in A \wedge \cdots \wedge a_n \in A &\rightarrow [A \models \psi \leftrightarrow \psi^A], \\ a_1 \in A \wedge \cdots \wedge a_n \in A &\rightarrow [A \models \eta \leftrightarrow \eta^A]. \end{aligned}$$

Therefore

$$\begin{aligned} a_1 \in A \wedge \cdots \wedge a_n \in A &\rightarrow [A \models \psi \wedge A \models \eta \leftrightarrow \psi^A \wedge \eta^A] \\ &\rightarrow [A \models \varphi \leftrightarrow \varphi^A]. \end{aligned}$$

If φ is of the form $(\forall x) \psi(x)$ and if all of the free variables of φ are among a_1, \dots, a_n then there is an x that is not among a_1, \dots, a_n and all of the free variables of $\psi(x)$ are among a_1, \dots, a_n, x . From the induction hypotheses

$$\begin{aligned} a_1 \in A \wedge \cdots \wedge a_n \in A \wedge x \in A &\rightarrow [A \models \psi(x) \leftrightarrow \psi^A(x)]. \\ a_1 \in A \wedge \cdots \wedge a_n \in A &\rightarrow [[x \in A \rightarrow A \models \psi(x)] \leftrightarrow [x \in A \rightarrow \psi^A(x)]]. \end{aligned}$$

Since x is not among a_1, \dots, a_n we generalize on x and distribute to obtain

$$\begin{aligned} a_1 \in A \wedge \cdots \wedge a_n \in A &\rightarrow [(\forall x \in A) A \models \psi(x) \leftrightarrow (\forall x \in A) \psi^A(x)] \\ &\rightarrow [A \models \varphi \leftrightarrow \varphi^A]. \end{aligned}$$

Remark. Theorem 12.5 is a basic result. It assures us that if φ is closed A is a model of φ if and only if φ^A is a theorem in ZF.

Suppose that $A \models \varphi$ and φ is equivalent to ψ i.e. $\varphi \leftrightarrow \psi$. Does it follow that $A \models \psi$? To answer this question we need a result from logic for which we review the axioms for our logic and the rules of inference.

Axioms.

- 1) $\varphi \rightarrow [\psi \rightarrow \varphi]$.
- 2) $[\varphi \rightarrow [\psi \rightarrow \eta]] \rightarrow [[\varphi \rightarrow \psi] \rightarrow [\varphi \rightarrow \eta]]$.
- 3) $[\neg \varphi \rightarrow \neg \psi] \rightarrow [\psi \rightarrow \varphi]$.
- 4) $(\forall x) [\varphi \rightarrow \psi] \rightarrow [\varphi \rightarrow (\forall x) \psi]$ where x is not free in φ .
- 5) $(\forall x) \varphi(x) \rightarrow \varphi(a)$ where x has no free occurrence in a wf part of φ of the form $(\forall a) \psi$.

Rules of Inference.

- 1) From $\varphi \rightarrow \psi$ and φ to infer ψ .
- 2) From φ to infer $(\forall x) \varphi$.

Theorem 12.6. *If $\vdash \varphi$ and if A is a nonempty class that satisfies each nonlogical axiom that occurs in some proof of φ then*

- 1) φ^A if φ is closed.
- 2) $a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow \varphi^A$ if all of the free variable of φ are among a_1, \dots, a_n .

Proof. We regard 1) as the special case of 2) with $n = 0$. Since φ is a theorem it has a proof and indeed by hypothesis a proof in which each nonlogical axiom is satisfied by A . Suppose that the sequence of wff

$$\eta_1, \dots, \eta_m$$

is such a proof. Then η_m is φ and each η_k is either an axiom or is inferred from previous formulas in the sequence by one of the rules of inference. Our procedure is to show that the sequence

$$\eta_1, \dots, \eta_m$$

can be modified to produce a proof of 2). More precisely we will prove by induction that for each $\eta_k, k = 1, \dots, m$, if all of the free variables in η_k are among b_1, \dots, b_p then

$$b_1 \in A \wedge \dots \wedge b_p \in A \rightarrow \eta_k^A.$$

Case 1. η_k is an axiom. If η_k is of the form $\psi \rightarrow [\eta \rightarrow \psi]$ then η_k^A is

$$\psi^A \rightarrow [\eta^A \rightarrow \psi^A]$$

i.e. η_k^A is an axiom. Hence

$$b_1 \in A \wedge \dots \wedge b_p \in A \rightarrow \eta_k^A.$$

If η_k is of the form $[\psi \rightarrow [\theta \rightarrow \zeta]] \rightarrow [[\psi \rightarrow \theta] \rightarrow [\psi \rightarrow \zeta]]$ or $[\neg \psi \rightarrow \neg \theta] \rightarrow [\theta \rightarrow \psi]$ an argument similar to the foregoing leads to

$$b_1 \in A \wedge \dots \wedge b_p \in A \rightarrow \eta_k^A.$$

If η_k is of the form $(\forall x) [\psi \rightarrow \theta] \rightarrow [\psi \rightarrow (\forall x)\theta]$ where x is not free in ψ then from the tautology $[p \rightarrow [q \rightarrow r]] \rightarrow [q \rightarrow [p \rightarrow r]]$ we have

$$[x \in A \rightarrow [\psi^A \rightarrow \theta^A]] \rightarrow [\psi^A \rightarrow [x \in A \rightarrow \theta^A]],$$

$$(\forall x) [x \in A \rightarrow [\psi^A \rightarrow \theta^A]] \rightarrow [\psi^A \rightarrow (\forall x) [x \in A \rightarrow \theta^A]],$$

$$b_1 \in A \wedge \dots \wedge b_p \in A \rightarrow \eta_k^A.$$

If η_k is of the form $(\forall x) \psi(x) \rightarrow \psi(a)$ then as an instance of this same axiom we have

$$\begin{aligned} (\forall x) [x \in A \rightarrow \psi(x)] &\rightarrow [a \in A \rightarrow \psi(a)], \\ a \in A &\rightarrow [(\forall x \in A) \psi(x) \rightarrow \psi(a)] \end{aligned}$$

and hence

$$b_1 \in A \wedge \cdots \wedge b_p \in A \rightarrow \eta_k^A.$$

If η_k is an axiom of ZF then by hypothesis $A \models \eta_k$, and from Theorem 12.5

$$b_1 \in A \wedge \cdots \wedge b_p \in A \rightarrow \eta_k^A.$$

Case 2. If η_k is inferred by modus ponens from η_i and $\eta_i \rightarrow \eta_k$ and if all of the free variables of η_i are among $b_1, \dots, b_p, c_1, \dots, c_q$ with c_1, \dots, c_q all distinct and none of them occur among b_1, \dots, b_p then from our induction hypothesis

$$\begin{aligned} b_1 \in A \wedge \cdots \wedge b_p \in A \wedge c_1 \in A \wedge \cdots \wedge c_q \in A &\rightarrow \eta_i^A, \\ b_1 \in A \wedge \cdots \wedge b_p \in A \wedge c_1 \in A \wedge \cdots \wedge c_q \in A &\rightarrow [\eta_i^A \rightarrow \eta_k^A]. \end{aligned}$$

From the self distributive law of implication and modus ponens

$$b_1 \in A \wedge \cdots \wedge b_p \in A \wedge c_1 \in A \wedge \cdots \wedge c_{q-1} \in A \rightarrow [c_q \in A \rightarrow \eta_k^A].$$

Since c_q does not occur among $b_1, \dots, b_p, c_1, \dots, c_{q-1}$ we have by generalization

$$\begin{aligned} b_1 \in A \wedge \cdots \wedge b_p \in A \wedge c_1 \in A \wedge \cdots \wedge c_{q-1} \in A &\rightarrow (\forall c_q) [c_q \in A \rightarrow \eta_k^A] \\ &\rightarrow [(\exists c_q) [c_q \in A] \rightarrow \eta_k^A] \\ (\exists c_q) [c_q \in A] &\rightarrow [b_1 \in A \wedge \cdots \wedge b_p \in A \wedge c_1 \in A \wedge \cdots \wedge c_{q-1} \in A \rightarrow \eta_k^A]. \end{aligned}$$

Since $A \neq 0$

$$b_1 \in A \wedge \cdots \wedge b_p \in A \wedge c_1 \in A \wedge \cdots \wedge c_{q-1} \in A \rightarrow \eta_k^A.$$

With $q - 1$ repetitions we obtain

$$b_1 \in A \wedge \cdots \wedge b_p \in A \rightarrow \eta_k^A.$$

Case 3. If η_k is inferred from $\eta_i(x)$ by generalization then there is an x not among b_1, \dots, b_p . From the induction hypothesis

$$b_1 \in A \wedge \cdots \wedge b_p \in A \wedge x \in A \rightarrow \eta_i^A(x).$$

Since x is not among b_1, \dots, b_p we have by generalization

$$b_1 \in A \wedge \cdots \wedge b_p \in A \rightarrow (\forall x \in A) \eta_i^A(x).$$

Remark. From Theorem 12.6 we see that if a proof of a wff φ requires only the logical axioms then every nonempty class A will be a model of φ . In particular every nonempty class A is a model of the logical axioms and if two wffs are logically equivalent i.e.

$$\vdash_{\text{LA}} \varphi \leftrightarrow \psi$$

then a nonempty class A is a model of φ iff it is a model of ψ .

We are interested in classes A that are models of ZF. Since there are infinitely many axioms for ZF the assertion that A is a model of ZF is the assertion that each wff in a certain infinite collection of wffs is a theorem in ZF. This assertion we abbreviate as the metastatement, $A \models \text{ZF}$.

From Theorem 12.6 we see that if $A \neq 0$ and $A \models \text{ZF}$ then every theorem of ZF holds in A , that is, A satisfies each theorem of ZF. In the next section we will give conditions on A that assure that $A \models \text{ZF}$. One requirement for most results of that section is that A be transitive.

Definition 12.7. $\text{STM}(A, \varphi) \stackrel{\Delta}{\longleftrightarrow} A \neq 0 \wedge \text{Tr}(A) \wedge A \models \varphi$.

Remark. By a standard transitive model of ZF we mean a nonempty transitive class A that satisfies each axiom of ZF, i.e. for each axiom φ

$$\text{STM}(A, \varphi).$$

Although we restrict our discussion to standard transitive models of ZF this theory nevertheless encompasses a large class of models of ZF as we see from the following theorem.

Theorem 12.8 (Mostowski). *If $R \subseteq A^2 \wedge R \text{ Wfr } A \wedge (\forall x \in A) (\forall y \in A) (\forall z) [zRx \leftrightarrow zRy] \rightarrow x = y$ then there exists a B and F such that*

- 1) $\text{Tr}(B)$.
- 2) $F \text{ Isom}_{R,E}(A, B)$.
- 3) $[A, R] \models \varphi \leftrightarrow B \models \varphi$ if φ is closed.
- 4) *If all of the free variables of φ are among a_1, \dots, a_n and if*

$$a_1 \in A \wedge \dots \wedge a_n \in A$$

then

$$[[A, R] \models \varphi(a_1, \dots, a_n) \leftrightarrow B \models \varphi(F'a_1, \dots, F'a_n)].$$

Proof. 1) If $R \text{ Wfr } A \wedge x \in A$ then $R^{-1}\{x\}$ is a set. Therefore if $f \mathcal{F}_n R^{-1}\{x\}$ then $\mathcal{W}(f)$ is a set. If

$$K = \{f \mid (\exists a \subseteq A) [f \mathcal{F}_n a \wedge (\forall x \in a) [f'x = \{f'y \mid yRx\} \wedge R^{-1}\{x\} \subseteq a]]\}$$

then any two functions in K have the same values at any point common to their domains: Otherwise there would exist an f and g in K and an x in $\mathcal{D}(f) \cap \mathcal{D}(g)$ such that $f(x) \neq g(x)$. If $c = \{x \in \mathcal{D}(f) \cap \mathcal{D}(g) \mid f'x \neq g'x\}$

then $c \neq 0 \wedge c \subseteq A$. Therefore $(\exists x \in c) [c \cap R^{-1}\{x\} = 0]$. Since $x \in c$

$$R^{-1}\{x\} \subseteq \mathcal{D}(f) \wedge R^{-1}\{x\} \subseteq \mathcal{D}(g).$$

Then $yRx \rightarrow f'y = g'y$ and hence

$$f'x = \{f'y | yRx\} = \{g'y | yRx\} = g'x.$$

Furthermore each f in K is one-to-one for otherwise

$$(\exists f \in K) (\exists x \in \mathcal{D}(f)) (\exists y \in \mathcal{D}(f)) [x \neq y \wedge f'x = f'y].$$

If $c = \{x \in \mathcal{D}(f) | (\exists y \in \mathcal{D}(f)) [x \neq y \wedge f'x = f'y]\}$ then $c \neq 0 \wedge c \subseteq A$. Therefore c has an R-minimal element i.e. $(\exists x \in c) [c \cap R^{-1}\{x\} = 0]$. Since $x \in c$,

$$(\exists y \in \mathcal{D}(f)) [x \neq y \wedge f'x = f'y].$$

Then

$$f'x = \{f'z | zRx\} = \{f'w | wRy\} = f'y.$$

But

$$\begin{aligned} zRx &\rightarrow f'z \in f'x \\ &\rightarrow f'z \in f'y \\ &\rightarrow (\exists w) [wRy \wedge f'w = f'z]. \end{aligned}$$

Since x is an R-minimal element of c and zRx

$$f'w = f'z \rightarrow w = z$$

that is $zRx \rightarrow zRy$.

By a similar argument we obtain $zRy \rightarrow zRx$ and hence from the hypotheses of our theorem we conclude that $x = y$. This is a contradiction.

If

$$F = \cup(K)$$

and if $\langle a, x \rangle \in F \wedge \langle a, y \rangle \in F$ then $(\exists f \in K) (\exists g \in K) [x = f'a \wedge y = g'a]$. But since $a \in \mathcal{D}(f) \cap \mathcal{D}(g)$, $f'a = g'a$ i.e., $x = y$. Therefore F is a function.

Furthermore $(\forall f \in K) (\forall x \in \mathcal{D}(f)) [F'x = f'x]$; consequently

$$F'x = f'x = \{f'y | yRx\} = \{F'y | yRx\}.$$

From this it follows that F is one-to-one for if not there is an x and a y in $\mathcal{D}(F)$ for which $F'x = F'y$ but $x \neq y$. From Theorem 9.4 it then follows that there is an R-minimal x in $\mathcal{D}(F)$ for which $(\exists y \in \mathcal{D}(F)) [x \neq y \wedge F'x = F'y]$. Then

$$F'x = \{F'z | zRx\} = \{F'w | wRy\} = F'y.$$

From this and the defining property of x it then follows that $zRy \leftrightarrow zRx$ and hence $x = y$. This is a contradiction.

Since

$$\mathcal{D}(F) = \bigcup_{f \in K} \mathcal{D}(f)$$

and $f \in K \rightarrow \mathcal{D}(f) \subseteq A$ it follows that $\mathcal{D}(F) \subseteq A$. If $A - \mathcal{D}(F) \neq \emptyset$ then by Theorem 9.4, $A - \mathcal{D}(F)$ has an R-minimal element, that is

$$(\exists x \in A - \mathcal{D}(F)) [A - \mathcal{D}(F) \cap R^{-1}\{x\} = \emptyset].$$

By Theorem 9.3 there is a subset of A that is the R^{-1} closure of $\{x\}$ i.e.

$$(\exists a) [\{x\} \subseteq a \subseteq A \wedge (\forall y) (\forall z) [yRz \wedge z \in a \rightarrow y \in a]].$$

Furthermore each element of a is “connected” to x by a finite R-chain i.e.

$$(\forall y \in a)(\exists n)(\exists f)[f:n+1 \rightarrow a \wedge f(0) = x \wedge f(n) = y \wedge (\forall i < n)[f(i+1)Rf(i)]].$$

Since x is an R-minimal element of $A - \mathcal{D}(F)$

$$A \cap R^{-1}\{x\} \subseteq \mathcal{D}(F).$$

By definition of K if $z \in \mathcal{D}(F)$ then $R^{-1}\{z\} \subseteq \mathcal{D}(F)$. Since each element of a is connected to x by a finite R-chain it follows by induction on the length of such chains that $a - \{x\} \subseteq \mathcal{D}(F)$. We then define

$$g = (F \upharpoonright (a - \{x\})) \cup \{\langle x, F^{-1}\{x\} \rangle\}.$$

Thus $\mathcal{D}(g) \subseteq A \wedge (\forall z \in \mathcal{D}(g)) [z = x \vee z \neq x]$.

If $z = x$ then

$$g^{-1}x = F^{-1}\{x\} = \{F^{-1}y \mid yRx\}.$$

Furthermore $yRx \rightarrow y \in a - \{x\}$ and hence $g^{-1}y = F^{-1}y$. Therefore

$$g^{-1}x = \{g^{-1}y \mid yRx\}.$$

If $z \neq x$ then since $z \in \mathcal{D}(g)$, $g^{-1}z = F^{-1}z$. But

$$F^{-1}z = \{F^{-1}y \mid yRz\}$$

and $yRz \wedge z \in a \rightarrow y \in a$. Furthermore since z is connected to x by a finite R-chain and since R is well founded $y \neq x$. Then $g^{-1}y = F^{-1}y$ and

$$g^{-1}z = \{g^{-1}y \mid yRz\}.$$

Since a is the domain of g and a is closed under R^{-1} it follows that $g \in K$. Hence $x \in a \subseteq \mathcal{D}(F)$. This is a contradiction from which we conclude that $\mathcal{D}(F) = A$.

Thus if $B = F^{-1}A$ then

$$F : A \xrightarrow{\text{onto}} B.$$

Furthermore $a \in b \wedge b \in B \rightarrow (\exists x \in A) [a \in b \wedge b = F^{-1}x]$. But

$$F^{-1}x = \{F^{-1}y \mid yRx\}.$$

Thus $(\exists y) [yRx \wedge a = F'y]$ i.e., $a \in B$ and hence B is transitive.

2) Also $a \in A \wedge b \in A \wedge aRb \rightarrow F'a \in \{F'y \mid yRb\} = F'b$. Therefore

$$F \text{ Isom}_{R,E}(A, B).$$

3) We consider 3) to be the special case of 4) with $n = 0$.

4) (By induction on the number, n , of logical symbols in φ .)
 If $n = 0$ then φ is of the form $a \in b$. Then

$$[A, R] \models a \in b \leftrightarrow aRb.$$

Since $F \text{ Isom}_{R,E}(A, B)$

$$a \in A \wedge b \in A \rightarrow [aRb \leftrightarrow F'a \in F'b].$$

But since $F'a \in B$ and $F'b \in B$ we have

$$a \in A \wedge b \in A \rightarrow [F'a \in F'b \leftrightarrow B \models F'a \in F'b].$$

Therefore

$$a \in A \wedge b \in A \rightarrow [[A, R] \models a \in b \leftrightarrow B \models F'a \in F'b].$$

If φ is of the form $\neg\psi$ and all of the free variables of φ are among a_1, \dots, a_n then so are the free variables of ψ . From the induction hypothesis if $a_1 \in A \wedge \dots \wedge a_n \in A$ then

$$\begin{aligned} [A, R] \models \psi(a_1, \dots, a_n) &\leftrightarrow B \models \psi(F'a_1, \dots, F'a_n) \quad , \\ \neg [A, R] \models \psi(a_1, \dots, a_n) &\leftrightarrow \neg B \models \psi(F'a_1, \dots, F'a_n) \quad , \\ [A, R] \models \neg \psi(a_1, \dots, a_n) &\leftrightarrow B \models \neg \psi(F'a_1, \dots, F'a_n) \quad . \end{aligned}$$

If φ is of the form $\psi \wedge \eta$ and all of the free variables of φ are among a_1, \dots, a_n then so are the free variables of ψ and of η . From the induction hypothesis if $a_1 \in A \wedge \dots \wedge a_n \in A$ then

$$\begin{aligned} [A, R] \models \psi(a_1, \dots, a_n) &\leftrightarrow B \models \psi(F'a_1, \dots, F'a_n) \\ [A, R] \models \eta(a_1, \dots, a_n) &\leftrightarrow B \models \eta(F'a_1, \dots, F'a_n) \quad . \end{aligned}$$

Therefore

$$\begin{aligned} [A, R] \models \psi(a_1, \dots, a_n) \wedge [A, R] \models \eta(a_1, \dots, a_n) &\leftrightarrow B \models \psi(F'a_1, \dots, F'a_n) \\ &\wedge B \models \eta(F'a_1, \dots, F'a_n) \quad , \end{aligned}$$

$$\begin{aligned} [A, R] \models [\psi(a_1, \dots, a_n) \wedge \eta(a_1, \dots, a_n)] &\leftrightarrow B \models [\psi(F'a_1, \dots, F'a_n) \\ &\wedge \eta(F'a_1, \dots, F'a_n)] \quad . \end{aligned}$$

If φ is of the form $(\forall x) \psi(x)$ and if all of the free variables of φ are among a_1, \dots, a_n then there is an x not among a_1, \dots, a_n and all of the

free variables of $\psi(x)$ are among a_1, \dots, a_n, x . From the induction hypothesis if $a_1 \in A \wedge \dots \wedge a_n \in A$ then

$$x \in A \rightarrow [[A, R] \models \psi(x, a_1, \dots, a_n) \leftrightarrow B \models \psi(F'x, F'a_1, \dots, F'a_n)].$$

From the self distributive law for implication

$$[x \in A \rightarrow [A, R] \models \psi(x, a_1, \dots, a_n)] \leftrightarrow [x \in A \rightarrow B \models \psi(F'x, F'a_1, \dots, F'a_n)].$$

Since x is not among a_1, \dots, a_n we have on generalizing and distributing

$$\begin{aligned} & [(\forall x) [x \in A \rightarrow [A, R] \models \psi(x, a_1, \dots, a_n)]] \\ & \leftrightarrow (\forall x) [x \in A \rightarrow B \models \psi(F'x, F'a_1, \dots, F'a_n)] . \end{aligned}$$

Since F maps A one-to-one onto B

$$\begin{aligned} & (\forall x) [x \in A \rightarrow B \models \psi(F'x, F'a_1, \dots, F'a_n)] \\ & \leftrightarrow (\forall x) [x \in B \rightarrow B \models \psi(x, F'a_1, \dots, F'a_n)] . \end{aligned}$$

Therefore

$$[A, R] \models (\forall x) \psi(x, a_1, \dots, a_n) \leftrightarrow B \models (\forall x) \psi(x, F'a_1, \dots, F'a_n) .$$

13 Absoluteness

A basic part of the interpretation of our theory is that each wff $\varphi(x)$ expresses a property that a given individual a has or does not have according as $\varphi(a)$ holds or does not hold. Then $\varphi^A(x)$ expresses the “same” or “corresponding” property for the universe A .

Consider, for example, the existence of an empty set. Earlier we proved that there exists an individual a , called the empty set, having the property

$$(\forall x) [x \notin a].$$

From the Axiom of Regularity it follows that every nonempty class A , as a universe, has this property. In particular, the class of infinite cardinal numbers N' contains an individual a with the property

$$(\forall x \in N') [x \notin a].$$

The set in N' that plays the role of the empty set is \aleph_0 , a set that is far from empty. Thus when viewed from within the universe N' , \aleph_0 is empty but when viewed from “without” i.e., in V , \aleph_0 is not empty.

There are however properties $\varphi(x)$ and universes A such that an individual of A has the property when viewed from within A iff it has the property when viewed from without. Such a property is said to be absolute with respect to A .

Definition 13.1.

- 1) $\varphi \text{ Abs } A \stackrel{\Delta}{\longleftrightarrow} [\varphi^A \longleftrightarrow \varphi]$ if φ is closed.
- 2) $\varphi \text{ Abs } A \stackrel{\Delta}{\longleftrightarrow} a_1 \in A \wedge \cdots \wedge a_n \in A \rightarrow [\varphi^A \longleftrightarrow \varphi]$ where a_1, \dots, a_n is a complete list of all of the free variables in φ .

Theorem 13.2. *If $\varphi \text{ Abs } A \wedge \psi \text{ Abs } A$ then*

- 1) $\varphi \wedge \psi \text{ Abs } A$.
- 2) $\neg \varphi \text{ Abs } A$.

The proofs are left to the reader.

Remark. From Theorem 13.2 we see that if φ and ψ are each absolute with respect to A then $\varphi \vee \psi$, $\varphi \rightarrow \psi$, and $\varphi \longleftrightarrow \psi$ are also absolute with

respect to A . The interesting questions about absoluteness center around quantifiers.

Theorem 13.3. *If $\varphi \text{ Abs } A \wedge \psi \text{ Abs } A$, if $a_1, \dots, a_m, b_1, \dots, b_n$ is a list of distinct variables containing all of the free variables of φ and of ψ , and if*

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge \varphi(a_1, \dots, a_m, b_1, \dots, b_n) \rightarrow a_1 \in A \wedge \dots \wedge a_m \in A$$

then

$$(\forall a_1) \dots (\forall a_m) [\varphi \rightarrow \psi] \text{ Abs } A .$$

Proof. Clearly if $b_1 \in A \wedge \dots \wedge b_n \in A$ then

$$[(\forall a_1) \dots (\forall a_m) [\varphi \rightarrow \psi] \rightarrow (\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi]] .$$

The formal details consist of observing that on the hypothesis

$$(\forall a_1) \dots (\forall a_m) [\varphi \rightarrow \psi]$$

we can deduce $\varphi \rightarrow \psi$ and hence

$$a_m \in A \rightarrow [\varphi \rightarrow \psi] .$$

Then by generalization

$$(\forall a_m) [a_m \in A \rightarrow [\varphi \rightarrow \psi]] .$$

By iteration

$$(\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi] .$$

On the other hand we can deduce from the hypotheses

$$b_1 \in A \wedge \dots \wedge b_n \in A, (\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi], \varphi$$

the following wffs

$$\begin{aligned} & (\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi] , \\ & a_1 \in A \rightarrow (\forall a_2 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi] . \end{aligned}$$

But a basic hypothesis of our theorem is that under the hypothesis listed above

$$a_1 \in A .$$

Hence by modus ponens we can deduce

$$(\forall a_2 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi] .$$

Repeating this we arrive finally at

$$\psi$$

from which one application of the deduction theorem gives that on the hypotheses

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge (\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi]$$

we can deduce

$$\varphi \rightarrow \psi .$$

Then by generalization we deduce

$$(\forall a_1) \dots (\forall a_m) [\varphi \rightarrow \psi]$$

and finally, by the deduction theorem,

$$b_1 \in A \wedge \dots \wedge b_n \in A \rightarrow [(\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi] \\ \rightarrow (\forall a_1) \dots (\forall a_m) [\varphi \rightarrow \psi]] .$$

We then have

$$b_1 \in A \wedge \dots \wedge b_n \in A \rightarrow [(\forall a_1) \dots (\forall a_m) [\varphi \rightarrow \psi] \\ \leftrightarrow (\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi]] .$$

Since φ and ψ are each absolute w.r.t. A

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge a_1 \in A \wedge \dots \wedge a_m \in A \rightarrow [\varphi \leftrightarrow \varphi^A] ,$$

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge a_1 \in A \wedge \dots \wedge a_m \in A \rightarrow [\psi \leftrightarrow \psi^A] .$$

Hence by generalization, properties of equivalence, and the self distributive law if $b_1 \in A \wedge \dots \wedge b_n \in A$ then

$$[(\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi \rightarrow \psi] \leftrightarrow (\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi^A \rightarrow \psi^A]] .$$

We then conclude that if $b_1 \in A \wedge \dots \wedge b_n \in A$ then

$$[(\forall a_1) \dots (\forall a_m) [\varphi \rightarrow \psi] \leftrightarrow (\forall a_1 \in A) \dots (\forall a_m \in A) [\varphi^A \rightarrow \psi^A]] .$$

Corollary 13.4. *If $\varphi \text{ Abs } A \wedge \psi \text{ Abs } A$, if $a_1, \dots, a_m, b_1, \dots, b_n$ is a list of distinct variables containing all of the free variables in φ and in ψ , and if*

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge \varphi(a_1, \dots, a_m, b_1 \dots b_n) \rightarrow a_1 \in A \wedge \dots \wedge a_m \in A ,$$

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge \psi(a_1, \dots, a_m, b_1 \dots b_n) \rightarrow a_1 \in A \wedge \dots \wedge a_m \in A$$

then $(\forall a_1) \dots (\forall a_m) [\varphi \leftrightarrow \psi] \text{ Abs } A$.

The proof is left to the reader.

Theorem 13.5. *If $\varphi \text{ Abs } A$, if $a_1, \dots, a_m, b_1, \dots, b_n$ is a list of distinct variables containing all of the free variables in φ and if*

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge \varphi(a_1, \dots, a_m, b_1, \dots, b_n) \rightarrow a_1 \in A \wedge \dots \wedge a_m \in A$$

then

$$(\exists a_1) \dots (\exists a_m) \varphi \text{ Abs } A .$$

Proof. Since

$$b_1 \in A \wedge \dots \wedge b_n \in A \wedge \varphi(a_1, \dots, a_m, b_1, \dots, b_n) \rightarrow a_1 \in A \wedge \dots \wedge a_m \in A$$

we have that

$$b_1 \in A \wedge \cdots \wedge b_n \in A \rightarrow [(\exists a_1) \dots (\exists a_m) \varphi \leftrightarrow (\exists a_1 \in A) \dots (\exists a_m \in A) \varphi].$$

Since φ is absolute w.r.t. A

$$b_1 \in A \wedge \cdots \wedge b_n \in A \wedge a_1 \in A \wedge \cdots \wedge a_m \in A \rightarrow [\varphi \leftrightarrow \varphi^A].$$

Therefore if $b_1 \in A \wedge \cdots \wedge b_n \in A$ then

$$[(\exists a_1 \in A) \dots (\exists a_m \in A) \varphi \leftrightarrow (\exists a_1 \in A) \dots (\exists a_m \in A) \varphi^A]$$

and hence

$$[(\exists a_1) \dots (\exists a_m) \varphi \leftrightarrow (\exists a_1 \in A) \dots (\exists a_m \in A) \varphi^A].$$

Theorem 13.6. *If $\vdash [\varphi \leftrightarrow \psi]$ and if A is a nonempty class that satisfies each nonlogical axiom in some proof of $\varphi \leftrightarrow \psi$ then*

$$\varphi \text{ Abs } A \leftrightarrow \psi \text{ Abs } A.$$

Proof. If all of the free variables of φ and of ψ are among b_1, \dots, b_n then by Theorem 12.6

$$b_1 \in A \wedge \cdots \wedge b_n \in A \rightarrow [\varphi^A \leftrightarrow \psi^A].$$

Therefore if $b_1 \in A \wedge \cdots \wedge b_n \in A$

$$[\varphi \leftrightarrow \varphi^A] \leftrightarrow [\psi \leftrightarrow \psi^A].$$

Theorem 13.7. *If $\vdash (\exists x) \varphi(x)$, if A is a nonempty class that satisfies each nonlogical axiom in some proof of $(\exists x) \varphi(x)$, and if $\varphi(x) \text{ Abs } A$ then*

$$(\exists x) \varphi(x) \text{ Abs } A.$$

Proof. If all of the free variables of $(\exists x) \varphi(x)$ are among a_1, \dots, a_n then by Theorem 12.6

$$a_1 \in A \wedge \cdots \wedge a_n \in A \rightarrow (\exists x \in A) \varphi^A(x).$$

Then

$$a_1 \in A \wedge \cdots \wedge a_n \in A \rightarrow [(\exists x) \varphi(x) \rightarrow (\exists x \in A) \varphi^A(x)].$$

Furthermore there exists an x distinct from a_1, \dots, a_n . Then since $\varphi(x)$ is absolute with respect to A .

$$a_1 \in A \wedge \cdots \wedge a_n \in A \wedge x \in A \rightarrow [\varphi(x) \leftrightarrow \varphi^A(x)].$$

Therefore

$$\begin{aligned} a_1 \in A \wedge \cdots \wedge a_n \in A \rightarrow & [(\exists x \in A) \varphi(x) \leftrightarrow (\exists x \in A) \varphi^A(x)] \\ & \rightarrow [(\exists x \in A) \varphi^A(x) \rightarrow (\exists x) \varphi(x)]. \end{aligned}$$

Theorem 13.8. *If $\vdash [\varphi \leftrightarrow (\forall x) \psi(x)]$ and $\vdash [\varphi \leftrightarrow (\exists x) \eta(x)]$, if A is a nonempty class that satisfies each nonlogical axiom in some proof of*

$\varphi \longleftrightarrow (\forall x) \psi(x)$ and some proof of $\varphi \longleftrightarrow (\exists x) \eta(x)$, and if $\psi(x) \text{ Abs } A \wedge \eta(x) \text{ Abs } A$ then

$$\varphi \text{ Abs } A .$$

Proof. If all of the variables of φ , $(\forall x) \psi(x)$, and $(\exists x) \eta(x)$ are among a_1, \dots, a_n then by Theorem 12.6

$$a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [\varphi^A \longleftrightarrow (\forall x \in A) \psi^A(x)] ,$$

$$a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [\varphi^A \longleftrightarrow (\exists x \in A) \eta^A(x)] .$$

Also since $\psi(x)$ and $\eta(x)$ are absolute

$$a_1 \in A \wedge \dots \wedge a_n \in A \wedge x \in A \rightarrow [\psi(x) \longleftrightarrow \psi^A(x)] ,$$

$$a_1 \in A \wedge \dots \wedge a_n \in A \wedge x \in A \rightarrow [\eta(x) \longleftrightarrow \eta^A(x)] .$$

From this, choosing x distinct from a_1, \dots, a_n

$$a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [(\forall x \in A) \psi(x) \longleftrightarrow (\forall x \in A) \psi^A(x)] ,$$

$$a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [(\exists x \in A) \eta(x) \longleftrightarrow (\exists x \in A) \eta^A(x)] .$$

Then since $\varphi \longleftrightarrow (\forall x) \psi(x)$ is a theorem

$$\begin{aligned} a_1 \in A \wedge \dots \wedge a_n \in A \wedge \varphi &\rightarrow (\forall x) \psi(x) \\ &\rightarrow (\forall x \in A) \psi(x) \\ &\rightarrow (\forall x \in A) \psi^A(x) \\ &\rightarrow \varphi^A . \end{aligned}$$

Also

$$\begin{aligned} a_1 \in A \wedge \dots \wedge a_n \in A \wedge \varphi^A &\rightarrow (\exists x \in A) \eta^A(x) \\ &\rightarrow (\exists x \in A) \eta(x) \\ &\rightarrow (\exists x) \eta(x) \\ &\rightarrow \varphi . \end{aligned}$$

Remark. Satisfaction and absoluteness have been defined for wffs. Most of our theorems in ZF are however wffs in the wider sense (Def. 4.1). It is therefore convenient to extend our definitions to wffs in the wider sense.

Definition 13.9. If φ is a wff in the wider sense then

- 1) $[A, R] \models \varphi \xleftrightarrow{\Delta} [A, R] \models \varphi^*$.
- 2) $\varphi^A \triangleq (\varphi^*)^A$.
- 3) $\varphi \text{ Abs } A \xleftrightarrow{\Delta} \varphi^* \text{ Abs } A$.

It would however be helpful to be able to determine the absoluteness of wffs in the wider sense without first reducing them to primitive terms. For this purpose the following substitution theorem is useful.

Theorem 13.10. If $A \neq 0$, if $\varphi(b_1, \dots, b_n) \text{ Abs } A$ and if $\mathcal{M}(B_1) \wedge \dots \wedge \mathcal{M}(B_n) \wedge b_1 = B_1 \text{ Abs } A \wedge \dots \wedge b_n = B_n \text{ Abs } A$ then $\varphi(B_1, \dots, B_n) \text{ Abs } A$.

Proof (by induction on n). If $n = 1$ then

$$\begin{aligned} \varphi(B_1) &\longleftrightarrow (\forall b_1) [b_1 = B_1 \rightarrow \varphi(b_1)] \\ &\longleftrightarrow (\exists b_1) [b_1 = B_1 \wedge \varphi(b_1)]. \end{aligned}$$

If $\varphi(b_1) \text{ Abs } A \wedge b_1 = B_1 \text{ Abs } A$ then $[b_1 = B_1 \rightarrow \varphi(b_1)] \text{ Abs } A$ and $[b_1 = B_1 \wedge \varphi(b_1)] \text{ Abs } A$. Then by Theorem 13.8, $\varphi(B_1) \text{ Abs } A$.

The induction step is obvious and hence omitted.

Definition 13.11.

- 1) $x^A = x \cap A$.
- 2) $\{x | \varphi(x)\}^A \triangleq \{x \in A | \varphi^A(x)\}$.

Theorem 13.12. If A is transitive and $x \in A$ then

- 1) $x^A = x$.
- 2) $[x \in y]^A \longleftrightarrow x^A \in y^A$.
- 3) $[[x \in \{y | \varphi(y)\}]^A \longleftrightarrow [x^A \in \{y | \varphi(y)\}^A]]$.
- 4) $[[\{y | \varphi(y)\} \in x]^A \longleftrightarrow [\{y | \varphi(y)\}^A \in x^A]]$.
- 5) $[\{x | \varphi(x)\} \in \{y | \psi(y)\}]^A \longleftrightarrow [\{x | \varphi(x)\}^A \in \{y | \psi(y)\}^A]$.

Proof.

- 1) If A is transitive and $x \in A$ then $x \subseteq A$. Therefore $x^A = x \cap A = x$.
- 2) Obvious from 1).
- 3) $[x \in \{y | \varphi(y)\}]^* \longleftrightarrow \varphi(x)$.

Hence

$$[x \in \{y | \varphi(y)\}]^A \longleftrightarrow \varphi^A(x).$$

Then $x \in A$ implies

$$\begin{aligned} [x \in \{y | \varphi(y)\}]^A &\longleftrightarrow x \in A \wedge \varphi^A(x) \\ &\longleftrightarrow x \in \{y \in A | \varphi^A(y)\} \\ &\longleftrightarrow x^A \in \{y | \varphi(y)\}^A \end{aligned}$$

- 4) $[\{y | \varphi(y)\} \in x]^* \longleftrightarrow (\exists z) [z \in x \wedge (\forall y) [y \in z \longleftrightarrow \varphi(y)]]$.

Hence

$$[\{y | \varphi(y)\} \in x]^A \longleftrightarrow (\exists z \in A) [z \in x \wedge (\forall y \in A) [y \in z \longleftrightarrow \varphi^A(y)]] .$$

Then A transitive and $x \in A$ implies

$$\begin{aligned} [\{y|\varphi(y)\} \in x]^A &\longleftrightarrow (\exists z \in A) [z \in x \wedge (\forall y \in A) [y \in x \longleftrightarrow \varphi^A(y)]] \\ &\longleftrightarrow (\exists z \in A) [z \in x \wedge z = \{y \in A | \varphi^A(y)\}] \\ &\longleftrightarrow \{y \in A | \varphi^A(y)\} \in x \\ &\longleftrightarrow \{y|\varphi(y)\}^A \in x^A. \end{aligned}$$

$$\begin{aligned} 5) \quad &[\{x|\varphi(x)\} \in \{y|\psi(y)\}]^* \\ &\longleftrightarrow (\exists z) [(\forall x) [x \in z \longleftrightarrow \varphi(x)] \wedge z \in \{y|\psi(y)\}]. \end{aligned}$$

Therefore from 2)

$$\begin{aligned} &[\{x|\varphi(x)\} \in \{y|\psi(y)\}]^A \\ &\longleftrightarrow (\exists z \in A) (\forall x \in A) [x \in z \longleftrightarrow \varphi^A(x) \wedge z \in \{y|\psi(y)\}^A] \\ &\longleftrightarrow (\exists z \in A) [z = \{x \in A | \varphi^A(x)\} \wedge z \in \{y|\psi(y)\}^A] \\ &\longleftrightarrow \{x \in A | \varphi^A(x)\} \in \{y|\psi(y)\}^A \\ &\longleftrightarrow \{x|\varphi(x)\}^A \in \{y|\psi(y)\}^A. \end{aligned}$$

Definition 13.13.

- 1) $B \text{ Abs } A \xleftrightarrow{\Delta} [B^A = B]$ if B is a term containing no free variables.
- 2) $B \text{ Abs } A \xleftrightarrow{\Delta} a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [B^A = B]$ where a_1, \dots, a_n is a complete list of all the free variables in B .

Theorem 13.14. *If $\varphi(x) \text{ Abs } A$ and if all of the free variables of $\varphi(x)$ are among a_1, \dots, a_n, x then*

$$a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [\{x|\varphi(x)\}^A = \{x|\varphi(x)\} \cap A].$$

Proof. If $a_1 \in A \wedge \dots \wedge a_n \in A$ then since $\varphi(x)$ is absolute with respect to A

$$\begin{aligned} x \in \{x|\varphi(x)\}^A &\longleftrightarrow x \in A \wedge \varphi^A(x) \\ &\longleftrightarrow x \in A \wedge \varphi(x) \\ &\longleftrightarrow x \in \{x|\varphi(x)\} \cap A. \end{aligned}$$

Remark. The class $\{x|\varphi(x)\}^A$ is the class $\{x|\varphi(x)\}$ relativized to A . By definition $\{x|\varphi(x)\}^A$ is the class of individuals in A for which $\varphi^A(x)$ holds. From Theorem 13.14 we see that if $\varphi(x)$ is absolute with respect to A then $\{x|\varphi(x)\}^A$ is simply the class of individuals in A for which $\varphi(x)$ holds.

Theorem 13.15. *If A is nonempty and transitive if*

$$\begin{aligned} \mathcal{M}(\{y|\varphi(y)\}) \text{ and if } \mathcal{M}(\{y|\varphi(y)\}) \text{ Abs } A \text{ then } [x = \{y|\varphi(y)\}] \text{ Abs } A \\ \text{iff } \{y|\varphi(y)\} \text{ Abs } A. \end{aligned}$$

Proof. If all of the free variables of $\varphi(y)$ are among a_1, \dots, a_n, y then

$[x = \{y | \varphi(y)\}] \text{ Abs } A$

$\longleftrightarrow a_1 \in A \wedge \dots \wedge a_n \in A \wedge x \in A \rightarrow [(\forall y) [y \in x \longleftrightarrow \varphi(y)]]$

$\longleftrightarrow (\forall y \in A) [y \in x \longleftrightarrow \varphi^A(y)]$

$\longleftrightarrow a_1 \in A \wedge \dots \wedge a_n \in A \wedge x \in A \rightarrow [(\forall y) [y \in x \longleftrightarrow \varphi(y)]]$

$\longleftrightarrow (\forall y) [y \in x \longleftrightarrow y \in A \wedge \varphi^A(y)]$

$\longleftrightarrow a_1 \in A \wedge \dots \wedge a_n \in A \wedge x \in A \rightarrow [x = \{y | \varphi(y)\} \longleftrightarrow x = \{y \in A | \varphi^A(y)\}]$

$\longleftrightarrow a_1 \in A \wedge \dots \wedge a_n \in A \rightarrow [\{y | \varphi(y)\} = \{y | \varphi(y)\}^A]$

$\longleftrightarrow \{y | \varphi(y)\} \text{ Abs } A .$

Remark. We turn now to the problem of establishing the absoluteness properties of certain wffs and terms. Our ultimate goal is to find conditions on A that will assure us that A is a standard transitive model of ZF.

Theorem 13.16. *If φ is quantifier free then $\varphi \text{ Abs } A$.*

Proof. If φ is quantifier free then $\varphi^A \longleftrightarrow \varphi$.

Theorem 13.17. *$a \in b \text{ Abs } A$.*

Proof. $a \in b$ is quantifier free.

Theorem 13.18. *If A is nonempty and transitive then*

1) $a \subseteq b \text{ Abs } A$.

2) $a = b \text{ Abs } A$.

Proof.

1) $a \subseteq b \longleftrightarrow (\forall x) [x \in a \rightarrow x \in b]$.

Since A is transitive $a \in A \rightarrow a \subseteq A$, i.e., $x \in a \wedge a \in A \rightarrow x \in A$. From Theorems 13.17, 13.3, and 13.6 it then follows that

$a \subseteq b \text{ Abs } A .$

2) $a = b \longleftrightarrow a \subseteq b \wedge b \subseteq a$.

Remark. The requirement in Theorem 13.18 that A be transitive cannot be dropped. For example if $A = \{0, 1, \{0, 1, 2\}, \{0, 1, 3\}\}$ then from an internal vantage point the sets $\{0, 1, 2\}$ and $\{0, 1, 3\}$ are indistinguishable i.e.

$(\forall x \in A) [x \in \{0, 1, 2\} \longleftrightarrow x \in \{0, 1, 3\}] .$

Since the membership property is absolute with respect to any nonempty class (Theorem 13.17) if $b \in A$ then those individuals in A that play the role of elements of b are individuals in V that *are* elements

in b . But not conversely. In the foregoing example we have relative to A

$$0 \underset{A}{\in} \{0, 1, 2\}, \quad 1 \underset{A}{\in} \{0, 1, 2\}, \quad 2 \underset{A}{\notin} \{0, 1, 2\}.$$

Similarly with subsets, if A is a nonempty transitive class then containment is absolute with respect to A . This means that if $b \in A$ then every element of A that is a subset of b relative to A is a subset of b in the “real” universe V . But not conversely. There may be a subset of b that is not an element of A . Indeed if A is transitive but not supertransitive there must be at least one element of A having a subset that is not in A .

Theorem 13.19. *If A is nonempty and transitive then*

- 1) $0 \text{ Abs } A$,
- 2) $[a \cup b] \text{ Abs } A$,
- 3) $\{a, b\} \text{ Abs } A$,
- 4) $\cup(a) \text{ Abs } A$.
- 5) $[a - b] \text{ Abs } A$.

Proof. 1) Since $x \neq x \text{ Abs } A$ we have

$$0^A = \{x \in A \mid [x \neq x]^A\} = \{x \in A \mid x \neq x\} = 0.$$

- 2) If $a \in A \wedge b \in A$ then

$$\begin{aligned} [a \cup b]^A &= \{x \in A \mid [x \in a \vee x \in b]^A\} \\ &= \{x \in A \mid x \in a \vee x \in b\} \\ &= \{x \mid x \in a \vee x \in b\} \\ &= a \cup b. \end{aligned}$$

- 3) If $a \in A \wedge b \in A$ then

$$\begin{aligned} \{a, b\}^A &= \{x \in A \mid [x = a \vee x = b]^A\} \\ &= \{x \mid x = a \vee x = b\} \\ &= \{a, b\}. \end{aligned}$$

- 4) If $a \in A$ then

$$\begin{aligned} [\cup(a)]^A &= \{x \in A \mid (\exists y \in A) [x \in y \wedge y \in a]^A\} \\ &= \{x \in A \mid (\exists y \in A) [x \in y \wedge y \in a]\} \\ &= \{x \mid (\exists y) [x \in y \wedge y \in a]\} \\ &= \cup(a). \end{aligned}$$

5) If $a \in A \wedge b \in A$ then

$$\begin{aligned} [a - b]^A &= \{x \in A \mid [x \in a \wedge x \notin b]^A\} \\ &= \{x \in A \mid x \in a \wedge x \notin b\} \\ &= a - b. \end{aligned}$$

Remark. The proofs of several of the theorems to follow are similar to the proof of Theorem 13.18 involving repeated applications of foregoing theorems on absoluteness. To avoid rather dull repetitions we omit most of the details.

Theorem 13.20. *If A is nonempty and transitive then*

- 1) $\text{Tr}(a) \text{ Abs } A$,
- 2) $\text{Ord}(a) \text{ Abs } A$.

Proof.

- 1) $\text{Tr}(a) \longleftrightarrow (\forall x) [x \in a \rightarrow x \subset a]$.
- 2) $\text{Ord}(a) \longleftrightarrow \text{Tr}(a) \wedge (\forall x) (\forall y) [x \in a \wedge y \in a \rightarrow x \in y \vee x = y \vee y \in x]$.

Remark. Theorem 13.20 assures us that restricting the definition of ordinal number to a nonempty transitive class does not enable any new objects to qualify as ordinals. Consequently if A is a standard transitive model of ZF then the class of ordinals “in” A is a subclass of On , i.e.

$$\text{On}^A = \{x \in A \mid [\text{Ord}(x)]^A\} = \{x \in A \mid \text{Ord}(x)\} = \text{On} \cap A.$$

Theorem 13.21. *If A is nonempty and transitive then*

- 1) $[a \in \omega] \text{ Abs } A$,
- 2) $[a = \omega] \text{ Abs } A$ if $\omega \subseteq A$.

Proof.

- 1) $a \in \omega \longleftrightarrow a \cup \{a\} \subseteq K_1$
 $\longleftrightarrow (\forall x) [x \in a \vee x = a \rightarrow x \in K_1]$
 $\longleftrightarrow (\forall x) [x \in a \vee x = a \rightarrow x = 0 \vee (\exists y) [\text{Ord}(y) \wedge x = y \cup \{y\}]]$.
- 2) $a = \omega \longleftrightarrow (\forall x) [x \in a \longleftrightarrow x \in \omega]$.

Theorem 13.22. *If A is nonempty and transitive then*

- 1) $[\alpha < \beta] \text{ Abs } A$,
- 2) $[\alpha = \beta] \text{ Abs } A$,
- 3) $[\gamma = \max(\alpha, \beta)] \text{ Abs } A$.

Proof.

- 1) $[\alpha < \beta] \iff \text{Ord}(\alpha) \wedge \text{Ord}(\beta) \wedge \alpha \in \beta.$
- 2) $[\alpha = \beta] \iff \text{Ord}(\alpha) \wedge \text{Ord}(\beta) \wedge \alpha = \beta.$
- 3) $[\gamma = \max(\alpha, \beta)] \iff \text{Ord}(\gamma) \wedge \text{Ord}(\alpha) \wedge \text{Ord}(\beta) \wedge \gamma = \alpha \cup \beta.$

Theorem 13.23. *If A is nonempty and transitive then*

- 1) $[\langle \alpha, \beta \rangle \text{Le} \langle \gamma, \delta \rangle] \text{ Abs } A,$
- 2) $[\langle \alpha, \beta \rangle R_0 \langle \gamma, \delta \rangle] \text{ Abs } A.$

Proof.

- 1) $[\langle \alpha, \beta \rangle \text{Le} \langle \gamma, \delta \rangle] \iff [\alpha < \gamma \vee [\alpha = \gamma \wedge \beta < \delta]].$
- 2) $[\langle \alpha, \beta \rangle R_0 \langle \gamma, \delta \rangle] \iff [\max(\alpha, \beta) < \max(\gamma, \delta) \vee [\max(\alpha, \beta) = \max(\gamma, \delta) \wedge \langle \alpha, \beta \rangle \text{Le} \langle \gamma, \delta \rangle]].$

Exercises. *In Exercises 1–29 determine whether or not the given predicate is absolute with respect to A , A being nonempty and transitive.*

- | | |
|---|--|
| 1) $\mathcal{M}(a).$ | 16) $f : a \xrightarrow{\text{onto}} b.$ |
| 2) $\mathcal{P}_r(a).$ | 17) $f \text{ Isom}_{r_1, r_2}(a_1, a_2).$ |
| 3) $\mathcal{R}el(a).$ | 18) $\text{Orf}(f).$ |
| 4) $\mathcal{U}n(a).$ | 19) $\text{Smo}(f).$ |
| 5) $\mathcal{U}n_2(a).$ | 20) $a \simeq b.$ |
| 6) $\mathcal{F}nc(a).$ | 21) $\text{Fin}(a).$ |
| 7) $\mathcal{F}nc_2(a).$ | 22) $\text{Inf}(a).$ |
| 8) $a \mathcal{F}n b.$ | 23) $\text{cof}(\alpha, \beta).$ |
| 9) $a \mathcal{F}n_2 b.$ | 24) $\text{Reg}(\alpha).$ |
| 10) $r \text{ Fr } a.$ | 25) $\text{Inacc}_w(\aleph_\alpha).$ |
| 11) $r \text{ Wfr } a.$ | 26) $\text{Inacc}(\aleph_\alpha).$ |
| 12) $r \text{ We } a.$ | 27) $\text{Cl}(r, a).$ |
| 13) $f : a \rightarrow b.$ | 28) $\text{Cl}_2(r, a).$ |
| 14) $f : a \xrightarrow{1-1} b.$ | 29) $\text{St}(a).$ |
| 15) $f : a \xrightarrow[1-1]{\text{onto}} b.$ | |

In Exercise 30–55 determine whether or not the given term is absolute with respect to A , A nonempty and transitive.

- | | |
|---------------------------|-----------------------|
| 30) $x = a \cap b.$ | 32) $x = \cap(a).$ |
| 31) $x = \mathcal{P}(a).$ | 33) $x = a \times b.$ |

- | | | | |
|-----|----------------------------|-----|--------------------------------|
| 34) | $x = a^{-1}.$ | 45) | $\gamma = \alpha \cdot 1.$ |
| 35) | $x = \mathcal{D}(a).$ | 46) | $\gamma = \alpha \cdot \beta.$ |
| 36) | $x = \mathcal{W}(a).$ | 47) | $\gamma = \alpha^0.$ |
| 37) | $x = a \Gamma b.$ | 48) | $\gamma = \alpha^1.$ |
| 38) | $x = a^{\ast} b.$ | 49) | $\gamma = \alpha^{\beta}.$ |
| 39) | $x = a \circ b.$ | 50) | $\alpha = \bar{\bar{a}}.$ |
| 40) | $x = a^{\ast} b.$ | 51) | $f = \aleph.$ |
| 41) | $\gamma = \alpha + 0.$ | 52) | $c = a^b.$ |
| 42) | $\gamma = \alpha + 1.$ | 53) | $\beta = \text{cf}(\alpha).$ |
| 43) | $\gamma = \alpha + \beta.$ | 54) | $x = R(\alpha).$ |
| 44) | $\gamma = \alpha \cdot 0.$ | 55) | $\alpha = \text{rank}(x).$ |

Remark. We turn now to an investigation of conditions on a class A that are necessary for A to be a model of ZF, that is, for A to be a model of the followings wffs.

Axiom 1. (*Extensionality*) $(\forall a) (\forall x) (\forall y) [x = y \wedge x \in a \rightarrow y \in a].$

Axiom 2. (*Pairing*) $(\forall a) (\forall b) \mathcal{M}(\{a, b\}).$

Axiom 3. (*Unions*) $(\forall a) \mathcal{M}(\cup(a)).$

Axiom 4. (*Powers*) $(\forall a) \mathcal{M}(\mathcal{P}(a)).$

Axiom 5. (*Schema of Replacement*)

$$(\forall a) [(\forall u) (\forall v) (\forall w) [\varphi(u, v) \wedge \varphi(u, w) \rightarrow v = w] \\ \rightarrow (\exists b) (\forall y) [y \in b \leftrightarrow (\exists x) [x \in a \wedge \varphi(x, y)]]].$$

Axiom 6. (*Regularity*) $(\forall a) [a \neq 0 \rightarrow (\exists x) [x \in a \wedge x \cap a = 0]].$

Axiom 7. (*Infinity*) $\mathcal{M}(\omega).$

Theorem 13.24. $A \neq 0 \wedge \text{Tr}(A) \rightarrow \text{STM}(A, \text{Ax 1}).$

Proof. $\text{Ax. 1} \leftrightarrow (\forall a) (\forall x) (\forall y) [x = y \wedge x \in a \rightarrow y \in a].$ Since Axiom 1 holds for each x, y and a , in particular it holds for each x, y and a in A i.e.,

$$(\forall a \in A) (\forall x \in A) (\forall y \in A) [x = y \wedge x \in a \rightarrow y \in a].$$

Since $x = y \wedge x \in a \rightarrow y \in a$ is absolute with respect to A

$$a \in A \wedge x \in A \wedge y \in A \rightarrow [[x = y \wedge x \in a \rightarrow y \in a] \\ \leftrightarrow [x = y \wedge x \in a \rightarrow y \in a]^A].$$

Therefore

$$(\forall a \in A) (\forall x \in A) (\forall y \in A) [x = y \wedge x \in a \rightarrow y \in a]^A.$$

But this is Axiom 1 relativized to A i.e., $[\text{Ax. 1}]^A$.

Since Axiom 1 is closed we have from Theorem 12.5

$$A \models \text{Ax. 1} \longleftrightarrow [\text{Ax. 1}]^A.$$

Hence A satisfies Axiom 1, i.e.,

$$A \models \text{Ax. 1}.$$

Since A is nonempty and transitive we conclude that A is a standard transitive model of Axiom 1.

Theorem 13.25. *If A is nonempty and transitive then A is a standard transitive model of the Axiom of Pairing iff*

$$(\forall a \in A) (\forall b \in A) [\{a, b\} \in A].$$

Proof. $\text{Ax. 2} \longleftrightarrow (\forall a) (\forall b) (\exists x) [x = \{a, b\}]$.

Since A is nonempty and transitive A is a standard transitive model of Axiom 2 if and only if

$$(\forall a \in A) (\forall b \in A) (\exists x \in A) [x = \{a, b\}]^A.$$

However since $x = \{a, b\}$ is absolute w.r.t. A , this holds if and only if

$$(\forall a \in A) (\forall b \in A) (\exists x \in A) [x = \{a, b\}]$$

i.e., if and only if

$$(\forall a \in A) (\forall b \in A) [\{a, b\} \in A].$$

Remark. Earlier we proved that the empty set, the union of two sets, unordered pairs, and the union of a set are absolute terms with respect to a nonempty transitive class A . Curiously ordered pairs are not absolute with respect to such classes.

Suppose that A is the nonempty transitive class $\{0, 1\}$ then

$$\begin{aligned} [\langle 0, 1 \rangle]^A &= \{x \in A \mid [x = \{0\} \vee x = \{0, 1\}]^A\} \\ &= \{x \in A \mid x = \{0\} \vee x = \{0, 1\}\} = \{\{0\}\} = \langle 0, 0 \rangle. \end{aligned}$$

This pathology disappears if A is a nonempty transitive model of the Axiom of Pairing.

Theorem 13.26. $\text{STM}(A, \text{Ax. 2}) \rightarrow \langle a, b \rangle \text{ Abs } A$.

Proof. If A is a standard transitive model of Axiom 2 and if $a \in A \wedge b \in A$ then $\{a\} \in A$ and $\{a, b\} \in A$. Then

$$\begin{aligned} [\langle a, b \rangle]^A &= \{x \in A \mid [x = \{a\} \vee x = \{a, b\}]^A\} \\ &= \{x \in A \mid x = \{a\} \vee x = \{a, b\}\} \\ &= \{x \mid x = \{a\} \vee x = \{a, b\}\} \\ &= \langle a, b \rangle. \end{aligned}$$

Remark. From Theorem 13.26 and the substitution property, Theorem 13.10 it follows that ordered triples, ordered quadruples, etc. are absolute with respect to standard transitive models of the Axiom of Pairing.

Theorem 13.27. *If A is a nonempty, transitive model of the Axiom of Pairing then*

- 1) $\mathcal{R}el(a) \text{ Abs } A.$
- 2) $\mathcal{U}n(a) \text{ Abs } A.$
- 3) $\mathcal{U}n_2(a) \text{ Abs } A.$
- 4) $\mathcal{F}nc(a) \text{ Abs } A.$
- 5) $\mathcal{F}nc_2(a) \text{ Abs } A.$

Proof.

- 1) $\mathcal{R}el(a) \longleftrightarrow (\forall x) [x \in a \rightarrow (\exists y) (\exists z) [x = \langle y, z \rangle]].$
- 2) $\mathcal{U}n(a) \longleftrightarrow (\forall x) (\forall y) (\forall z) [\langle x, y \rangle \in a \wedge \langle x, z \rangle \in a \rightarrow y = z].$
- 3) $\mathcal{U}n_2(a) \longleftrightarrow \mathcal{U}n(a)$
 $\wedge (\forall x) (\forall y) (\forall z) [\langle x, z \rangle \in a \wedge \langle y, z \rangle \in a \rightarrow x = y].$
- 4) $\mathcal{F}nc(a) \longleftrightarrow \mathcal{R}el(a) \wedge \mathcal{U}n(a).$
- 5) $\mathcal{F}nc_2(a) \longleftrightarrow \mathcal{R}el(a) \wedge \mathcal{U}n_2(a).$

Theorem 13.28. *If A is a nonempty transitive model of the Axiom of Pairing then*

- 1) $a \times b \text{ Abs } A.$
- 2) $a^{-1} \text{ Abs } A.$
- 3) $\mathcal{D}(a) \text{ Abs } A.$
- 4) $\mathcal{W}(a) \text{ Abs } A.$
- 5) $a \cdot b \text{ Abs } A.$
- 6) $a \text{ `` } b \text{ Abs } A.$
- 7) $a \Gamma b \text{ Abs } A.$

Proof. If $a \in A \wedge b \in A$ then

- 1) $[a \times b]^A = \{x \in A \mid (\exists y \in A) (\exists z \in A) [y \in a \wedge z \in b \wedge x = \langle y, z \rangle]^A\}$
 $= \{x \in A \mid (\exists y \in A) (\exists z \in A) [y \in a \wedge z \in b \wedge x = \langle y, z \rangle]\}$
 $= \{x \mid (\exists y) (\exists z) [y \in a \wedge z \in b \wedge x = \langle y, z \rangle]\}$
 $= a \times b.$

- 2) $[a^{-1}]^A = \{x \in A \mid (\exists y \in A) (\exists z \in A) [x = \langle y, z \rangle \wedge \langle z, y \rangle \in a]^A\}$
 $= \{x \in A \mid (\exists y \in A) (\exists z \in A) [x = \langle y, z \rangle \wedge \langle z, y \rangle \in a]\}$
 $= \{x \in A \mid (\exists y) (\exists z) [x = \langle y, z \rangle \wedge \langle z, y \rangle \in a]\}$
 $= \{x \mid (\exists y) (\exists z) [x = \langle y, z \rangle \wedge \langle z, y \rangle \in a]\}$
 $= a^{-1}.$
- 3) $[\mathcal{D}(a)]^A = \{x \in A \mid (\exists y \in A) [\langle x, y \rangle \in a]^A\}$
 $= \{x \in A \mid (\exists y \in A) [\langle x, y \rangle \in a]\}$
 $= \{x \mid (\exists y) [\langle x, y \rangle \in a]\}$
 $= \mathcal{D}(a).$
- 4) $[\mathcal{W}(a)]^A = \{x \in A \mid (\exists y \in A) [\langle y, x \rangle \in a]^A\} = \mathcal{W}(a).$
- 5) $[a' b]^A = \{x \in A \mid (\exists y \in A) [x \in y \wedge \langle b, y \rangle \in a]^A \wedge (\exists! y \in A) [\langle b, y \rangle \in a]^A\}$
 $= \{x \mid (\exists y) [x \in y \wedge \langle b, y \rangle \in a] \wedge (\exists! y) [\langle b, y \rangle \in a]\}.$
 $= a' b.$
- 6) $[a'' b]^A = \{x \in A \mid (\exists y \in A) [y \in b \wedge \langle y, x \rangle \in a]^A\}$
 $= \{x \mid (\exists y) [y \in b \wedge \langle y, x \rangle \in a]\} = a'' b.$
- 7) The proof is left to the reader.

Theorem 13.29. *If A is a nonempty transitive model of the Axiom of Pairing then*

- 1) $f \mathcal{F}n a \text{ Abs } A.$
- 2) $f \mathcal{F}n_2 a \text{ Abs } A.$

Proof.

- 1) $f \mathcal{F}n a \iff \mathcal{F}nc(f) \wedge \mathcal{D}(f) = a.$
- 2) $f \mathcal{F}n_2 a \iff \mathcal{F}nc_2(f) \wedge \mathcal{D}(f) = a.$

Theorem 13.30. *If A is a nonempty transitive model of the Axiom of Pairing and if $x R_1 y$ and $x R_2 y$ are each absolute with respect to A then $f \text{ Isom}_{R_1, R_2}(a, b) \text{ Abs } A.$*

Proof. $f \text{ Isom}_{R_1, R_2}(a, b) \iff f \mathcal{F}n_2 a \wedge \mathcal{W}(f) = b$
 $\wedge (\forall x) (\forall y) [x \in a \wedge y \in a \wedge \langle x, y \rangle \in R_1 \rightarrow \langle f' x, f' y \rangle \in R_2].$

Theorem 13.31. *If A is nonempty and transitive then A is a standard transitive model of the Axiom of Unions iff*

$$(\forall a \in A) [\cup(a) \in A].$$

Proof. $\text{Ax. 3} \iff (\forall a) (\exists x) [x = \cup(a)].$

Since A is nonempty and transitive we have

$$\text{STM}(A, \text{Ax. 3})$$

if and only if

$$(\forall a \in A) (\exists x \in A) [x = \cup(a)]^A.$$

Since $x = \cup(a)$ is absolute w.r.t. A , this holds if and only if

$$(\forall a \in A) (\exists x \in A) [x = \cup(a)].$$

But this is true if and only if

$$(\forall a \in A) [\cup(a) \in A].$$

Remark. Axiom 1 is absolute w.r.t. any transitive class. As a consequence every transitive class is a model of this axiom. Furthermore we note that if $b \in A$ then the extent of b as an individual in the universe A is $b \cap A$, that is, the collection of objects in A that play the role of elements of b is precisely the collection of objects in A that *are* elements of b . If A is transitive then $b \in A \rightarrow b \subset A$ and $b \cap A = b$, that is b as an individual in the universe A has the same extent it has as an individual in V . Consequently if A is transitive and a and b are in A then $\{a, b\} \cap A = \{a, b\}$ and $\cup(a) \cap A = \cup(a)$. Therefore in order for A to be a model of the Axiom of Pairing and the Axiom of Unions we must have

$$\{a, b\} \in A \quad \text{for } a, b \in A,$$

$$\cup(a) \in A \quad \text{for } a \in A.$$

The Power Set Axiom however presents a different situation. If A is transitive but not super transitive then $a \in A$ does not imply that all subsets of a are in A thus

$$\mathcal{P}(a) \cap A \subseteq \mathcal{P}(a)$$

and equality need not hold. That is, even if A is to be a transitive model of the Power Set Axiom, the object in A that plays the role of the power set of a need not be $\mathcal{P}(a)$.

Theorem 13.32. *If A is nonempty and transitive then A is a standard transitive model of the Axiom of Powers iff*

$$(\forall a \in A) [\mathcal{P}(a) \cap A \in A].$$

Proof. $\text{Ax. 4} \leftrightarrow (\forall a) (\exists x) (\forall y) [y \in x \leftrightarrow y \subseteq a]$. Since A is nonempty and transitive we have

$$\text{STM}(A, \text{Ax. 4})$$

if and only if

$$(\forall a \in A) (\exists x \in A) (\forall y \in A) [y \in x \leftrightarrow y \subseteq a]^A.$$

But since $y \in x \leftrightarrow y \subseteq a$ is absolute w.r.t. A , this holds if and only if

$$(\forall a \in A) (\exists x \in A) (\forall y \in A) [y \in x \leftrightarrow y \subseteq a].$$

Since A is transitive it follows that if $x \in A$ then

$$(\forall y \in A) [y \in x \leftrightarrow y \subseteq a]$$

holds if and only if

$$(\forall y) [y \in x \leftrightarrow y \subseteq a \wedge y \in A]$$

i.e., if and only if

$$(\forall y) [y \in x \leftrightarrow y \in \mathcal{P}(a) \cap A].$$

Thus A is a transitive model of Axiom 4 if and only if

$$(\forall a \in A) (\exists x \in A) (\forall y) [y \in x \leftrightarrow y \in \mathcal{P}(a) \cap A]$$

that is, if and only if

$$(\forall a \in A) [\mathcal{P}(a) \cap A \in A].$$

Remark. Axiom 5 is of course not *an* axiom but an axiom schema. For each wff $\varphi(u, v)$ we have an instance of Axiom 5 that we will denote by Axiom 5_φ .

Theorem 13.33. *If A is nonempty and transitive then A is a standard transitive model of Axiom 5_φ , iff $\forall a \in A$*

$$(\forall u \in A) (\forall v \in A) (\forall w \in A) [\varphi^A(u, v) \wedge \varphi^A(u, w) \rightarrow v = w]$$

implies that

$$\{x \in A \mid (\exists y \in a) \varphi^A(y, x)\} \in A.$$

Proof. Since A is nonempty and transitive A is a standard transitive model of Axiom 5_φ iff $\forall a \in A$

$$(\forall u \in A) (\forall v \in A) (\forall w \in A) [\varphi^A(u, v) \wedge \varphi^A(u, w) \rightarrow v = w]$$

implies that

$$(\exists b \in A) (\forall x \in A) [x \in b \leftrightarrow (\exists y \in A) [\varphi^A(y, x) \wedge y \in a]].$$

But since A is transitive this is the case iff

$$(\exists b \in A) (\forall x) [x \in b \leftrightarrow (\exists y \in A) [\varphi^A(y, x) \wedge y \in a] \wedge x \in A]$$

i.e.

$$(\exists b \in A) [b = \{x \in A \mid (\exists y \in A) [\varphi^A(y, x) \wedge y \in a]\}]$$

Hence

$$\{x \in A \mid (\exists y \in A) [\varphi^A(y, x) \wedge y \in a]\} \in A.$$

Theorem 13.34. $STM(A, Ax. 5_\varphi) \rightarrow (\forall a \in A) [\{y \in a \mid \varphi^A(y)\} \in A].$

Proof. If A is a transitive model of Axiom 5_φ then A is a nonempty transitive class that satisfies all of the nonlogical axioms required to

prove the instance of Zermelo's Schema of Separation:

$$(\forall a)(\exists x)[x = \{y \in a \mid \varphi(y)\}].$$

(See Theorem 5.11.)

Theorem 13.35. $A \neq 0 \wedge \text{Tr}(A) \rightarrow \text{STM}(A, \text{Ax. 6})$.

Proof. $\text{Ax. 6} \leftrightarrow (\forall a)[a \neq 0 \rightarrow (\exists x)[x \in a \wedge x \cap a = 0]]$. In particular, from Axiom 6.

$$(\forall a \in A)[a \neq 0 \rightarrow (\exists x)[x \in a \wedge x \cap a = 0]].$$

But since A is transitive $x \in a \wedge a \in A \rightarrow x \in A$. Therefore

$$(\forall a \in A)[a \neq 0 \rightarrow (\exists x \in A)[x \in a \wedge x \cap a = 0]].$$

Furthermore $x \in a \wedge x \cap a = 0$ is absolute with respect to A . Therefore if $a \in A$ then

$$(\exists x \in A)[x \in a \wedge x \cap a = 0] \leftrightarrow (\exists x \in A)[x \in a \wedge x \cap a = 0]^A.$$

Since $a \neq 0$ is absolute with respect to A it follows that

$$(\forall a \in A)[a \neq 0 \rightarrow (\exists x)[x \in a \wedge x \cap a = 0]]^A$$

i.e., A is a model of Axiom 6.

Theorem 13.36. $A \neq 0 \wedge \text{Tr}(A) \rightarrow 0 \in A$.

Proof. By the Axiom of Regularity

$$\begin{aligned} A \neq 0 \wedge \text{Tr}(A) &\rightarrow (\exists x)[x \in A \wedge x \cap A = 0] \\ &\rightarrow (\exists x)[x \in A \wedge x = 0] \\ &\rightarrow 0 \in A. \end{aligned}$$

Theorem 13.37. $\text{STM}(A, \text{Ax. 2}) \wedge \text{STM}(A, \text{Ax. 3}) \rightarrow \omega \subseteq A$.

Proof (by induction). From Theorem 13.36, $0 \in A$. If $k \in A$ then since A is a model of Ax. 2, $\{k\} \in A$ and hence $\{k, \{k\}\} \in A$. Since A is also a model of Ax. 3 $\cup \{k, \{k\}\} \in A$, that is $k + 1 \in A$.

Theorem 13.38. *If A is a standard transitive model of the Axiom of Pairing and the Axiom of Unions then A is a standard transitive model of the Axiom of Infinity iff $\omega \in A$.*

Proof. Since A is nonempty and transitive A is a standard transitive model of the Axiom of Infinity if and only if

$$(\exists x \in A)[x = \omega]^A.$$

But Theorems 13.37 and 13.21 establish that $x = \omega$ is absolute w.r.t. A . Therefore A is a standard transitive model of the Axiom of Infinity if

and only if

$$(\exists x \in A) [x = \omega]$$

i.e., if and only if

$$\omega \in A.$$

Remark. We next prove the relative consistency of the Axiom of Regularity and the other axioms of ZF. In addition to being of interest in itself the proof gives an excellent illustration of Gödel's method of proving the relative consistency of ZF with the AC and the GCH. Our procedure is to prove without using the Axiom of Regularity that the class of well founded sets is a model of ZF (see Definition 9.11).

Theorem 13.39. *If $\mathcal{M} = \{x \mid \text{Wf}(x)\}$ then \mathcal{M} is a standard transitive model of ZF.*

Proof. Since $0 \in R'1$ we have $0 \in \mathcal{M}$ i.e. $\mathcal{M} \neq \emptyset$. If $x \in \mathcal{M}$ then $(\exists \alpha) [x \in R'\alpha]$. Since $R'\alpha$ is transitive $x \in R'\alpha \rightarrow x \subseteq R'\alpha$ and hence $x \subseteq \mathcal{M}$. Thus \mathcal{M} is transitive. Since \mathcal{M} is nonempty and transitive \mathcal{M} is a model of Axiom 1.

Recall that a set is well founded if each of its elements is well founded. Consequently if $a \in \mathcal{M} \wedge b \in \mathcal{M}$ then $\{a, b\} \in \mathcal{M}$. Thus \mathcal{M} is a model of Axiom 2.

If $a \in \mathcal{M}$ then $y \in \cup(a) \rightarrow (\exists x) [y \in x \wedge x \in a]$. Since \mathcal{M} is transitive it follows that $\cup(a) \subseteq \mathcal{M}$, hence $\cup(a) \in \mathcal{M}$. Thus \mathcal{M} is a model of Axiom 3.

If $a \in \mathcal{M} \wedge x \subseteq a$ then since \mathcal{M} is transitive $x \subseteq \mathcal{M}$ and hence $x \in \mathcal{M}$. Thus $\mathcal{P}(a) \subseteq \mathcal{M}$ and hence $\mathcal{P}(a) \in \mathcal{M}$. Thus \mathcal{M} is a model of Axiom 4.

If $(\forall u \in \mathcal{M}) (\forall v \in \mathcal{M}) (\forall w \in \mathcal{M}) [\varphi^{\mathcal{M}}(u, v) \wedge \varphi^{\mathcal{M}}(u, w) \rightarrow v = w]$ and if $a \in \mathcal{M}$ then

$$\{y \in \mathcal{M} \mid (\exists z \in a) \varphi^{\mathcal{M}}(z, y)\} \subseteq \mathcal{M}$$

and hence $\{y \in \mathcal{M} \mid (\exists z \in a) \varphi^{\mathcal{M}}(z, y)\} \in \mathcal{M}$. Thus \mathcal{M} is a model of Axiom 5.

If $a \in \mathcal{M}$ then since \mathcal{M} is transitive $a \subseteq \mathcal{M}$. Thus each element of a has a rank. If $a \neq \emptyset$ then a contains an element of smallest rank i.e.

$$\begin{aligned} a \neq \emptyset &\rightarrow (\exists x \in a) (\forall y \in a) [\text{rank } x \leq \text{rank } y] \\ &\rightarrow (\exists x \in a) (x \cap a = \emptyset). \end{aligned}$$

Again since \mathcal{M} is transitive $x \in a \rightarrow x \in \mathcal{M}$ i.e.

$$(\forall a \in \mathcal{M}) [a \neq \emptyset \rightarrow (\exists x \in \mathcal{M}) [x \in a \wedge x \cap a = \emptyset]].$$

Thus, since $[x \in a \wedge x \cap a = \emptyset] \text{ Abs } \mathcal{M}$, \mathcal{M} is a standard transitive model of Axiom 6.

Since \mathcal{M} is a standard transitive model of Axioms 2 and 3 it follows that $\omega \subseteq \mathcal{M}$ (Theorem 13.37). From Theorem 9.12 it then follows that $\omega \in \mathcal{M}$. Thus \mathcal{M} is a model of Axiom 7.

***Theorem 13.40.** *If α is inaccessible then $R'\alpha$ is a standard transitive model of ZF.*

Proof. Since $R'\alpha$ is nonempty and transitive it is a model of Axioms 1 and 6. If $a \in R'\alpha \wedge b \in R'\alpha$ then

$$\text{rank}(\{a, b\}) = \max(\text{rank}(a), \text{rank}(b)) + 1 < \alpha.$$

Hence $\{a, b\} \in R'\alpha$ and $R'\alpha$ is a model of Axiom 2.

If $a \in R'\alpha$ then

$$\text{rank}(\cup(a)) = \mu_\gamma((\forall x \in \cup(a)) [\text{rank}(x) < \gamma]) \leq \text{rank}(a).$$

Therefore $\cup(a) \in R'\alpha$ and $R'\alpha$ is a model of Axiom 3.

If $a \in R'\alpha$ then $\mathcal{P}(a) \subseteq R'\alpha$ and hence $\mathcal{P}(a) \cap R'\alpha = \mathcal{P}(a)$. Furthermore since $x \subseteq a \rightarrow \text{rank}(x) \leq \text{rank}(a)$ and since $a \in \mathcal{P}(a)$

$$\text{rank}(\mathcal{P}(a)) = \text{rank}(a) + 1 < \alpha.$$

Thus $\mathcal{P}(a) \in R'\alpha$ and hence $R'\alpha$ is a model of Axiom 4.

From the AC it follows by induction that

$$(\forall \beta < \alpha) [\overline{\overline{R'\beta}} < \alpha].$$

Indeed $\overline{\overline{R'0}} = 0 < \alpha$. If $\overline{\overline{R'\beta}} < \alpha$ for $\beta < \alpha$ then since α is inaccessible

$$\overline{\overline{R'(\beta + 1)}} = \overline{\overline{\mathcal{P}(R'\beta)}} < \alpha.$$

If $\beta \in K_{II}$ and if $\gamma < \beta \rightarrow \overline{\overline{R'\gamma}} < \alpha$ then

$$\overline{\overline{R'\beta}} = \overline{\overline{\bigcup_{\gamma < \beta} R'\gamma}} = \bigcup_{\gamma < \beta} \overline{\overline{R'\gamma}} \leq \alpha.$$

If $\overline{\overline{R'\beta}} = \alpha$ and if $f(\gamma) \triangleq \overline{\overline{R'\gamma}}$ then $f: \beta \rightarrow \alpha$, f is monotone increasing and $\cup(f''\beta) = \alpha$. Hence α is cofinal with β . This is a contradiction from which we conclude that

$$\overline{\overline{R'\beta}} < \alpha.$$

If for some $a \in R'\alpha$ and some wff φ we have

$$(\forall u \in R'\alpha) (\forall v \in R'\alpha) (\forall w \in R'\alpha) [\varphi^{R'\alpha}(u, v) \wedge \varphi^{R'\alpha}(u, w) \rightarrow v = w]$$

and

$$\{y \in R'\alpha \mid (\exists x \in a) \varphi^{R'\alpha}(x, y)\} \notin R'\alpha$$

then since $(\exists \beta < \alpha) [a \subseteq R'\beta]$, $\bar{a} \leq \overline{\overline{R'\beta}} < \alpha$ i.e. $(\exists \gamma < \alpha) [\gamma \simeq a]$. Since

$$\text{rank}(\{y \in R'\alpha \mid (\exists x \in a) \varphi^{R'\alpha}(x, y)\}) = \alpha$$

it then follows that α is cofinal with some ordinal smaller than or equal to γ . This is a contradiction and hence $R'\alpha$ is a model of Axiom 5.

Since $\omega \in R'\alpha$, $R'\alpha$ is a model of Axiom 7.

Exercises. *Prove the following*

- 1) $\text{STM}(R'\alpha, \text{Ax. 1}) \longleftrightarrow \alpha > 0.$
- 2) $\text{STM}(R'\alpha, \text{Ax. 2}) \longleftrightarrow \alpha \in K_{\text{II}}.$
- 3) $\text{STM}(R'\alpha, \text{Ax. 3}) \longleftrightarrow \alpha > 0.$
- 4) $\text{STM}(R'\alpha, \text{Ax. 4}) \longleftrightarrow \alpha \in K_{\text{II}}.$
- 5) $\text{STM}(R'\omega, \text{Ax. 5}) \wedge \neg \text{STM}(R(\omega 2), \text{Ax. 5}).$
- 6) $\text{STM}(R'\alpha, \text{Ax. 6}) \longleftrightarrow \alpha > 0.$
- 7) $\text{STM}(R'\alpha, \text{Ax. 7}) \longleftrightarrow \alpha > \omega.$

In Exercises 8–15 assume α an inaccessible cardinal.

- 8) $\mathcal{P}(a) \text{ Abs } R'\alpha.$
- 9) $a \simeq b \text{ Abs } R'\alpha.$
- 10) $\text{cf}(\beta) \text{ Abs } R'\alpha.$
- 11) $\beta \in N \text{ Abs } R'\alpha.$
- 12) $\beta \in N' \text{ Abs } R'\alpha.$
- 13) $\text{Reg } (\beta) \text{ Abs } R'\alpha.$
- 14) $\text{Inacc}_w(\beta) \text{ Abs } R'\alpha.$
- 15) $\text{Inacc}(\beta) \text{ Abs } R'\alpha.$

Remark. There are several interesting conclusions to be drawn from the foregoing exercises and Theorem 13.40. First we have that if α is inaccessible then $R'\alpha$ is a standard transitive model of ZF. The proof requires the AC. From this and Exercise 15 it follows that it is consistent with ZF to assume that there are no inaccessible cardinals. We argue in the following way. Suppose the statement “there exists an inaccessible cardinal” were provable in ZF. Let α be the smallest such cardinal. Then $R'\alpha$ is a standard transitive model of ZF. But from Exercise 15 above, if there were an inaccessible cardinal in $R'\alpha$ that cardinal would be inaccessible in V and smaller than the smallest inaccessible cardinal. This is a contradiction. Therefore it is not possible to prove in ZF that inaccessible cardinals exist. It may, however, be possible to prove in ZF that there does not exist an inaccessible cardinal. No proof is known at the time of this writing.

From Exercises 1–7, $R'\omega$ is a standard transitive model of Axioms 1–6 but not of Axiom 7. Thus Axiom 7 is independent of Axioms 1–6. Also $R(\omega 2)$ is a standard transitive model of all axioms except Axiom 5.

14 The Fundamental Operations

Gödel proved the relative consistency of the AC and the GCH by showing that a certain class L is a model of $ZF + AC + GCH$. This class L he defined initially as the union of a sequence of sets A_α , $\alpha \in \text{On}$ which were so defined that $x \in A_{\alpha+1}$ iff there exists a wff $\varphi(x_0, x_1, \dots, x_n)$ having no free variables other than x_0, x_1, \dots, x_n and there exist $a_1, \dots, a_n \in A_\alpha$ such that $x = \{y \mid A_\alpha \models \varphi(y, a_1, \dots, a_n)\}$.

The foregoing condition describes a sense in which $A_{\alpha+1}$ is the collection of sets that are definable from A_α . However, to properly define “definable” we must avoid quantification on wffs. This we will do in later chapters¹.

Later Gödel discovered that his class of constructible sets could be defined as the range of a certain function F defined on On by transfinite recursion from eight basic operations. In Section 15, we will follow Gödel’s second development. In anticipation of that development, we now establish certain conditions involving Gödel’s eight fundamental operations to be defined below that are sufficient for a class M to be a standard transitive model of ZF.

Definition 14.1.

- 1) $\text{Cnv}_2(A) = \{\langle x, y, z \rangle \mid \langle z, x, y \rangle \in A\}$.
- 2) $\text{Cnv}_3(A) = \{\langle x, y, z \rangle \mid \langle x, z, y \rangle \in A\}$.

Remark. $\text{Cnv}_2(A)$ and $\text{Cnv}_3(A)$ are read “the second converse of A ” and “the third converse of A ” respectively.

Definition 14.2 (The Fundamental Operations).

$$\mathcal{F}_1(a, b) \triangleq \{a, b\}.$$

$$\mathcal{F}_2(a, b) \triangleq a \cap b.$$

$$\mathcal{F}_3(a, b) \triangleq a - b.$$

$$\mathcal{F}_4(a, b) \triangleq a \cap b.$$

¹ For a more detailed discussion of definability see Takeuti and Zaring: *Axiomatic Set Theory*. Springer-Verlag.

$$\mathcal{F}_5(a, b) \triangleq a \cap \mathcal{D}(b).$$

$$\mathcal{F}_6(a, b) \triangleq a \cap b^{-1}.$$

$$\mathcal{F}_7(a, b) \triangleq a \cap \text{Cnv}_2(b).$$

$$\mathcal{F}_8(a, b) \triangleq a \cap \text{Cnv}_3(b).$$

Theorem 14.3. *If \mathcal{M} is a standard transitive model of ZF then \mathcal{M} is closed under the eight fundamental operations.*

Proof. From Theorem 13.19

$$a \in \mathcal{M} \wedge b \in \mathcal{M} \wedge \text{STM}(\mathcal{M}, \text{Ax. 2}) \rightarrow \mathcal{F}_1(a, b) \in \mathcal{M}.$$

Since \mathcal{M} is a model of Axiom 2, $x = \langle a, b \rangle$ and $x = \langle a, b, c \rangle$ are each absolute with respect to \mathcal{M} . Since \mathcal{M} is also a model of Axiom 5 it follows from Theorem 13.34 and properties of absoluteness that for $a \in \mathcal{M} \wedge b \in \mathcal{M}$

$$\begin{aligned} \mathcal{F}_2(a, b) &= \{x \in a \mid (\exists c) (\exists d) [x = \langle c, d \rangle \wedge c \in d]\} \\ &= \{x \in a \mid (\exists c \in \mathcal{M}) (\exists d \in \mathcal{M}) [x = \langle c, d \rangle \wedge c \in d]^{\mathcal{M}}\} \in \mathcal{M}. \end{aligned}$$

$$\mathcal{F}_3(a, b) = \{x \in a \mid x \notin b\} = \{x \in a \mid [x \notin b]^{\mathcal{M}}\} \in \mathcal{M}.$$

$$\begin{aligned} \mathcal{F}_4(a, b) &= \{x \in a \mid (\exists c) (\exists d) [x = \langle c, d \rangle \wedge c \in b]\} \\ &= \{x \in a \mid (\exists c \in \mathcal{M}) (\exists d \in \mathcal{M}) [x = \langle c, d \rangle \wedge c \in b]^{\mathcal{M}}\} \in \mathcal{M}. \end{aligned}$$

$$\mathcal{F}_5(a, b) = \{x \in a \mid (\exists y) [\langle x, y \rangle \in b]\} = \{x \in a \mid (\exists y \in \mathcal{M}) [\langle x, y \rangle \in b]^{\mathcal{M}}\} \in \mathcal{M}.$$

$$\begin{aligned} \mathcal{F}_6(a, b) &= \{x \in a \mid (\exists c) (\exists d) [x = \langle c, d \rangle \wedge \langle d, c \rangle \in b]\} \\ &= \{x \in a \mid (\exists c \in \mathcal{M}) (\exists d \in \mathcal{M}) [x = \langle c, d \rangle \wedge \langle d, c \rangle \in b]\} \in \mathcal{M}. \end{aligned}$$

$$\begin{aligned} \mathcal{F}_7(a, b) &= \{x \in a \mid (\exists c) (\exists d) (\exists e) [x = \langle c, d, e \rangle \wedge \langle e, c, d \rangle \in b]\} \\ &= \{x \in a \mid (\exists c \in \mathcal{M}) (\exists d \in \mathcal{M}) (\exists e \in \mathcal{M}) \\ &\quad [x = \langle c, d, e \rangle \wedge \langle e, c, d \rangle \in b]^{\mathcal{M}}\} \in \mathcal{M}. \end{aligned}$$

$$\begin{aligned} \mathcal{F}_8(a, b) &= \{x \in a \mid (\exists c) (\exists d) (\exists e) [x = \langle c, d, e \rangle \wedge \langle c, e, d \rangle \in b]\} \\ &= \{x \in a \mid (\exists c \in \mathcal{M}) (\exists d \in \mathcal{M}) (\exists e \in \mathcal{M}) \\ &\quad [x = \langle c, d, e \rangle \wedge \langle c, e, d \rangle \in b]^{\mathcal{M}}\} \in \mathcal{M}. \end{aligned}$$

Remark. An examination of the foregoing proof reveals that the full strength of the hypothesis that \mathcal{M} is a standard transitive model of ZF was not used. All that is required is that \mathcal{M} be a standard transitive model of Axiom 2 and of seven instances of Axiom 5. In view of this it is not reasonable to expect the condition that \mathcal{M} be closed under the eight fundamental operations to be also sufficient for a nonempty transitive class \mathcal{M} to be a standard transitive model of ZF. Surprisingly in addition to closure under the eight fundamental operations we need only the added condition that every subset of \mathcal{M} have an extension in \mathcal{M} .

Definition 14.4. \mathcal{M} is almost universal iff

$$(\forall x) [x \subseteq \mathcal{M} \rightarrow (\exists a \in \mathcal{M}) [x \subseteq a]].$$

Remark. Note that if \mathcal{M} is almost universal then \mathcal{M} is not empty.

To prove that if \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations then \mathcal{M} is a standard transitive model of ZF we need a few preliminary results.

Theorem 14.5. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations then $\forall a, b \in \mathcal{M}$*

- | | | |
|--|----------------------------------|--------------------------------|
| 1) $\{a, b\} \in \mathcal{M},$ | 3) $a \times b \in \mathcal{M},$ | 5) $a \cap b \in \mathcal{M},$ |
| 2) $\langle a, b \rangle \in \mathcal{M},$ | 4) $a - b \in \mathcal{M},$ | 6) $a \cup b \in \mathcal{M}.$ |

Proof.

1) $\{a, b\} = \mathcal{F}_1(a, b) \in \mathcal{M}.$

2) $\langle a, b \rangle = \{\{a\}, \{a, b\}\} \in \mathcal{M}.$

3) Since \mathcal{M} is transitive $a \subseteq \mathcal{M} \wedge b \subseteq \mathcal{M}$. Therefore $a \times b \subseteq \mathcal{M} \wedge b \times a \subseteq \mathcal{M}$. But \mathcal{M} is also almost universal. Therefore

$$(\exists c, d \in \mathcal{M}) [a \times b \subseteq c \wedge b \times a \subseteq d].$$

Then

$$\begin{aligned} a \times b &= [c \cap (a \times V)] \cap [d \cap (b \times V)]^{-1} \\ &= \mathcal{F}_6(\mathcal{F}_4(c, a), \mathcal{F}_4(d, b)) \in \mathcal{M}. \end{aligned}$$

4) $a - b = \mathcal{F}_3(a, b) \in \mathcal{M}.$

5) $a \cap b = a - (a - b) \in \mathcal{M}.$

6) Since \mathcal{M} is transitive $a \cup b \subseteq \mathcal{M}$. Also \mathcal{M} is almost universal. Therefore

$$(\exists c \in \mathcal{M}) [a \cup b \subseteq c].$$

Then $a \cup b = c - [(c - a) - b] \in \mathcal{M}.$

Theorem 14.6. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations then*

$$(\forall a_1, \dots, a_n \in \mathcal{M}) [a_1 \times a_2 \times \dots \times a_n \in \mathcal{M}].$$

Proof. Obvious from Theorem 14.5.3 by induction.

Theorem 14.7. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations then $\forall a \in \mathcal{M}$.*

- | | | |
|-------------------------------------|--------------------------------------|---------------------------------------|
| 1) $a^n \in \mathcal{M}, n \geq 1,$ | 3) $\mathcal{D}(a) \in \mathcal{M},$ | 5) $\text{Cnv}_2(a) \in \mathcal{M},$ |
| 2) $a^{-1} \in \mathcal{M},$ | 4) $\mathcal{W}(a) \in \mathcal{M},$ | 6) $\text{Cnv}_3(a) \in \mathcal{M}.$ |

Proof. 1) Obvious from Theorem 14.6.

2) Since \mathcal{M} is transitive $\langle x, y \rangle \in a \wedge a \in \mathcal{M}$ implies $x, y \in \mathcal{M}$. This in turn implies that $\langle y, x \rangle \in \mathcal{M}$, i.e., $a^{-1} \subseteq \mathcal{M}$. Then $(\exists b \in \mathcal{M}) [a^{-1} \subseteq b]$ and hence

$$a^{-1} = b \cap a^{-1} = \mathcal{F}_6(b, a) \in \mathcal{M}.$$

3) $x \in \mathcal{D}(a) \rightarrow (\exists y) [\langle x, y \rangle \in a]$. Again from transitivity it follows that $\mathcal{D}(a) \subseteq \mathcal{M}$ and hence $(\exists b \in \mathcal{M}) [\mathcal{D}(a) \subseteq b]$. Therefore

$$\mathcal{D}(a) = b \cap \mathcal{D}(a) = \mathcal{F}_5(b, a) \in \mathcal{M}.$$

4) $\mathcal{W}(a) = \mathcal{D}(a^{-1}) \in \mathcal{M}$.

5) Since \mathcal{M} is transitive $\text{Cnv}_2(a) \subseteq \mathcal{M}$. Therefore

$$(\exists b \in \mathcal{M}) [\text{Cnv}_2(a) \subseteq b].$$

Then

$$\text{Cnv}_2(a) = b \cap \text{Cnv}_2(a) = \mathcal{F}_7(b, a) \in \mathcal{M}.$$

6) The proof is left to the reader.

Theorem 14.8. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations, if (i_1, i_2, i_3) is a permutation of 1, 2, 3 and if $a \in \mathcal{M}$ then*

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_{i_1}, x_{i_2}, x_{i_3} \rangle \in a\} \in \mathcal{M}.$$

Proof.

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_1, x_3, x_2 \rangle \in a\} = \text{Cnv}_3(a) \in \mathcal{M}.$$

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_3, x_1, x_2 \rangle \in a\} = \text{Cnv}_2(a) \in \mathcal{M}.$$

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_3, x_2, x_1 \rangle \in a\} = \text{Cnv}_2(\text{Cnv}_3(a)) \in \mathcal{M}.$$

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_1, x_2, x_3 \rangle \in a\} = \text{Cnv}_3(\text{Cnv}_3(a)) \in \mathcal{M}.$$

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_2, x_1, x_3 \rangle \in a\} = \text{Cnv}_3(\text{Cnv}_2(a)) \in \mathcal{M}.$$

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_2, x_3, x_1 \rangle \in a\} = \text{Cnv}_2(\text{Cnv}_2(a)) \in \mathcal{M}.$$

Theorem 14.9. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations then $\forall a, b \in \mathcal{M}$ if $a \subseteq \mathcal{M}^2$*

1) $\{\langle x, y, z \rangle \mid \langle x, y \rangle \in a \wedge z \in b\} \in \mathcal{M}$,

2) $\{\langle x, z, y \rangle \mid \langle x, y \rangle \in a \wedge z \in b\} \in \mathcal{M}$,

3) $\{\langle z, x, y \rangle \mid \langle x, y \rangle \in a \wedge z \in b\} \in \mathcal{M}$.

Proof. Obvious from Theorem 14.8 and the fact that $a \times b \in \mathcal{M}$.

Theorem 14.10. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations and if $\varphi(x_1, \dots, x_m)$ is a wff all of whose free variables are among $b_1, \dots, b_n, x_1, \dots, x_m$ then $(\forall c_1, \dots, c_m \in \mathcal{M})$ if $b_1 \in \mathcal{M} \wedge \dots \wedge b_m \in \mathcal{M}$ then*

$$a \triangleq \{ \langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge \varphi^{\mathcal{M}}(x_1, \dots, x_m) \} \in \mathcal{M} .$$

Proof. (By induction on k the number of logical symbols in $\varphi(x_1, \dots, x_m)$.)

If $k=0$ then $\varphi(x_1, \dots, x_m)$ either 1) contains none of the variables x_1, \dots, x_m or it is of the form 2) $x_i \in x_j$ or 3) $x_i \in b_j$ or 4) $b_i \in x_j$.

Case 1) If $\varphi(x_1, \dots, x_m)$ contains none of the variables x_1, \dots, x_m then $a = c_1 \times c_2 \times \dots \times c_m$ or $a = 0$ according as $\varphi^{\mathcal{M}}(x_1, \dots, x_m)$ holds or does not hold. In either case $a \in \mathcal{M}$.

Case 2) If $\varphi(x_1, \dots, x_m)$ is $x_i \in x_j$ then $i < j$ or $i = j$ or $j < i$. If $i < j$ then since $(c_i \times c_j) \cap E \in \mathcal{M}$, and $c_1 \times \dots \times c_{i-1} \in \mathcal{M}$ we have from Theorem 14.9.3 that

$$\{ \langle x_1, \dots, x_i, x_j \rangle \mid \langle x_i, x_j \rangle \in [(c_i \times c_j) \cap E] \wedge \langle x_1, \dots, x_{i-1} \rangle \in c_1 \times \dots \times c_{i-1} \} \in \mathcal{M} .$$

From this we obtain after $j - (i + 1)$ applications of 14.9.2

$$\{ \langle x_1, \dots, x_j \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_j \in c_j \wedge x_i \in x_j \} \in \mathcal{M} .$$

With $m - (j + 1)$ applications of Theorem 14.9.1 we have

$$\{ \langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge x_i \in x_j \} \in \mathcal{M} .$$

If $i = j$ then $a = 0 \in \mathcal{M}$. If $j < i$ then $[(c_i \times c_j) \cap E]^{-1} \in \mathcal{M} \wedge c_1 \times \dots \times c_{j-1} \in \mathcal{M}$. Then

$$\begin{aligned} \{ \langle x_1, \dots, x_j, x_i \rangle \mid \langle x_j, x_i \rangle \in [(c_i \times c_j) \cap E]^{-1} \\ \wedge \langle x_1, \dots, x_{j-1} \rangle \in c_1 \times \dots \times c_{j-1} \} \in \mathcal{M} . \end{aligned}$$

We then proceed as before.

Case 3) If $\varphi(x_1, \dots, x_m)$ is $x_i \in b_j$ then

$$\begin{aligned} \{ \langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge x_i \in b_j \} \\ = (c_1 \times \dots \times c_m) \cap (c_1 \times \dots \times c_{i-1} \times b_j \times c_{j+1} \times \dots \times c_m) \in \mathcal{M} . \end{aligned}$$

Case 4) If $\varphi(x_1, \dots, x_m)$ is $b_j \in x_i$ then

$$\{ x_i \in c_i \mid b_j \in x_i \} = \mathcal{W}(\{ \{ b_j \} \times c_i \} \cap E) \in \mathcal{M} .$$

Then

$$\begin{aligned} \{ \langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge b_j \in x_i \} \\ = c_1 \times \dots \times c_{i-1} \times \{ x_i \in c_i \mid b_j \in x_i \} \times c_{i+1} \times \dots \times c_m \in \mathcal{M} . \end{aligned}$$

If $k > 0$ then $\varphi(x_1, \dots, x_m)$ is of the form 1) $\neg \psi(x_1, \dots, x_m)$ or 2) $\psi(x_1, \dots, x_m) \wedge \eta(x_1, \dots, x_m)$ or 3) $(\exists x) \psi(x_1, \dots, x_m, x)$.

Case 1) If $\varphi(x_1, \dots, x_m)$ is $\neg \psi(x_1, \dots, x_m)$ then as our induction hypothesis we have

$$\{\langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge \psi^{\mathcal{M}}(x_1, \dots, x_m)\} \in \mathcal{M}.$$

Then

$$\begin{aligned} & \{\langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge \neg \psi^{\mathcal{M}}(x_1, \dots, x_m)\} \\ &= c_1 \times \dots \times c_m - \{\langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge \psi^{\mathcal{M}}(x_1, \dots, x_m) \in \mathcal{M}\}. \end{aligned}$$

Case 2) If $\varphi(x_1, \dots, x_m)$ is $\psi(x_1, \dots, x_m) \wedge \eta(x_1, \dots, x_m)$ then from the induction hypothesis

$$\begin{aligned} & \{\langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge \psi^{\mathcal{M}}(x_1, \dots, x_m) \wedge \eta^{\mathcal{M}}(x_1, \dots, x_m)\} \\ &= \{\langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge \psi^{\mathcal{M}}(x_1, \dots, x_m)\} \\ &\quad \cap \{\langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge \eta^{\mathcal{M}}(x_1, \dots, x_m)\} \in \mathcal{M}. \end{aligned}$$

Case 3) If $\varphi(x_1, \dots, x_m)$ is $(\exists x) \psi(x_1, \dots, x_m, x)$ and

$$\begin{aligned} F^c \langle x_1, \dots, x_m \rangle &= \{x \in \mathcal{M} \mid \psi^{\mathcal{M}}(x_1, \dots, x_m, x) \wedge (\forall y \in \mathcal{M}) [\psi^{\mathcal{M}}(x_1, \dots, x_m, y) \\ &\quad \rightarrow \text{rank}(x) \leq \text{rank}(y)] \text{ if } x_i \in c_i \text{ for } 1 \leq i \leq n, \\ &= 0 \text{ otherwise} \end{aligned}$$

then $F^c(c_1 \times \dots \times c_m)$ is a set and $\cup F^c(c_1 \times \dots \times c_m) \subseteq \mathcal{M}$. Since \mathcal{M} is almost universal

$$(\exists c \in \mathcal{M}) [\cup F^c(c_1 \times \dots \times c_m) \subseteq c]$$

then $(\exists x \in \mathcal{M}) \psi^{\mathcal{M}}(x_1, \dots, x_m, x) \leftrightarrow (\exists x \in c) \psi^{\mathcal{M}}(x_1, \dots, x_m, x)$. But by our induction hypothesis

$$\{\langle x_1, \dots, x_m, x \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge x \in c \wedge \psi^{\mathcal{M}}(x_1, \dots, x_m, x)\} \in \mathcal{M}.$$

Then

$$\begin{aligned} & \{\langle x_1, \dots, x_m \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge (\exists x \in \mathcal{M}) \psi^{\mathcal{M}}(x_1, \dots, x_m, x)\} \\ &= \mathcal{D} \{\langle x_1, \dots, x_m, x \rangle \mid x_1 \in c_1 \wedge \dots \wedge x_m \in c_m \wedge x \in c \wedge \psi^{\mathcal{M}}(x_1, \dots, x_m, x)\} \in \mathcal{M}. \end{aligned}$$

Theorem 14.11. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations then \mathcal{M} is a standard transitive model of ZF and $\text{On} \subseteq \mathcal{M}$.*

Proof. Since \mathcal{M} is almost universal $\mathcal{M} \neq 0$. Therefore, \mathcal{M} is a standard transitive model of the Axiom of Extensionality (Axiom 1) and the Axiom of Regularity (Axiom 6). By Theorem 14.5 $(\forall a, b \in \mathcal{M}) [\{a, b\} \in \mathcal{M}]$. Therefore \mathcal{M} is a model of the Axiom of Pairing (Axiom 2).

Since \mathcal{M} is transitive, $(\forall a \in \mathcal{M}) [\cup(a) \subseteq \mathcal{M}]$. Hence

$$(\exists b \in \mathcal{M}) [\cup(a) \subseteq b].$$

Since by Theorem 14.5, $b \times a \in \mathcal{M}$ and since \mathcal{M} is closed under the eight fundamental operations

$$(b \times a) \cap E = \mathcal{F}_2(b \times a, a) \in \mathcal{M} .$$

Then from Theorem 14.7

$$\cup(a) = \mathcal{D}((b \times a) \cap E) \in \mathcal{M} .$$

Thus \mathcal{M} is a model of the Axiom of Unions (Axiom 3).

Since $(\forall a \in \mathcal{M}) [\mathcal{P}(a) \cap \mathcal{M} \subset \mathcal{M}]$, $(\exists b \in \mathcal{M}) [\mathcal{P}(a) \cap \mathcal{M} \subseteq b]$. Then by Theorem 14.10

$$\mathcal{P}(a) \cap \mathcal{M} = \{x \mid x \in b \wedge [x \subseteq a]^{\mathcal{M}}\} \in \mathcal{M} .$$

Therefore \mathcal{M} is a model of the Axiom of Powers (Axiom 4).

If $(\forall u, v, w \in \mathcal{M}) [\varphi^{\mathcal{M}}(u, v) \wedge \varphi^{\mathcal{M}}(u, w) \rightarrow v = w]$ then $\forall a \in \mathcal{M}$

$$F \triangleq \{\langle x, y \rangle \in \mathcal{M}^2 \mid x \in a \wedge \varphi^{\mathcal{M}}(x, y)\}$$

is a function. Since $\mathcal{D}(F) \subseteq a$ both $\mathcal{D}(F) \wedge \mathcal{W}(F)$ are sets. Therefore since $F''a \subseteq \mathcal{M}$, $(\exists b \in \mathcal{M}) [F''a \subseteq b]$. Then

$$\begin{aligned} \{y \in \mathcal{M} \mid (\exists x \in a) \varphi^{\mathcal{M}}(x, y)\} &= \mathcal{W}(\{\langle x, y \rangle \in \mathcal{M}^2 \mid x \in a \wedge \varphi^{\mathcal{M}}(x, y)\}) \\ &= \mathcal{W}(\{\langle x, y \rangle \mid x \in a \wedge y \in b \wedge \varphi^{\mathcal{M}}(x, y)\}) \in \mathcal{M} . \end{aligned}$$

Thus \mathcal{M} is a model of the Axiom Schema of Replacement (Axiom 5).

Since \mathcal{M} is a standard transitive model of the Axiom of Pairing and the Axiom of Unions it follows from Theorem 13.37 that $\omega \subseteq \mathcal{M}$. Since \mathcal{M} is almost universal

$$(\exists a \in \mathcal{M}) [\omega \subseteq a] .$$

Since $[x \in \omega] \text{ Abs } \mathcal{M}$ and \mathcal{M} is a model of the Axiom Schema of Replacement it follows from Theorem 13.34 that

$$\omega = \{x \in a \mid x \in \omega\} = \{x \in a \mid [x \in \omega]^{\mathcal{M}}\} \in \mathcal{M} .$$

Therefore \mathcal{M} is a model of the Axiom of Infinity (Axiom 7).

Thus \mathcal{M} is a standard transitive model of ZF. That $\text{On} \subseteq \mathcal{M}$ we prove by induction.

If $\alpha \subseteq \mathcal{M}$ then since \mathcal{M} is almost universal

$$(\exists a \in \mathcal{M}) [\alpha \subseteq a] .$$

Then

$$\{x \in a \mid \text{Ord}(x)\} = \{x \in a \mid [\text{Ord}(x)]^{\mathcal{M}}\} \in \mathcal{M} .$$

Therefore

$$\cup \{x \in a \mid \text{Ord}(x)\} \in \mathcal{M} .$$

But the union of a collection of ordinals is an ordinal i.e.

$$(\exists \beta \in \mathcal{M}) (\forall \gamma \in a) [\gamma \leq \beta] .$$

Then $(\forall \gamma \in a) [\gamma \in \beta + 1]$. Since $\alpha \subseteq a$ it follows that $\alpha \subseteq \beta + 1$. Since $\beta \in \mathcal{M}$ it follows that $\beta + 1 = \beta \cup \{\beta\} \in \mathcal{M}$. Hence $\alpha \in \mathcal{M}$.

Remark. With the proof of Theorem 14.11 we have achieved our major objective for this section. There is however an interesting theory of classes in which certain results of this section are generalized. We state the main results of this theory leaving the proofs as exercises.

Definition 14.12. A is \mathcal{M} -constructible $\leftrightarrow A \subseteq \mathcal{M} \wedge (\forall x \in \mathcal{M}) [x \cap A \in \mathcal{M}]$.

Theorem 14.13. *If \mathcal{M} is transitive, and almost universal, and if a is \mathcal{M} -constructible then $a \in \mathcal{M}$.*

Remark. Theorem 14.13 tells us that if \mathcal{M} is transitive and almost universal then every \mathcal{M} -constructible set is an element in \mathcal{M} . As an application of this theorem it can be proved that if in addition \mathcal{M} is closed under the eight fundamental operation then $a \in \mathcal{M}$ implies $\cup(a)$ is \mathcal{M} -constructible. It then follows that $\cup(a) \in \mathcal{M}$ and hence \mathcal{M} satisfies the Axiom of Unions.

Theorem 14.14. *If \mathcal{M} is transitive then \mathcal{M} is \mathcal{M} -constructible.*

Theorem 14.15. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations and if A and B are \mathcal{M} -constructible then*

- 1) $E \cap \mathcal{M}$ is \mathcal{M} -constructible.
- 2) $A - B$ is \mathcal{M} -constructible.
- 3) $A \cap B$ is \mathcal{M} -constructible.
- 4) $A \cup B$ is \mathcal{M} -constructible.

Definition 14.16.

$$Q_4 = \{\langle x, \langle x, y \rangle \rangle \mid x \in V \wedge y \in V\},$$

$$Q_5 = \{\langle \langle x, y \rangle, x \rangle \mid x \in V \wedge y \in V\},$$

$$Q_6 = \{\langle \langle x, y \rangle, \langle y, x \rangle \rangle \mid x \in V \wedge y \in V\},$$

$$Q_7 = \{\langle \langle x, y, z \rangle, \langle y, z, x \rangle \rangle \mid x \in V \wedge y \in V \wedge z \in V\},$$

$$Q_8 = \{\langle \langle x, y, z \rangle, \langle x, z, y \rangle \rangle \mid x \in V \wedge y \in V \wedge z \in V\}.$$

Theorem 14.17.

- 1) $Q_4^*A = A \times V$ and $\mathcal{F}_4(a, b) = a \cap Q_4^*b$.
- 2) $Q_5^*A = \mathcal{D}(A) \wedge \mathcal{U}_n(Q_5)$ and $\mathcal{F}_5(a, b) = a \cap Q_5^*b$.
- 3) $Q_6^*A = A^{-1} \wedge \mathcal{U}_n(Q_6)$ and $\mathcal{F}_6(a, b) = a \cap Q_6^*b$.
- 4) $Q_7^*A = \text{Cnv}_2(A) \wedge \mathcal{U}_n(Q_7)$ and $\mathcal{F}_7(a, b) = a \cap Q_7^*b$.
- 5) $Q_8^*A = \text{Cnv}_3(A) \wedge \mathcal{U}_n(Q_8)$ and $\mathcal{F}_8(a, b) = a \cap Q_8^*b$.

Theorem 14.18. If \mathcal{M} is transitive and closed under the eight fundamental operation and if $a \in \mathcal{M}$ then

$$Q_n^{\cdot} a \in \mathcal{M}, \quad n = 5, 6, 7, 8.$$

Lemma. If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations, if A is \mathcal{M} -constructible, if $a \in \mathcal{M}$ and if G_n is a function on a defined by

$$G_n^{\cdot} b = \{c \in A \mid \langle c, b \rangle \in Q_n \wedge (\forall x \in A) [\langle x, b \rangle \in Q_n \rightarrow \text{rank}(c) \leq \text{rank}(x)]\}$$

then

- 1) $(\exists d \in \mathcal{M}) [\cup(G_n^{\cdot} a) \subseteq d]$,
- 2) $(\exists d_0 \in \mathcal{M}) [a \cap Q_n^{\cdot} d_0 = a \cap Q_n^{\cdot} A]$.

Theorem 14.19. If \mathcal{M} is transitive, almost universal and closed under the eight fundamental operations and if A is \mathcal{M} -constructible then

- 1) $\mathcal{M} \cap Q_4^{\cdot} A$ is \mathcal{M} -constructible.
- 2) $Q_n^{\cdot} A$ is \mathcal{M} -constructible $n = 5, 6, 7, 8$.

Corollary 14.20. If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations and if A is \mathcal{M} -constructible then

- 1) $\mathcal{M} \cap (A \times V)$ is \mathcal{M} -constructible.
- 2) $\mathcal{D}(A)$ is \mathcal{M} -constructible.
- 3) A^{-1} is \mathcal{M} -constructible.
- 4) $\text{Cnv}_2(A)$ is \mathcal{M} -constructible.
- 5) $\text{Cnv}_3(A)$ is \mathcal{M} -constructible.

Theorem 14.21. If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations and if A and B are each \mathcal{M} -constructible then

- 1) $A \times B$ is \mathcal{M} -constructible.
- 2) A^n is \mathcal{M} -constructible.
- 3) $\mathcal{W}(A)$ is \mathcal{M} -constructible.
- 4) $A \Gamma B$ is \mathcal{M} -constructible.
- 5) $A^{\cdot} B$ is \mathcal{M} -constructible.

Theorem 14.22. If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations and if A and B are \mathcal{M} -constructible with $A \subseteq \mathcal{M}^2$ then

- 1) $\{x \in \mathcal{M} \mid a \in x\}$ is \mathcal{M} -constructible.
- 2) $\{\langle x, y, z \rangle \mid \langle x, y \rangle \in A \wedge z \in B\}$ is \mathcal{M} -constructible.
- 3) $\{\langle x, z, y \rangle \mid \langle x, y \rangle \in A \wedge z \in B\}$ is \mathcal{M} -constructible.
- 4) $\{\langle z, x, y \rangle \mid \langle x, y \rangle \in A \wedge z \in B\}$ is \mathcal{M} -constructible.

Theorem 14.23. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations and if $\varphi(x_1, \dots, x_m)$ is a wff all of whose free variables are among $b_1, \dots, b_n, x_1, \dots, x_m$ then*

$$b_1 \in \mathcal{M} \wedge \dots \wedge b_n \in \mathcal{M}$$

implies

$$A \triangleq \{ \langle x_1, \dots, x_m \rangle \in \mathcal{M}^m \mid \varphi^{\mathcal{M}}(x_1, \dots, x_m) \}$$

is \mathcal{M} -constructible.

Theorem 14.24. *If \mathcal{M} is transitive, almost universal, and closed under the eight fundamental operations then $\forall a \in \mathcal{M}$*

- 1) $\cup(a)$ is \mathcal{M} -constructible.
- 2) $\mathcal{P}(a) \cap \mathcal{M}$ is \mathcal{M} -constructible.
- 3) *If all of the free variables of $\varphi(u, v)$ are among u, v, b_1, \dots, b_n , if $b_1 \in \mathcal{M} \wedge \dots \wedge b_n \in \mathcal{M}$ and*

$$(\forall u, v, w \in \mathcal{M}) [\varphi^{\mathcal{M}}(u, v) \wedge \varphi^{\mathcal{M}}(u, w) \rightarrow v = w]$$

then $\{y \in \mathcal{M} \mid (\exists x \in a) \varphi^{\mathcal{M}}(x, y)\}$ is \mathcal{M} -constructible.

Remark. From Theorems 14.24 and 14.13 we have another proof of Theorem 14.11.

15 The Gödel Model

In Section 7 we defined a relation R_0 on On^2 . We proved that R_0 well orders On^2 and, with respect to R_0 , initial segments of On^2 are sets. Consequently there is an order isomorphism J_0 such that

$$J_0 \text{ Isom}_{R_0, E}(\text{On}^2, \text{On}).$$

This isomorphism we illustrate with the following diagram:

$\alpha \backslash \beta$	0	1	2	...	ω
0	0	1	4	...	ω
1	2	3	5	...	$\omega + 1$
2	6	7	8	...	$\omega + 2$
.
.
.
ω	$\omega 2$	$\omega 2 + 1$	$\omega 2 + 2$...	$\omega 3$

Here the element in the α^{th} row and β^{th} column is $J_0' \langle \alpha, \beta \rangle$. From the diagram it is apparent that the entry in the α^{th} row and β^{th} column i.e. $J_0' \langle \alpha, \beta \rangle$ is at least as large as the maximum of α and β . It is also easily proved that the cardinality of $J_0' \langle \alpha, \beta \rangle$ does not exceed the maximum of $\bar{\alpha}$ and $\bar{\beta}$ for α or β infinite.

Theorem 15.1.

- 1) $\max(\alpha, \beta) \leq J_0' \langle \alpha, \beta \rangle$.
- 2) $\alpha < \aleph_\gamma \wedge \beta < \aleph_\gamma \rightarrow J_0' \langle \alpha, \beta \rangle < \aleph_\gamma$.

Proof. 1) If $(\forall \alpha) F' \alpha \triangleq J_0' \langle 0, \alpha \rangle$ then F is a strictly monotonic ordinal function and hence $\alpha \leq F' \alpha$. In particular if $\gamma = \max(\alpha, \beta)$ then

$$\max(\alpha, \beta) = \gamma \leq F' \gamma = J_0' \langle 0, \gamma \rangle.$$

But

$$\langle 0, \gamma \rangle R_0 \langle \alpha, \beta \rangle \vee \langle 0, \gamma \rangle = \langle \alpha, \beta \rangle.$$

Therefore

$$J'_0 \langle 0, \gamma \rangle \leq J'_0 \langle \alpha, \beta \rangle$$

i.e.

$$\max(\alpha, \beta) \leq J'_0 \langle \alpha, \beta \rangle.$$

2) If $J'_0 \langle \alpha, \beta \rangle < \aleph_0$ then $J'_0 \langle \alpha, \beta \rangle < \aleph_\gamma$. If $J'_0 \langle \alpha, \beta \rangle \geq \aleph_0$ then since order isomorphisms map initial segments onto initial segments

$$J'_0 \langle \alpha, \beta \rangle \simeq R_0^{-1} \{ \langle \alpha, \beta \rangle \}.$$

But

$$\begin{aligned} \langle \gamma, \delta \rangle \in R_0^{-1} \langle \alpha, \beta \rangle &\rightarrow \langle \gamma, \delta \rangle R_0 \langle \alpha, \beta \rangle \\ &\rightarrow \max(\gamma, \delta) \leq \max(\alpha, \beta) \\ &\rightarrow [\gamma < \max(\alpha, \beta) + 1] \wedge [\delta < \max(\alpha, \beta) + 1] \\ &\rightarrow \langle \gamma, \delta \rangle \in [\max(\alpha, \beta) + 1] \times [\max(\alpha, \beta) + 1]. \end{aligned}$$

Thus

$$R_0^{-1} \langle \alpha, \beta \rangle \subseteq [\max(\alpha, \beta) + 1] \times [\max(\alpha, \beta) + 1].$$

Consequently $J'_0 \langle \alpha, \beta \rangle$ is equivalent to a subset of

$$[\max(\alpha, \beta) + 1] \times [\max(\alpha, \beta) + 1].$$

From this we see that if $\max(\alpha, \beta)$ were finite then $J'_0 \langle \alpha, \beta \rangle$ would also be finite. Since $J'_0 \langle \alpha, \beta \rangle \geq \aleph_0$ it then follows that $\max(\alpha, \beta) \geq \omega$. Therefore by Theorems 10.22 and 10.34

$$\overline{J'_0 \langle \alpha, \beta \rangle} \leq \overline{[\max(\alpha, \beta) + 1] \times [\max(\alpha, \beta) + 1]} = \overline{\max(\alpha, \beta) + 1} < \aleph_\gamma.$$

Hence

$$J'_0 \langle \alpha, \beta \rangle < \aleph_\gamma.$$

Remark. From the relation R_0 we can define a relation S on $\text{On}^2 \times 9$ that will be used to define the Gödel model. This relation S well orders $\text{On}^2 \times 9$ and is well founded on $\text{On}^2 \times 9$. Consequently the ordering is order isomorphic to On . Indeed for our purposes this order isomorphism is of greater interest than S . We therefore choose to define it directly from J_0 .

Definition 15.2. $J' \langle \alpha, \beta, m \rangle \triangleq 9 \cdot J'_0 \langle \alpha, \beta \rangle + m, m < 9$.

Theorem 15.3. $J : \text{On}^2 \times 9 \xrightarrow[1\text{-}1]{\text{onto}} \text{On}$.

Proof. Clearly α, β and m uniquely determine $9 \cdot J'_0 \langle \alpha, \beta \rangle + m$. Therefore J is a function on $\text{On}^2 \times 9$. If

$$J' \langle \alpha, \beta, m \rangle = J' \langle \gamma, \delta, n \rangle$$

then

$$9 \cdot J'_0 \langle \alpha, \beta \rangle + m = 9 \cdot J'_0 \langle \gamma, \delta \rangle + n.$$

From the uniqueness property in the division theorem for ordinals (Theorem 8.27) it then follows that

$$J'_0 \langle \alpha, \beta \rangle = J'_0 \langle \gamma, \delta \rangle \wedge m = n.$$

But J_0 is one-to-one. Hence $\alpha = \gamma \wedge \beta = \delta$. Therefore

$$\langle \alpha, \beta, m \rangle = \langle \gamma, \delta, n \rangle$$

and hence J is one-to-one.

Again from Theorem 8.27

$$(\forall \gamma) (\exists \delta) (\exists m < 9) [\gamma = 9 \cdot \delta + m].$$

Since $\delta \in \text{On}$ and J_0 is onto

$$(\exists \alpha) (\exists \beta) [\delta = J'_0 \langle \alpha, \beta \rangle].$$

Hence $J' \langle \alpha, \beta, m \rangle = 9 \cdot J'_0 \langle \alpha, \beta \rangle + m = \gamma$ i.e. J is onto.

Definition 15.4.

$$S \triangleq \{ \langle \langle \alpha, \beta, m \rangle, \langle \gamma, \delta, n \rangle \rangle \mid m < 9 \wedge n < 9 \\ \wedge [\langle \alpha, \beta \rangle R_0 \langle \gamma, \delta \rangle \vee [\langle \alpha, \beta \rangle = \langle \gamma, \delta \rangle \wedge m < n]] \}.$$

Theorem 15.5. $J \text{ Isom}_{S,E}(\text{On}^2 \times 9, \text{On})$.

Theorem 15.6.

- 1) $S \text{ We}(\text{On}^2 \times 9)$.
- 2) $S \text{ Wfr}(\text{On}^2 \times 9)$.

The proofs are left to the reader.

Definition 15.7.

$$K_1 \triangleq \{ \langle \gamma, \alpha \rangle \mid (\exists n < 9) (\exists \beta) [\gamma = J' \langle \alpha, \beta, n \rangle] \}.$$

$$K_2 \triangleq \{ \langle \gamma, \beta \rangle \mid (\exists n < 9) (\exists \alpha) [\gamma = J' \langle \alpha, \beta, n \rangle] \}.$$

$$K_3 \triangleq \{ \langle \gamma, n \rangle \mid n < 9 \wedge (\exists \alpha) (\exists \beta) [\gamma = J' \langle \alpha, \beta, n \rangle] \}.$$

Remark. J maps $\text{On}^2 \times 9$ one-to-one onto On . Therefore for each γ in On there is one and only one ordered triple $\langle \alpha, \beta, m \rangle$ in $\text{On}^2 \times 9$ such that $\gamma = J' \langle \alpha, \beta, m \rangle$ that is γ determines an α, β and m such that $\gamma = J' \langle \alpha, \beta, m \rangle$. The function K_1, K_2 , and K_3 are so defined that $K'_1 \gamma, K'_2 \gamma$, and $K'_3 \gamma$ are respectively the first, second and third components of the ordered triple in $\text{On}^2 \times 9$ corresponding to γ under J .

Theorem 15.8. $\gamma = J' \langle K'_1 \gamma, K'_2 \gamma, K'_3 \gamma \rangle$.

Corollary 15.9. *If $m < 9$ then*

- 1) $K'_1 J' \langle \alpha, \beta, m \rangle = \alpha,$
- 2) $K'_2 J' \langle \alpha, \beta, m \rangle = \beta,$
- 3) $K'_3 J' \langle \alpha, \beta, m \rangle = m.$

Details are left to the reader.

Theorem 15.10.

- 1) $K'_1 \gamma \leq \gamma \wedge K'_2 \gamma \leq \gamma.$
- 2) $K'_3 \gamma \neq 0 \rightarrow K'_1 \gamma < \gamma \wedge K'_2 \gamma < \gamma.$

Proof. $\gamma = J' \langle K'_1 \gamma, K'_2 \gamma, K'_3 \gamma \rangle = 9 \cdot J'_0 \langle K'_1 \gamma, K'_2 \gamma \rangle + K'_3 \gamma.$ Therefore from properties of ordinal arithmetic (Corollary 8.5 and Theorem 8.21)

$$J'_0 \langle K'_1 \gamma, K'_2 \gamma \rangle \leq \gamma.$$

But by Theorem 15.1

$$\max(K'_1 \gamma, K'_2 \gamma) \leq J'_0 \langle K'_1 \gamma, K'_2 \gamma \rangle.$$

Thus

$$K'_1 \gamma \leq \gamma \quad \text{and} \quad K'_2 \gamma \leq \gamma.$$

If in addition $K'_3 \gamma \neq 0$ then

$$J'_0 \langle K'_1 \gamma, K'_2 \gamma \rangle < \gamma$$

and hence

$$K'_1 \gamma < \gamma \quad \text{and} \quad K'_2 \gamma < \gamma.$$

Theorem 15.11. $m < 9 \wedge \alpha < \aleph_\gamma \wedge \beta < \aleph_\gamma \rightarrow J' \langle \alpha, \beta, m \rangle < \aleph_\gamma.$

Proof. By Theorem 15.1 we have

$$J'_0 \langle \alpha, \beta \rangle < \aleph_\gamma.$$

If $J'_0 \langle \alpha, \beta \rangle < \aleph_0$ then

$$J' \langle \alpha, \beta, m \rangle = 9 \cdot J'_0 \langle \alpha, \beta \rangle + m < \aleph_0 \leq \aleph_\gamma.$$

If $J'_0 \langle \alpha, \beta \rangle \geq \aleph_0$ then since

$$9 \cdot J'_0 \langle \alpha, \beta \rangle \simeq 9 \times J'_0 \langle \alpha, \beta \rangle$$

we have that

$$\overline{\overline{9 \cdot J'_0 \langle \alpha, \beta \rangle}} = \overline{\overline{J'_0 \langle \alpha, \beta \rangle}} < \aleph_\gamma$$

and hence

$$J' \langle \alpha, \beta, m \rangle = 9 \cdot J'_0 \langle \alpha, \beta \rangle + m < \aleph_\gamma.$$

Theorem 15.12. $J' \langle 0, \aleph_\gamma, 0 \rangle = \aleph_\gamma.$

Proof. If it were the case that $K_1' \aleph_\gamma < \aleph_\gamma$ and $K_2' \aleph_\gamma < \aleph_\gamma$ then by Theorem 15.8 it would follow that

$$\aleph_\gamma = J' \langle K_1' \aleph_\gamma, K_2' \aleph_\gamma, K_3' \aleph_\gamma \rangle < \aleph_\gamma.$$

From this contradiction and Theorem 15.1 we conclude that

$$\aleph_\gamma \leq \max(K_1' \aleph_\gamma, K_2' \aleph_\gamma) \leq J_0' \langle K_1' \aleph_\gamma, K_2' \aleph_\gamma \rangle.$$

Furthermore it follows that

$$\langle 0, \aleph_\gamma \rangle R_0 \langle K_1' \aleph_\gamma, K_2' \aleph_\gamma \rangle \quad \text{or} \quad \langle 0, \aleph_\gamma \rangle = \langle K_1' \aleph_\gamma, K_2' \aleph_\gamma \rangle$$

and hence

$$\aleph_\gamma \leq J_0' \langle 0, \aleph_\gamma \rangle \leq J_0' \langle K_1' \aleph_\gamma, K_2' \aleph_\gamma \rangle.$$

Therefore

$$\aleph_\gamma \leq 9 \cdot J_0' \langle 0, \aleph_\gamma \rangle + 0 \leq 9 \cdot J_0' \langle K_1' \aleph_\gamma, K_2' \aleph_\gamma \rangle + K_3' \aleph_\gamma$$

i.e.

$$\aleph_\gamma \leq J' \langle 0, \aleph_\gamma, 0 \rangle \leq J' \langle K_1' \aleph_\gamma, K_2' \aleph_\gamma, K_3' \aleph_\gamma \rangle = \aleph_\gamma.$$

Thus

$$J' \langle 0, \aleph_\gamma, 0 \rangle = \aleph_\gamma.$$

Remark. We are now ready to define the Gödel model L . This is a standard transitive model that we define as the range of a special function F that is in turn defined by transfinite recursion in the following way.

Definition 15.13.

$$\begin{aligned} G'x &\triangleq \mathcal{W}(x) \quad \text{if} \quad K_3' \mathcal{D}(x) = 0 \\ &\triangleq \mathcal{F}_n(x' K_1' \mathcal{D}(x), x' K_2' \mathcal{D}(x)) \quad \text{if} \quad K_3' \mathcal{D}(x) = n \neq 0. \end{aligned}$$

$$F \mathcal{F}_n \text{ On } \wedge (\forall \alpha) [F' \alpha \triangleq G'(F \Gamma \alpha)].$$

Theorem 15.14.

$$\begin{aligned} F' \alpha &= F'' \alpha \quad \text{if} \quad K_3' \alpha = 0, \\ F' \alpha &= \mathcal{F}_n'(F' K_1' \alpha, F' K_2' \alpha) \quad \text{if} \quad K_3' \alpha = n \neq 0. \end{aligned}$$

Proof. Since $\mathcal{D}(F \Gamma \alpha) = \alpha$ we have

$$\begin{aligned} F' \alpha &= G'(F \Gamma \alpha) = \mathcal{W}(F \Gamma \alpha) = F'' \alpha \quad \text{if} \quad K_3' \alpha = 0, \\ F' \alpha &= G'(F \Gamma \alpha) = \mathcal{F}_n((F \Gamma \alpha)' K_1' \alpha, (F \Gamma \alpha)' K_2' \alpha) \quad \text{if} \quad K_3' \alpha = n \neq 0. \end{aligned}$$

But by Theorem 15.10

$$K_3' \alpha \neq 0 \rightarrow K_1' \alpha < \alpha \wedge K_2' \alpha < \alpha.$$

Thus

$$(F \ulcorner \alpha)' K_1' \alpha = F' K_1' \alpha \quad \text{and} \quad (F \ulcorner \alpha)' K_2' \alpha = F' K_2' \alpha.$$

Consequently

$$F' \alpha = \mathcal{F}_n'(F' K_1' \alpha, F' K_2' \alpha) \quad \text{if} \quad K_3' \alpha = n \neq 0.$$

Examples.

$$\begin{aligned} J' \langle 0, 0, 0 \rangle = 0 & \quad K_3' 0 = 0 & \quad F' 0 = F'' 0 = 0. \\ J' \langle 0, 0, 1 \rangle = 1 & \quad K_3' 1 = 1 & \quad F' 1 = \mathcal{F}_1'(F' 0, F' 0) = \{0\} = 1. \\ J' \langle 0, 0, 2 \rangle = 2 & \quad K_3' 2 = 2 & \quad F' 2 = \mathcal{F}_2'(F' 0, F' 0) = F' 0 \cap E = 0. \\ J' \langle 0, 0, 3 \rangle = 3 & \quad K_3' 3 = 3 & \quad F' 3 = \mathcal{F}_3'(F' 0, F' 0) = F' 0 - F' 0 = 0. \\ J' \langle 0, 0, 4 \rangle = 4 & \quad K_3' 4 = 4 & \quad F' 4 = \mathcal{F}_4'(F' 0, F' 0) = F' 0 \ulcorner F' 0 = 0. \\ J' \langle 0, 0, 5 \rangle = 5 & \quad K_3' 5 = 5 & \quad F' 5 = \mathcal{F}_5'(F' 0, F' 0) = F' 0 \cap \mathcal{D}(F' 0) = 0. \\ J' \langle 0, 0, 6 \rangle = 6 & \quad K_3' 6 = 6 & \quad F' 6 = \mathcal{F}_6'(F' 0, F' 0) = F' 0 \cap (F' 0)^{-1} = 0. \\ J' \langle 0, 0, 7 \rangle = 7 & \quad K_3' 7 = 7 & \quad F' 7 = \mathcal{F}_7'(F' 0, F' 0) = F' 0 \cap \text{Cnv}_2(F' 0) = 0. \\ J' \langle 0, 0, 8 \rangle = 8 & \quad K_3' 8 = 8 & \quad F' 8 = \mathcal{F}_8'(F' 0, F' 0) = F' 0 \cap \text{Cnv}_3(F' 0) = 0. \\ J' \langle 0, 1, 0 \rangle = 9 & \quad K_3' 9 = 0 & \quad F' 9 = F'' 9 = \{0, 1\} = 2. \end{aligned}$$

Definition 15.15. $L \triangleq F'' \text{On}$.

A set a is constructible iff $a \in L$.

Remark. We will prove that L is a model of ZF by proving that L is transitive, almost universal and closed under the eight fundamental operations. Those classes that are L -constructible in the sense of Definition 14.12 we will refer to simply as constructible classes. The elements of L we will call constructible sets. Indeed if $x = F' \alpha$ we refer to x as the set constructed at the α^{th} stage. From the foregoing example we see that 0 is the set constructed at the 0th, 2nd, 3rd, 4th, 5th, 6th, 7th, and 8th stages. Also 1 is constructed at the first stage and 2 is constructed at the 9th stage. Thus the constructible sets are those sets that can be “built” up from the empty set by a finite or transfinite number of applications of the eight fundamental operations.

Closely related to the notion of constructibility is the notion of relative constructibility. There are many ways to generalize the notion of constructibility. For Cohen’s work we wish to introduce an arbitrary set a of natural numbers at the $(\omega + 1)^{\text{th}}$ stage. Since we wish to construct models of ZF and every such model must contain ω and all of its elements we modify the first $(\omega + 1)$ -stage to introduce ω and its elements in a most direct and obvious way. A set is then constructible relative to a iff it can be built up from ω and its elements and from a by a finite or transfinite number of applications of the eight fundamental operations.

Although relative constructibility will not be needed until later we introduce it here because the definition and theorems of interest so closely parallel those for constructibility.

Definition 15.16. $\forall a \subseteq \omega$,

$$\begin{aligned} G_a'x &= \mathcal{W}(x) \quad \text{if } \mathcal{D}(x) < \omega + 1 \vee K_3' \mathcal{D}(x) = 0 \\ &= a \quad \text{if } \mathcal{D}(x) = \omega + 1 \\ &= \mathcal{F}_n(x'K_1' \mathcal{D}(x), x'K_2' \mathcal{D}(x)) \quad \text{if } \mathcal{D}(x) > \omega + 1 \wedge K_3' \mathcal{D}(x) = n \neq 0. \\ &F_a \mathcal{F}_n \text{ On} \wedge (\forall \alpha) [F_a' \alpha = G_a(F_a \uparrow \alpha)] \\ &L_a = F_a'' \text{ On}. \end{aligned}$$

Theorem 15.17.

$$\begin{aligned} F_a' \alpha &= \alpha, \quad \alpha \leq \omega \\ &= a, \quad \alpha = \omega + 1 \\ &= F_a'' \alpha, \quad \alpha > \omega + 1 \wedge K_3' \alpha = 0 \\ &= \mathcal{F}_n(F_a' K_1' \alpha, F_a' K_2' \alpha), \quad \alpha > \omega + 1 \wedge K_3' \alpha = n \neq 0. \end{aligned}$$

The proof is left to the reader.

Definition 15.18.

- 1) $\text{Od}'x = \mu_\alpha(x = F' \alpha).$
- 2) $\text{Od}'_a x = \mu_\alpha(x = F_a' \alpha).$

Remark. The symbol $\text{Od}'x$ is read “the order of x .” If x is constructible then $\text{Od}'x$ is the smallest ordinal α for which $x = F' \alpha$ i.e., $\text{Od}'x$ is the first stage at which x is constructed.

Theorem 15.19.

- 1) $x \in L \iff x = F' \text{Od}'x.$
- 2) $x \in L_a \iff x = F_a' \text{Od}'_a x.$

Proof. Definition 15.18.

Remark. We wish to prove that L is transitive. For this we prove that the set constructed at the α^{th} stage is constructed only from sets that were constructed at earlier stages.

Theorem 15.20.

- 1) $(\forall \alpha) [F' \alpha \subseteq F'' \alpha].$
- 2) $(\forall \alpha) [F_a' \alpha \subseteq F_a'' \alpha].$

Proof. 1) (By transfinite induction). If $\beta = K'_1\alpha, \gamma = K'_2\alpha$ and $n = K'_3\alpha$ then

$$\alpha = J'\langle\beta, \gamma, n\rangle.$$

If $n = 0$ then $F'\alpha = F''\alpha$ and hence $F'\alpha \subseteq F''\alpha$. If $n \neq 0$ then by Theorem 15.10, $\beta < \alpha, \gamma < \alpha$ and hence

$$F'\beta \in F''\alpha \wedge F'\gamma \in F''\alpha.$$

If $n = 1$ then

$$F'\alpha = \mathcal{F}_1(F'\beta, F'\gamma) = \{F'\beta, F'\gamma\} \subseteq F''\alpha.$$

If $n > 1$ then

$$F'\alpha = \mathcal{F}_n(F'\beta, F'\gamma) \subseteq F'\beta.$$

From the induction hypothesis and the fact that $\beta < \alpha$ we have

$$F'\beta \subseteq F''\beta \subseteq F''\alpha.$$

Therefore

$$F'\alpha \subseteq F''\alpha.$$

2) The proof is left to the reader.

Theorem 15.21.

1) $\text{Tr}(F''\alpha)$.

2) $\text{Tr}(F''_a\alpha)$.

Proof.

1) $x \in F''\alpha \rightarrow (\exists \beta < \alpha) [x = F'\beta]$. But from Theorem 15.20 and the fact that $\beta < \alpha$ we have

$$x = F'\beta \subseteq F''\beta \subseteq F''\alpha.$$

2) The proof is left to the reader.

Theorem 15.22.

1) $\text{Tr}(L)$.

2) $\text{Tr}(L_a)$.

Proof.

1) $x \in L \rightarrow (\exists \alpha) [x = F'\alpha]$. Then

$$x = F'\alpha \in F''(\alpha + 1).$$

Since $F''(\alpha + 1)$ is transitive

$$x \subseteq F''(\alpha + 1) \subseteq L.$$

2) The proof is left to the reader.

Theorem 15.23.

- 1) $x \in L \wedge y \in L \wedge x \in y \rightarrow \text{Od}'x < \text{Od}'y.$
- 2) $x \in L_a \wedge y \in L_a \wedge x \in y \rightarrow \text{Od}'_a x < \text{Od}'_a y.$

Proof.

- 1) $x \in y \wedge y \in L \rightarrow x \in F' \text{Od}'y$
 $\rightarrow x \in F'' \text{Od}'y$
 $\rightarrow (\exists \beta < \text{Od}'y) [x = F'\beta]$
 $\rightarrow \text{Od}'x < \text{Od}'y.$
- 2) The proof is left to the reader.

Theorem 15.24.

$$(\forall x \in L_a) (\exists \alpha > \omega) [x = F'_a \alpha].$$

The proof is left to the reader.

Theorem 15.25.

- 1) $(\forall x \in L) (\forall y \in L) [\mathcal{F}_n(x, y) \in L], n = 1, \dots, 8.$
- 2) $(\forall x \in L_a) (\forall y \in L_a) [\mathcal{F}_n(x, y) \in L_a], n = 1, \dots, 8.$

Proof. 1) If $\alpha = \text{Od}'x \wedge \beta = \text{Od}'y$ then $x = F'\alpha \wedge y = F'\beta$. If $\gamma = J'\langle \alpha, \beta, n \rangle$ then

$$\mathcal{F}_n(x, y) = \mathcal{F}_n(F'\alpha, F'\beta) = \mathcal{F}_n(F'K'_1\gamma, F'K'_2\gamma) = F'\gamma \in L.$$

- 2) The proof is left to the reader.

Theorem 15.26.

- 1) $b \subseteq L \rightarrow (\exists x \in L) [b \subseteq x].$
- 2) $b \subseteq L_a \rightarrow (\exists x \in L_a) [b \subseteq x].$

Proof. 1) Since Od is a function from V into On , $\text{Od}''b$ is a set of ordinals. Therefore

$$(\exists \alpha) [\text{Od}''b \subseteq \alpha].$$

If

$$\beta = J'\langle 0, \alpha, 0 \rangle$$

then $K'_3\beta = 0$ and hence

$$F'\beta = F''\beta.$$

Furthermore $\alpha \leq \beta$. Therefore if $x = F'\beta$

$$y \in b \rightarrow y \in L \wedge \text{Od}'y < \alpha \leq \beta,$$

and hence

$$(\exists \gamma < \beta) [y = F'\gamma].$$

Then

$$y = F'\gamma \in F''\beta = F'\beta = x$$

that is

$$b \subseteq x \wedge x \in L.$$

2) The proof is left to the reader.

Theorem 15.27. 1) L is a standard transitive model of ZF and $\text{On} \subseteq L$.

2) L_a is a standard transitive model of ZF and $\text{On} \subseteq L_a$.

Proof. Theorems 15.22, 15.25, 15.26, and 14.11.

Remark. We have now shown that L is a model of ZF and for each $a \subseteq \omega$, L_a is a model of ZF; but are these models different? It is not difficult to show that if a is constructible then $L_a = L$. Do there exist nonconstructible sets? From Cohen's work we know that this question is undecidable in ZF.

The assumption that every set is constructible is called the Axiom of Constructibility.

Axiom of Constructibility.

$$V = L.$$

Gödel's program for proving the consistency of the GCH and the AC consists of proving that the Axiom of Constructibility implies the GCH and the AC. It is then sufficient to establish the consistency of the Axiom of Constructibility with ZF. This is done by proving that L is a model of $V = L$. To prove this we must prove in ZF that

$$[V = L]^L$$

i.e. since $L = \{x \mid (\exists \alpha) [x = F'\alpha]\}$ we must prove that

$$L = \{x \in L \mid (\exists \alpha \in L) [x = F'\alpha]^L\}.$$

Since $\text{On} \subseteq L$ it is sufficient to prove that $x = F'\alpha$ is absolute with respect to L . This we will do by proving that $x = F'\alpha$ is absolute with respect to every standard transitive model of ZF. We need the following lemmas in which \mathcal{M} is a standard transitive model of ZF and G is as given in Definition 15.13.

Lemma 1. $\langle \alpha, \beta, m \rangle S \langle \gamma, \delta, n \rangle \text{ Abs } \mathcal{M}$.

Proof.

$$\langle \alpha, \beta, m \rangle S \langle \gamma, \delta, n \rangle \longleftrightarrow \langle \alpha, \beta \rangle R_0 \langle \gamma, \delta \rangle \vee [\langle \alpha, \beta \rangle = \langle \gamma, \delta \rangle \wedge m < n].$$

Lemma 2. $f \text{ Isom}_{S,E}(\beta^2 \times 9, \alpha) \text{ Abs } \mathcal{M}.$

Proof. Theorem 13.30.

Lemma 3.

- 1) $\beta = J^* \langle \gamma, \delta, m \rangle \text{ Abs } \mathcal{M}.$
- 2) $\beta = J_0^* \langle \gamma, \delta \rangle \text{ Abs } \mathcal{M}.$

Proof. 1) From properties of order isomorphisms (Theorem 7.51)

$$(\exists! f) (\exists! \alpha) [f \text{ Isom}_{S,E}(\mu \times \mu \times 9, \alpha)].$$

Therefore, from the definition of J

$$\begin{aligned} \beta = J^* \langle \gamma, \delta, m \rangle &\longleftrightarrow (\exists f) (\exists \alpha) [m < 9 \wedge f \text{ Isom}_{S,E}(\max(\gamma, \delta) \\ &\quad \times \max(\gamma, \delta) \times 9, \alpha) \wedge f^* \langle \gamma, \delta, m \rangle = \beta] \\ &\longleftrightarrow (\forall f) (\forall \alpha) [m < 9 \wedge f \text{ Isom}_{S,E}(\max(\gamma, \delta) \\ &\quad \times \max(\gamma, \delta) \times 9, \alpha) \rightarrow f^* \langle \gamma, \delta, m \rangle = \beta]. \end{aligned}$$

From Theorem 13.8 it then follows that

$$\beta = J^* \langle \gamma, \delta, m \rangle \text{ Abs } \mathcal{M}.$$

- 2) The proof is left to the reader.

Lemma 4.

- 1) $K_1^* \alpha = \beta \text{ Abs } \mathcal{M}.$
- 2) $K_2^* \alpha = \beta \text{ Abs } \mathcal{M}.$
- 3) $K_3^* \alpha = \beta \text{ Abs } \mathcal{M}.$

Proof. $K_1^* \alpha = \beta \longleftrightarrow (\exists m) (\exists \gamma) [m < 9 \wedge J^* \langle \beta, \gamma, m \rangle = \alpha].$
 $K_2^* \alpha = \beta \longleftrightarrow (\exists m) (\exists \gamma) [m < 9 \wedge J^* \langle \gamma, \beta, m \rangle = \alpha].$
 $K_3^* \alpha = \beta \longleftrightarrow \beta < 9 \wedge (\exists \gamma) (\exists \delta) [J^* \langle \gamma, \delta, m \rangle = \alpha].$

Since $J^* \langle \beta, \gamma, m \rangle = \alpha$ is absolute with respect to \mathcal{M} and $\max(\beta, \gamma) \leq J^* \langle \beta, \gamma, m \rangle = \alpha$ it follows that $\alpha \in \mathcal{M}$ implies $\beta, \gamma \in \mathcal{M}$. The results then follow from Theorem 13.5.

Lemma 5.

$$b = \mathcal{F}_n(c, d) \text{ Abs } \mathcal{M}, \quad n = 1, \dots, 8.$$

Proof.

$$b = \mathcal{F}_1(c, d) \longleftrightarrow b = \{c, d\}.$$

$$\begin{aligned} b = \mathcal{F}_2(c, d) &\longleftrightarrow b = c \cap E \\ &\longleftrightarrow (\forall x) [x \in b \longleftrightarrow x \in c \\ &\quad \wedge (\exists y) (\exists z) [x = \langle y, z \rangle \wedge y \in z]]. \end{aligned}$$

$$\begin{aligned} b = \mathcal{F}_3(c, d) &\longleftrightarrow b = c - d \\ &\longleftrightarrow (\forall x) [x \in b \longleftrightarrow x \in c \wedge x \notin d]. \end{aligned}$$

$$\begin{aligned} b = \mathcal{F}_4(c, d) &\longleftrightarrow b = c \Gamma d \\ &\longleftrightarrow (\forall x) [x \in b \longleftrightarrow x \in c \\ &\quad \wedge (\exists y) (\exists z) [x = \langle y, z \rangle \wedge y \in d]]. \end{aligned}$$

$$\begin{aligned} b = \mathcal{F}_5(c, d) &\longleftrightarrow b = c \cap \mathcal{D}(d) \\ &\longleftrightarrow (\forall x) [x \in b \longleftrightarrow x \in c \\ &\quad \wedge (\exists y) [y = \mathcal{D}(d) \wedge x \in y]]. \end{aligned}$$

$$\begin{aligned} b = \mathcal{F}_6(c, d) &\longleftrightarrow b = c \cap d^{-1} \\ &\longleftrightarrow (\forall x) [x \in b \longleftrightarrow x \in c \\ &\quad \wedge (\exists y) [y = d^{-1} \wedge x \in y]]. \end{aligned}$$

$$\begin{aligned} b = \mathcal{F}_7(c, d) &\longleftrightarrow b = c \cap \text{Cnv}_2(d) \\ &\longleftrightarrow (\forall x) [x \in b \longleftrightarrow x \in c \\ &\quad \wedge (\exists u) (\exists v) (\exists w) [x = \langle u, v, w \rangle \wedge \langle v, w, u \rangle \in d]]. \end{aligned}$$

$$\begin{aligned} b = \mathcal{F}_8(c, d) &\longleftrightarrow b = c \cap \text{Cnv}_3(d) \\ &\longleftrightarrow (\forall x) [x \in b \longleftrightarrow x \in c \\ &\quad \wedge (\exists u) (\exists v) (\exists w) [x = \langle u, v, w \rangle \wedge \langle u, w, v \rangle \in d]]. \end{aligned}$$

Lemma 6.

$$b = G'(f \Gamma \beta) \text{ Abs } \mathcal{M}.$$

Proof.

$$\begin{aligned} b = G'(f \Gamma \beta) &\longleftrightarrow [K'_3 \beta = 0 \wedge b = f'' \beta] \vee [K'_3 \beta = 1 \\ \wedge b = \mathcal{F}_1(f' K'_1 \beta, f' K'_2 \beta)] &\vee \dots \vee [K'_3 \beta = 8 \wedge b = \mathcal{F}_8(f' K'_1 \beta, f' K'_2 \beta)]. \end{aligned}$$

Theorem 15.28.

- 1) $b = F' \alpha \text{ Abs } \mathcal{M}.$
- 2) $b = F'_a \alpha \text{ Abs } \mathcal{M}.$

Proof. 1) From the definition of F and Corollary 7.41

$$(\exists ! f) [f \mathcal{F}_n(\alpha + 1) \wedge (\forall \beta \leq \alpha) [f' \beta = G(f \Gamma \beta)]].$$

Therefore

$$b = F'\alpha \longleftrightarrow (\exists f) [f \mathcal{F}_n(\alpha + 1) \wedge (\forall \beta \leq \alpha) [f'\beta = G(f \Gamma \beta)] \wedge \langle \alpha, b \rangle \in f] \\ \longleftrightarrow (\forall f) [f \mathcal{F}_n(\alpha + 1) \wedge (\forall \beta \leq \alpha) [f'\beta = G(f \Gamma \beta)] \rightarrow \langle \alpha, b \rangle \in f].$$

From the preceding lemmas and Theorem 13.8 it then follows that $b = F'\alpha \text{ Abs } \mathcal{M}$.

2) The proof is left to the reader.

Theorem 15.29.

1) $\text{On} \subseteq \mathcal{M} \rightarrow L \subseteq \mathcal{M}$.

2) $\text{On} \subseteq \mathcal{M} \wedge a \in \mathcal{M} \rightarrow L_a \subseteq \mathcal{M}$.

Proof. 1) $(\forall \alpha) (\exists x) [x = F'\alpha]$.

Therefore

$$(\forall \alpha \in \mathcal{M}) (\exists x \in \mathcal{M}) [x = F'\alpha]^{\mathcal{M}}.$$

But since $\text{On} \subseteq \mathcal{M}$ and $x = F'\alpha \text{ Abs } \mathcal{M}$

$$(\forall \alpha) (\exists x \in \mathcal{M}) [x = F'\alpha].$$

Therefore

$$L \subseteq \mathcal{M}.$$

2) The proof is left to the reader.

Remark. From Theorem 15.29 we see that L is the smallest of all the standard transitive models that contain On . In particular $(\forall a \subseteq \omega) [L \subseteq L_a]$. Furthermore if a is constructible i.e. if $a \in L$ then $L_a \subseteq L$ i.e. $L = L_a$.

Theorem 15.30.

1) L is a model of $V = L$.

2) L_a is a model of $V = L_a$.

Proof. 1) $V = L \longleftrightarrow (\forall x) (\exists \alpha) [x = F'\alpha]$. From the definition of L

$$(\forall x \in L) (\exists \alpha) [x = F'\alpha].$$

Since $\text{On} \subseteq L$ and $x = F'\alpha \text{ Abs } L$ it follows that

$$(\forall x \in L) (\exists \alpha \in L) [x = F'\alpha]^L$$

i.e.

$$[V = L]^L.$$

2) The proof is left to the reader.

Definition 15.31.

- 1) $As = \{\langle x, y \rangle \in L^2 \mid y \in x \wedge (\forall z \in x) [Od'y \leq Od'z]\}$.
- 2) $As_a = \{\langle x, y \rangle \in L_a^2 \mid y \in x \wedge (\forall z \in x) [Od'_a y \leq Od'_a z]\}$.

Theorem 15.32.

- 1) $(\forall x \in L) [x \neq 0 \rightarrow As'x \in x]$.
- 2) $(\forall x \in L_a) [x \neq 0 \rightarrow As'_a x \in x]$.

Proof. 1) Since L is transitive $L - \{0\} \subseteq \mathcal{D}(As)$. Furthermore As is single valued. Therefore

$$x \in L \wedge x \neq 0 \rightarrow As'x \in x.$$

- 2) The proof is left to the reader.

Theorem 15.33.

- 1) $V = L \rightarrow AC$.
- 2) $V = L_a \rightarrow AC$.

Proof. Obvious from Theorem 15.32.

Theorem 15.34.

- 1) L is a model of the AC.
- 2) L_a is a model of the AC.

Proof. Theorems 15.30 and 15.33.

Remark. In Theorem 15.33 we have a result that is in fact stronger than the strong form of the AC. The strong form of the AC asserts the existence of a universal choice function. In the proof of Theorem 15.33 we have exhibited such a function.

We turn now to a proof that the Axiom of Constructibility implies the GCH. It is of course sufficient to prove that it implies the Aleph Hypothesis:

$$(\forall \alpha) [\overline{\overline{2^{\aleph_\alpha}}} = \aleph_{\alpha+1}].$$

The key to the proof lies in proving two results. First, we prove that the cardinality of $F''\aleph_\alpha$ is \aleph_α . From this we deduce the AH by proving that if $V = L$ then every subset of $F''\aleph_\alpha$ is constructed before the $\aleph_{\alpha+1}$ th stage.

Definition 15.35.

- 1) $C'\alpha \triangleq Od'As'F'\alpha$.
- 2) $C'_a\alpha \triangleq Od'_aAs'_aF'_a\alpha$.

Theorem 15.36.

- 1) $C'\alpha \leq \alpha.$
- 2) $C'_a\alpha \leq \alpha.$

Proof. 1) If $F'\alpha = 0$ then $As'F'\alpha = 0$ and $Od'As'F'\alpha = 0$ i.e. $C'\alpha = 0 \leq \alpha.$ If $F'\alpha \neq 0$ then $As'F'\alpha \in F'\alpha.$ Therefore

$$C'\alpha = Od'As'F'\alpha < Od'F'\alpha \leq \alpha.$$

- 2) The proof is left to the reader.

Theorem 15.37.

- 1) $\overline{\overline{F''\aleph_\alpha}} = \aleph_\alpha.$
- 2) $\overline{\overline{F'_a\aleph_\alpha}} = \aleph_\alpha.$

Proof. 1) Since F is a function it follows that

$$\overline{\overline{F''\aleph_\alpha}} \leq \aleph_\alpha.$$

Furthermore

$$F'J'\langle 0, \beta, 0 \rangle = F''J'\langle 0, \beta, 0 \rangle.$$

Therefore since $\gamma < \beta$ implies $J'\langle 0, \gamma, 0 \rangle < J'\langle 0, \beta, 0 \rangle$ it follows that if $\gamma < \beta$ then

$$F'J'\langle 0, \gamma, 0 \rangle \in F'J'\langle 0, \beta, 0 \rangle$$

that is

$$\gamma \neq \beta \rightarrow F'J'\langle 0, \gamma, 0 \rangle \neq F'J'\langle 0, \beta, 0 \rangle.$$

Thus if

$$H'\beta \triangleq F'J'\langle 0, \beta, 0 \rangle, \quad \beta \in \text{On}$$

then $H : \text{On} \xrightarrow{1-1} L.$ Since

$$\beta < \aleph_\alpha \rightarrow J'\langle 0, \beta, 0 \rangle < \aleph_\alpha$$

it follows that

$$H''\aleph_\alpha \subseteq F''\aleph_\alpha.$$

Since H is one-to-one

$$\aleph_\alpha = \overline{\overline{H''\aleph_\alpha}} \leq \overline{\overline{F''\aleph_\alpha}}.$$

Therefore $\overline{\overline{F''\aleph_\alpha}} = \aleph_\alpha.$

- 2) The proof is left to the reader.

Theorem 15.38.

- 1) $(\forall \alpha) [\mathcal{P}(F''\aleph_\alpha) \subseteq F''\aleph_{\alpha+1} \rightarrow \overline{2^{\aleph_\alpha}} = \aleph_{\alpha+1}]$.
- 2) $(\forall \alpha) [\mathcal{P}(F'_a''\aleph_\alpha) \subseteq F'_a''\aleph_{\alpha+1} \rightarrow \overline{2^{\aleph_\alpha}} = \aleph_{\alpha+1}]$.

Proof. 1) If $\mathcal{P}(F''\aleph_\alpha) \subseteq F''\aleph_{\alpha+1}$ then from Theorem 15.37

$$\overline{2^{\aleph_\alpha}} = \overline{\mathcal{P}(\aleph_\alpha)} = \overline{\mathcal{P}(F''\aleph_\alpha)} \leq \overline{F''\aleph_{\alpha+1}} = \aleph_{\alpha+1}.$$

Since by Cantor's Theorem $\overline{2^{\aleph_\alpha}} > \aleph_\alpha$ we have

$$\overline{2^{\aleph_\alpha}} = \aleph_{\alpha+1}.$$

- 2) The proof is left to the reader.

Remark. It now remains to be proved that if $V = L$ then each subset of $F''\aleph_\alpha$ is constructed before the $\aleph_{\alpha+1}$ th stage. This we do in the following sequence of theorems.

Theorem 15.39.

1) If $V = L$ and $(\forall b) (\forall \gamma) (\forall f) [9 \subseteq b \subseteq \text{On} \wedge C''b \subseteq b \wedge K_1''b \subseteq b \wedge K_2''b \subseteq b \wedge J''b^3 \subseteq b \wedge f \text{ Isom}_{E,E}(b, \gamma) \rightarrow (\forall \alpha \in b) (\forall \beta \in b) [F'\alpha \in F'\beta \leftrightarrow F'f'\alpha \in F'f'\beta]]$ then $(\forall \alpha) [\mathcal{P}(F''\aleph_\alpha) \subseteq F''\aleph_{\alpha+1}]$.

2) If $V = L_\alpha$ and $(\forall b) (\forall \gamma) (\forall f) [\omega + 2 \subseteq b \subseteq \text{On} \wedge C''_a b \subseteq b \wedge K_1''_a b \subseteq b \wedge K_2''_a b \subseteq b \wedge J''_a b^3 \subseteq b \wedge f \text{ Isom}_{E,E}(b, \gamma) \rightarrow (\forall \alpha \in b) (\forall \beta \in b) [F'_a \alpha \in F'_a \beta \leftrightarrow F'_a f'_a \alpha \in F'_a f'_a \beta]]$ then $(\forall \alpha) [\mathcal{P}(F'_a''\aleph_\alpha) \subseteq F'_a''\aleph_{\alpha+1}]$.

Proof. 1) If $x \subseteq F''\aleph_\alpha$ then from the Axiom of Constructibility

$$(\exists \delta) [x = F'\delta].$$

Since C, K_1, K_2 and J are each single valued and $\aleph_\alpha \cup \{\delta\}$ is infinite it follows from Theorem 11.29 that

$$(\exists b) [C''b \subseteq b \wedge K_1''b \subseteq b \wedge K_2''b \subseteq b \wedge J''b^3 \subseteq b \wedge \aleph_\alpha \cup \{\delta\} \subseteq b \subseteq \text{On} \wedge \bar{b} = \aleph_\alpha].$$

Since $V = L$ implies the AC, $(\exists \gamma) (\exists f) f \text{ Isom}_{E,E}(b, \gamma)$. By hypothesis

$$f \text{ Isom}_{E,E}(b, \gamma) \rightarrow (\forall \alpha \in b) (\forall \beta \in b) [F'\alpha \in F'\beta \leftrightarrow F'f'\alpha \in F'f'\beta].$$

In particular, since $\aleph_\alpha \subseteq b$ and f is order preserving

$$\beta < \aleph_\alpha \rightarrow f'\beta = \beta.$$

Thus, since $\delta \in b$, if $\beta < \aleph_\alpha$

$$[F'\beta \in F'\delta \leftrightarrow F'\beta \in F'f'\delta].$$

Consequently

$$F'\delta \cap F''\aleph_\alpha = F'f'\delta \cap F''\aleph_\alpha.$$

But since $\aleph_\alpha = J\langle 0, \aleph_\alpha, 0 \rangle$,

$$F''\aleph_\alpha = F'\aleph_\alpha.$$

Furthermore, since $F'\delta = x \subseteq F''\aleph_\alpha$,

$$F'\delta \cap F''\aleph_\alpha = F'\delta = x.$$

Then

$$\begin{aligned} x &= F'f'\delta \cap F'\aleph_\alpha \\ &= F'f'\delta - [F'f'\delta - F'\aleph_\alpha] \\ &= F'J\langle f'\delta, J\langle f'\delta, \aleph_\alpha, 3 \rangle, 3 \rangle. \end{aligned}$$

But since $\delta \in b \wedge \bar{b} = \aleph_\alpha$ and since f is an order isomorphism $f'\delta < \aleph_{\alpha+1}$. Therefore by Theorem 15.1

$$J\langle f'\delta, \aleph_\alpha, 3 \rangle < \aleph_{\alpha+1}$$

and hence by a second application of Theorem 15.1

$$J\langle f'\delta, J\langle f'\delta, \aleph_\alpha, 3 \rangle, 3 \rangle < \aleph_{\alpha+1}.$$

Consequently

$$x \in F''\aleph_{\alpha+1}.$$

2) The proof is left to the reader.

Remark. In somewhat over simplified terms Theorem 15.39 states that if every set is constructible and if certain ordinal isomorphisms preserve the order of constructibility then each subset of $F''\aleph_\alpha$ is constructed before the $\aleph_{\alpha+1}$ th stage.

Since for any set of ordinals b

$$(\exists! \gamma) (\exists! f) [f \text{ Isom}_{E,E}(b, \gamma)]$$

it remains to be proved that for appropriately chosen sets b , namely those closed under C, K_1, K_2, K_3 , and J, f does preserve the order of constructibility.

Theorem 15.40. *If $9 \subseteq b \subseteq \text{On} \wedge K_1''b \subseteq b \wedge K_2''b \subseteq b \wedge J''b^3 \subseteq b \wedge f \text{ Isom}_{E,E}(b, \eta)$ then*

$$1) \quad m < 9 \wedge \alpha \in b \wedge \beta \in b \rightarrow J\langle f'\alpha, f'\beta, m \rangle = f'J\langle \alpha, \beta, m \rangle$$

and

$$2) \quad J''\eta^3 \subseteq \eta.$$

Proof. 1) Since J is order preserving

$$J\langle f'\alpha, f'\beta, m \rangle < J\langle f'\gamma, f'\delta, n \rangle \longleftrightarrow \langle f'\alpha, f'\beta, m \rangle S \langle f'\gamma, f'\delta, n \rangle.$$

But f is also order preserving. Therefore

$$\langle f'\alpha, f'\beta, m \rangle S \langle f'\gamma, f'\delta, n \rangle \longleftrightarrow \langle \alpha, \beta, m \rangle S \langle \gamma, \delta, n \rangle:$$

If there exists an ordered triple $\langle \alpha, \beta, m \rangle$ such that

$$J'\langle f'\alpha, f'\beta, m \rangle \neq f'J'\langle \alpha, \beta, m \rangle$$

then there is an S -minimal such element. We will show that the assumption that $\langle \alpha, \beta, m \rangle$ is such an ordered triple leads to a contradiction.

If $J'\langle f'\alpha, f'\beta, m \rangle < f'J'\langle \alpha, \beta, m \rangle$ then since b is closed w.r.t. J and since α, β , and m are in b it follows that

$$J'\langle \alpha, \beta, m \rangle \in b$$

and hence

$$f'J'\langle \alpha, \beta, m \rangle \in \eta.$$

Since

$$J'\langle f'\alpha, f'\beta, m \rangle < f'J'\langle \alpha, \beta, m \rangle$$

it follows that

$$(\exists v \in b) [f'v = J'\langle f'\alpha, f'\beta, m \rangle].$$

If

$$\gamma = K_1'v \wedge \delta = K_2'v \wedge n = K_3'v$$

then since $v \in b$ and b is closed w.r.t. K_1, K_2, K_3

$$\gamma \in b \wedge \delta \in b \wedge n \in b.$$

Therefore

$$v = J'\langle \gamma, \delta, n \rangle \in b$$

and

$$f'J'\langle \gamma, \delta, n \rangle = f'v = J'\langle f'\alpha, f'\beta, m \rangle.$$

Since by hypothesis

$$J'\langle f'\alpha, f'\beta, m \rangle < f'J'\langle \alpha, \beta, m \rangle$$

we have that

$$f'J'\langle \gamma, \delta, n \rangle < f'J'\langle \alpha, \beta, m \rangle.$$

From this it follows that

$$\langle \gamma, \delta, n \rangle S \langle \alpha, \beta, m \rangle.$$

But from the defining property of $\langle \alpha, \beta, m \rangle$

$$J'\langle f'\gamma, f'\delta, n \rangle = f'J'\langle \gamma, \delta, n \rangle = J'\langle f'\alpha, f'\beta, m \rangle.$$

Since both J and f are one-to-one this implies that

$$\langle \gamma, \delta, n \rangle = \langle \alpha, \beta, m \rangle$$

which is a contradiction.

If $f'J'\langle \alpha, \beta, m \rangle < J'\langle f'\alpha, f'\beta, m \rangle$ and if

$$\gamma = K'_1 f'J'\langle \alpha, \beta, m \rangle \wedge \delta = K'_2 f'J'\langle \alpha, \beta, m \rangle \wedge n = K'_3 f'J'\langle \alpha, \beta, m \rangle$$

then

$$f'J'\langle \alpha, \beta, m \rangle = J'\langle \gamma, \delta, n \rangle < J'\langle f'\alpha, f'\beta, m \rangle$$

hence

$$\langle \gamma, \delta, n \rangle S \langle f'\alpha, f'\beta, m \rangle.$$

Therefore

$$\gamma \leq \max(f'\alpha, f'\beta) \wedge \delta \leq \max(f'\alpha, f'\beta).$$

Since $\alpha \in b \wedge \beta \in b$,

$$f'\alpha \in \eta \wedge f'\beta \in \eta$$

i.e.

$$\max(f'\alpha, f'\beta) < \eta.$$

Then $\gamma \in \eta \wedge \delta \in \eta$, and consequently

$$(\exists \gamma_0 \in b) (\exists \delta_0 \in b) [\gamma = f'\gamma_0 \wedge \delta = f'\delta_0].$$

Since

$$\begin{aligned} \langle \gamma, \delta, n \rangle S \langle f'\alpha, f'\beta, m \rangle, \\ \langle f'\gamma_0, f'\delta_0, n \rangle S \langle f'\alpha, f'\beta, m \rangle \end{aligned}$$

and hence

$$\langle \gamma_0, \delta_0, n \rangle S \langle \alpha, \beta, m \rangle.$$

Again from the defining property of $\langle \alpha, \beta, m \rangle$

$$f'J'\langle \gamma_0, \delta_0, n \rangle = J'\langle f'\gamma_0, f'\delta_0, n \rangle = J'\langle \gamma, \delta, n \rangle = f'J'\langle \alpha, \beta, m \rangle.$$

Since f and J are one-to-one we conclude that

$$\langle \gamma_0, \delta_0, n \rangle = \langle \alpha, \beta, m \rangle$$

which is a contradiction.

2) If $\gamma \in \eta \wedge \delta \in \eta$ then

$$(\exists \gamma_0 \in b) (\exists \delta_0 \in b) [\gamma = f'\gamma_0 \wedge \delta = f'\delta_0].$$

Therefore from 1)

$$J^{\ast}\langle\gamma, \delta, m\rangle = J^{\ast}\langle f^{\ast}\gamma_0, f^{\ast}\delta_0, m\rangle = f^{\ast}J^{\ast}\langle\gamma_0, \delta_0, m\rangle.$$

Since b is closed w.r.t. J and f maps b into η ,

$$J^{\ast}\langle\gamma, \delta, m\rangle \in \eta$$

i.e. η is closed w.r.t. J .

Theorem 15.41. *If $\mathfrak{Q} \subseteq b \subseteq \text{On} \wedge \mathfrak{Q} \subseteq c \subseteq \text{On} \wedge K_1^{\ast}b \subseteq b \wedge K_2^{\ast}b \subseteq b \wedge J^{\ast}b^3 \subseteq b \wedge K_1^{\ast}c \subseteq c \wedge K_2^{\ast}c \subseteq c \wedge J^{\ast}c^3 \subseteq c \wedge f \text{ Isom}_{E,E}(b, c)$ then*

$$1) \quad \alpha \in b \wedge \beta \in b \wedge m < \mathfrak{Q} \rightarrow J^{\ast}\langle f^{\ast}\alpha, f^{\ast}\beta, m\rangle = f^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle$$

and

$$2) \quad \alpha \in b \rightarrow K_1^{\ast}f^{\ast}\alpha = f^{\ast}K_1^{\ast}\alpha \wedge K_2^{\ast}f^{\ast}\alpha = f^{\ast}K_2^{\ast}\alpha \wedge K_3^{\ast}\alpha = K_3^{\ast}f^{\ast}\alpha.$$

Proof. 1) Since $f \text{ Isom}_{E,E}(b, c)$ it follows that for some η, f_1 , and f_2 we have

$$f_1 \text{ Isom}_{E,E}(b, \eta) \wedge f_2 \text{ Isom}_{E,E}(c, \eta) \wedge f_2 \circ f = f_1.$$

Since b is closed w.r.t. J

$$\alpha \in b \wedge \beta \in b \wedge m < \mathfrak{Q} \rightarrow J^{\ast}\langle\alpha, \beta, m\rangle \in b$$

and hence

$$f^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle \in c.$$

If

$$\gamma = K_1^{\ast}f^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle \wedge \delta = K_2^{\ast}f^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle \wedge n = K_3^{\ast}f^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle$$

then since c is closed w.r.t. K_1 , and K_2

$$\gamma \in c \wedge \delta \in c \wedge n < \mathfrak{Q}$$

and

$$f^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle = J^{\ast}\langle\gamma, \delta, n\rangle.$$

From Theorem 15.40 we then have that

$$f_1^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle = (f_2 \circ f)^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle = f_2^{\ast}J^{\ast}\langle\gamma, \delta, n\rangle = J^{\ast}\langle f_2^{\ast}\gamma, f_2^{\ast}\delta, n\rangle.$$

On the other hand we also have from Theorem 15.40 that

$$f_1^{\ast}J^{\ast}\langle\alpha, \beta, m\rangle = J^{\ast}\langle f_1^{\ast}\alpha, f_1^{\ast}\beta, m\rangle.$$

Therefore since J is one-to-one

$$f_1^{\ast}\alpha = f_2^{\ast}\gamma \wedge f_1^{\ast}\beta = f_2^{\ast}\delta \wedge m = n$$

that is

$$\gamma = f^{\ast}\alpha \wedge \delta = f^{\ast}\beta \wedge m = n.$$

Therefore

$$f' J' \langle \alpha, \beta, m \rangle = J' \langle \gamma, \delta, n \rangle = J' \langle f' \alpha, f' \beta, m \rangle .$$

2) Since b is closed w.r.t. K_1 and K_2 we have that

$$\alpha \in b \rightarrow K'_1 \alpha \in b \wedge K'_2 \alpha \in b .$$

Since

$$\alpha = J' \langle K'_1 \alpha, K'_2 \alpha, K'_3 \alpha \rangle$$

we have from 1)

$$f' \alpha = f' J' \langle K'_1 \alpha, K'_2 \alpha, K'_3 \alpha \rangle = J' \langle f' K'_1 \alpha, f' K'_2 \alpha, K'_3 \alpha \rangle .$$

Therefore

$$K'_1 f' \alpha = f' K'_1 \alpha \wedge K'_2 f' \alpha = f' K'_2 \alpha \wedge K'_3 f' \alpha = K'_3 \alpha .$$

Remark. For our next theorem we need several results that we prove as lemmas.

Lemma 1. *If $\mathcal{O} \subseteq b \subseteq \text{On}$ and b is closed w.r.t. J then $F''b$ is closed w.r.t. the fundamental operations.*

Proof. If $x \in F''b \wedge y \in F''b$ then

$$(\exists \alpha \in b) (\exists \beta \in b) [x = F' \alpha \wedge y = F' \beta]$$

and hence

$$\mathcal{F}_n(x, y) = \mathcal{F}_n(F' \alpha, F' \beta) = F' J' \langle \alpha, \beta, n \rangle .$$

Since b is closed w.r.t. J it follows that

$$J' \langle \alpha, \beta, n \rangle \in b$$

and hence

$$\mathcal{F}_n(x, y) \in F''b .$$

Lemma 2. *If $\mathcal{O} \subseteq b \subseteq \text{On}$, if b is closed w.r.t. C and J , and if $x \in F''b$ then*

$$\text{Od}' x \in b .$$

Proof. From Lemma 1

$$\begin{aligned} x \in F''b &\rightarrow \{x\} \in F''b \\ &\rightarrow (\exists \alpha \in b) [\{x\} = F' \alpha] . \end{aligned}$$

But

$$\text{Od}' x = \text{Od}' \text{As}' \{x\} = \text{Od}' \text{As}' F' \alpha = C' \alpha .$$

Since b is closed w.r.t. C and $\alpha \in b$ we have that

$$C' \alpha \in b$$

i.e.

$$\text{Od}'x \in b .$$

Lemma 3. *If $9 \subseteq b \subseteq \text{On}$ and b is closed w.r.t. C then*

$$x \in F''b \wedge x \neq 0 \rightarrow x \cap F''b \neq 0 .$$

Proof. $x \in F''b \rightarrow (\exists \alpha \in b) [x = F'\alpha]$.

Since b is closed w.r.t. C

$$\begin{aligned} \alpha \in b &\rightarrow C'\alpha \in b \\ &\rightarrow F'C'\alpha \in F''b . \end{aligned}$$

But

$$F'C'\alpha = F'\text{Od}'\text{As}'F'\alpha = \text{As}'x .$$

If $x \neq 0$ then

$$\text{As}'x \in x$$

i.e.

$$F'C'\alpha \in x \cap F''b .$$

Lemma 4. *If $9 \subseteq b \subseteq \text{On}$ and b is closed w.r.t. C and J then*

- 1) $\{x, y\} \in F''b \rightarrow x \in F''b \wedge y \in F''b ,$
- 2) $\langle x, y \rangle \in F''b \rightarrow x \in F''b \wedge y \in F''b ,$
- 3) $\langle x, y, z \rangle \in F''b \rightarrow x \in F''b \wedge y \in F''b \wedge z \in F''b .$

Proof. 1) Since $\{x, y\} \neq 0$ we have from Lemma 3 that

$$\{x, y\} \cap F''b \neq 0 .$$

Therefore

$$x \in F''b \vee y \in F''b .$$

If $x \in F''b \wedge x \neq y$ then from Lemma 1, $\{x\} \in F''b$ and

$$\{y\} = \{x, y\} - \{x\} \in F''b .$$

'Since $\{y\} \in F''b \wedge \{y\} \neq 0$ it follows from Lemma 3 that $y \in F''b$.

Similarly if $y \in F''b$ then $x \in F''b$.

2)–3) The proofs are left to the reader.

Lemma 5. *If $9 \subseteq b \subseteq \text{On}$ and b is closed w.r.t. C and J then*

$$\langle x, y \rangle \in Q_n \wedge y \in F''b \rightarrow x \in F''b , \quad n = 4, 6, 7, 8 .$$

Proof. $\langle x, y \rangle \in Q_4 \rightarrow (\exists z) [y = \langle x, z \rangle]$. Then by Lemma 4

$$y \in F''b \rightarrow x \in F''b .$$

$$\langle x, y \rangle \in Q_6 \rightarrow (\exists z) (\exists w) [x = \langle z, w \rangle \wedge y = \langle w, z \rangle] .$$

Then by Lemmas 1 and 4

$$y \in F''b \rightarrow x \in F''b .$$

$$\langle x, y \rangle \in Q_7 \rightarrow (\exists z) (\exists w) (\exists u) [x = \langle u, w, z \rangle \wedge y = \langle w, z, u \rangle] .$$

$$\langle x, y \rangle \in Q_8 \rightarrow (\exists z) (\exists w) (\exists u) [x = \langle u, w, z \rangle \wedge y = \langle u, z, w \rangle] .$$

In each case we see from Lemmas 1 and 4 that

$$y \in F''b \rightarrow x \in F''b .$$

Lemma 6. *If $\mathcal{Q} \subseteq b \subseteq \text{On}$ and if b is closed w.r.t. C and J then*

$$x \in F''(b \cap \eta) \wedge y \in x \cap F''b \rightarrow y \in F''(b \cap \eta) .$$

Proof. $x \in F''(b \cap \eta) \rightarrow x \in F''b \wedge x \in F''\eta$. Therefore

$$\text{Od}'x \in \eta$$

and by Lemma 2

$$\text{Od}'x \in b .$$

Also

$$\begin{aligned} y \in x \cap F''b &\rightarrow y \in x \wedge y \in F''b \\ &\rightarrow \text{Od}'y < \text{Od}'x \wedge \text{Od}'y \in b \\ &\rightarrow \text{Od}'y \in b \cap \eta \\ &\rightarrow y \in F''(b \cap \eta) . \end{aligned}$$

Lemma 7. *If $\mathcal{Q} \subseteq b \subseteq \text{On}$ and if b is closed w.r.t. C and J then*

$$x \in F'\eta \cap F''b \rightarrow x \in F''(b \cap \eta) .$$

Proof. $x \in F'\eta \rightarrow \text{Od}'x < \text{Od}'F'\eta \leq \eta$. By Lemma 2

$$x \in F''b \rightarrow \text{Od}'x \in b .$$

Then

$$x \in F''(b \cap \eta) .$$

Lemma 8. *If $\mathcal{Q} \subseteq b \subseteq \text{On}$ and b is closed w.r.t. C and J then*

- 1) $\{x, y\} \in F''(b \cap \eta) \rightarrow x \in F''(b \cap \eta) \wedge y \in F''(b \cap \eta)$.
- 2) $\langle x, y \rangle \in F''(b \cap \eta) \rightarrow x \in F''(b \cap \eta) \wedge y \in F''(b \cap \eta)$.
- 3) $\langle x, y, z \rangle \in F''(b \cap \eta) \rightarrow x \in F''(b \cap \eta) \wedge y \in F''(b \cap \eta) \wedge z \in F''(b \cap \eta)$.

Proof. 1) Since $F''(b \cap \eta) \subseteq F''b$, we have by Lemma 4.

$$\begin{aligned} \{x, y\} \in F''(b \cap \eta) &\rightarrow \{x, y\} \in F''b \\ &\rightarrow x \in F''b \wedge y \in F''b \end{aligned}$$

then

$$\{x, y\} \in F''(b \cap \eta) \wedge x \in \{x, y\} \cap F''b \wedge y \in \{x, y\} \cap F''b.$$

From Lemma 6

$$x \in F''(b \cap \eta) \wedge y \in F''(b \cap \eta).$$

2)–3) The proofs are left to the reader.

Lemma 9. *If $\emptyset \subseteq b \subseteq \text{On} \wedge \emptyset \subseteq c \subseteq \text{On}$, if b and c are each closed with respect to C and J , if $f \text{ Isom}_{E,E}(b, c)$ and if $H \text{ Isom}_{E,E}(F''(b \cap \eta), F''f''(b \cap \eta))$ for $\eta \in b$ then*

- 1) $\{x, y\} \in F''(b \cap \eta) \rightarrow H'\{x, y\} = \{H'x, H'y\},$
- 2) $\langle x, y \rangle \in F''(b \cap \eta) \rightarrow H'\langle x, y \rangle = \langle H'x, H'y \rangle,$
- 3) $\langle x, y, z \rangle \in F''(b \cap \eta) \rightarrow H'\langle x, y, z \rangle = \langle H'x, H'y, H'z \rangle,$
- 4) $(\forall x, y \in F''(b \cap \eta)) [\langle xy \rangle \in Q_n \leftrightarrow \langle H'x, H'y \rangle \in Q_n], n = 4, 5, 6, 7, 8.$

Proof. 1) From Lemma 8

$$\{x, y\} \in F''(b \cap \eta) \rightarrow x, y \in F''(b \cap \eta).$$

Therefore since $H \text{ Isom}_{E,E}(F''(b \cap \eta), F''f''(b \cap \eta))$ and since $x, y \in \{x, y\}$

$$H'x, H'y \in H'\{x, y\}$$

i.e.

$$\{H'x, H'y\} \subseteq H'\{x, y\}.$$

Either $H'\{x, y\} = \{H'x, H'y\}$ or $\exists z \in H'\{x, y\} - \{H'x, H'y\}$. In the latter case we note that $\eta \in b$ and hence

$$f''(b \cap \eta) = c \cap f'\eta.$$

Since $H'\{x, y\}, H'x, H'y \in F''c$ it follows from Lemma 1 that

$$H'\{x, y\} - \{H'x, H'y\} \in F''c.$$

From Lemma 3

$$\exists z \in (H'\{x, y\} - \{H'x, H'y\}) \cap F''c.$$

Therefore

$$H'\{x, y\} \in F''(c \cap f'\eta) \wedge z \in H'\{x, y\} \cap F''c$$

and hence by Lemma 6

$$z \in F''(c \cap f'\eta) = F''f''(b \cap \eta).$$

Consequently $(\exists w \in F''(b \cap \eta)) [z = H'w]$. But since $z \in H'\{x, y\}$ and $H \text{ Isom}_{E,E}(F''(b \cap \eta), F''f''(b \cap \eta))$,

$$w \in \{x, y\}$$

i.e. $z = H'x \vee z = H'y$. This is a contradiction. Hence

$$H'\{x, y\} = \{H'x, H'y\}.$$

2) From Lemma 8

$$\langle x, y \rangle \in F''(b \cap \eta) \rightarrow x, y, \{x\}, \{x, y\} \in F''(b \cap \eta).$$

Therefore from 1) above

$$\begin{aligned} H'\langle x, y \rangle &= H'\{\{x\}, \{x, y\}\} = \{H'\{x\}, H'\{x, y\}\} = \{\{H'x\}, \{H'x, H'y\}\} \\ &= \langle H'x, H'y \rangle. \end{aligned}$$

3) The proof is left to the reader.

4) $\langle x, y \rangle \in Q_4 \rightarrow (\exists z) [y = \langle x, z \rangle]$.

But from Lemma 8

$$y \in F''(b \cap \eta) \rightarrow x, z \in F''(b \cap \eta),$$

Then from 3) above

$$\begin{aligned} y = \langle x, z \rangle &\rightarrow H'y = \langle H'x, H'z \rangle \\ &\rightarrow \langle H'x, H'y \rangle \in Q_4. \end{aligned}$$

Conversely

$$\langle H'x, H'y \rangle \in Q_4 \rightarrow (\exists w) [H'y = \langle H'x, w \rangle].$$

Since $y \in F''(b \cap \eta)$, $H'y \in F''(c \cap f'\eta)$. Hence by Lemma 8, $w \in F''(c \cap f'\eta) = F''f''(b \cap \eta)$. Consequently $(\exists z \in F''(b \cap \eta)) [w = H'z]$.

Since $f^{-1} \text{Isom}_{E,E}(c, b) \wedge H^{-1} \text{Isom}_{E,E}(F''(c \cap f'\eta), F''f^{-1}(c \cap f'\eta))$ the foregoing argument gives

$$\begin{aligned} H'y = \langle H'x, H'z \rangle &\rightarrow y = \langle x, z \rangle \\ &\rightarrow \langle x, y \rangle \in Q_4. \end{aligned}$$

The arguments for Q_5, Q_6, Q_7 , and Q_8 are similar and are left to the reader.

Theorem 15.42. 1) If $9 \subseteq b \subseteq \text{On} \wedge 9 \subseteq c \subseteq \text{On}$, if b and c are each closed with respect to K_1, K_2, C , and J and if $f \text{Isom}_{E,E}(b, c)$ then

$$(\forall \alpha, \beta \in b) [[F'\alpha \in F'\beta \leftrightarrow F'f'\alpha \in F'f'\beta] \wedge [F'\alpha = F'\beta \leftrightarrow F'f'\alpha = F'f'\beta]].$$

2) If $\omega + 2 \subseteq b \subseteq \text{On} \wedge \omega + 2 \subseteq c \subseteq \text{On}$, if b and c are each closed with respect to K_1, K_2, C_a and J and if $f \text{Isom}_{E,E}(b, c)$ then

$$\begin{aligned} (\forall \alpha, \beta \in b) &[[F'_a\alpha \in F'_a\beta \leftrightarrow F'_af'\alpha \in F'_af'\beta] \\ &\wedge [F'_a\alpha = F'_a\beta \leftrightarrow F'_af'\alpha = F'_af'\beta]]. \end{aligned}$$

Proof. 1) (By induction on $\max(\alpha, \beta)$.) If $\eta = \max(\alpha, \beta)$ and if $\eta = \alpha = \beta$ then the result is true because

$$F'\alpha = F'\beta \wedge F'f'\alpha = F'f'\beta \wedge F'\alpha \notin F'\beta \wedge F'f'\alpha \notin F'f'\beta.$$

If $\alpha < \beta = \eta \vee \beta < \alpha = \eta$ then it is sufficient to prove that if $\gamma \in b \cap \eta$.

- i) $F'\gamma \in F'\eta \iff F'f'\gamma \in F'f'\eta,$
- ii) $F'\eta \in F'\gamma \iff F'f'\eta \in F'f'\gamma,$
- iii) $F'\gamma = F'\eta \iff F'f'\gamma = F'f'\eta.$

If $H = \{\langle F'\gamma, F'f'\gamma \rangle \mid \gamma \in b \cap \eta\}$ then by the induction hypothesis

$$(\forall \gamma, \delta \in b \cap \eta) [[F'\gamma \in F'\delta \iff F'f'\gamma \in F'f'\delta] \\ \wedge [F'\gamma = F'\delta \iff F'f'\gamma = F'f'\delta]]$$

consequently $H \text{ Isom}_{E,E}(F''(b \cap \eta), F''f''(b \cap \eta))$. With the aid of Lemma 9 we will prove i) i.e.

$$(\forall \gamma \in b \cap \eta) [F'\gamma \in F'\eta \iff F'f'\gamma \in F'f'\eta].$$

We argue by cases.

If $K'_3\eta = 0$ then $K'_3f'\eta = 0$ (Theorem 15.41). Therefore $F'\eta = F''\eta$ and $F''f'\eta = F''f'\eta$. Consequently

$$(\forall \gamma \in b \cap \eta) [F'\gamma \in F'\eta].$$

But since f is E-order preserving

$$\gamma \in b \cap \eta \Rightarrow f'\gamma \in f'\eta.$$

Hence

$$(\forall \gamma \in b \cap \eta) [F'f'\gamma \in F'f'\eta].$$

If $K'_3\eta \neq 0$ then $K'_1\eta < \eta \wedge K'_2\eta < \eta$. Since $\eta \in b$ and b is closed with respect to K_1 and K_2 ,

$$K'_1\eta \in b \cap \eta \wedge K'_2\eta \in b \cap \eta.$$

From Theorem 15.41

$$K'_1f'\eta = f'K'_1\eta, \quad K'_2f'\eta = f'K'_2\eta, \quad K'_3f'\eta = K'_3\eta.$$

If $K'_3\eta = 1$

$$F'\eta = \{F'K'_1\eta, F'K'_2\eta\} \wedge F'f'\eta = \{F'f'K'_1\eta, F'f'K'_2\eta\}.$$

Then from the induction hypothesis if $\gamma \in b \cap \eta$

$$F'\gamma \in F'\eta \iff F'\gamma = F'K'_1\eta \vee F'\gamma = F'K'_2\eta \\ \iff F'f'\gamma = F'f'K'_1\eta \vee F'f'\gamma = F'f'K'_2\eta \\ \iff F'f'\gamma \in F'f'\eta.$$

If $K'_3\eta = 2$

$$F'\eta = E \cap F'K'_1\eta \wedge F'f'\eta = E \cap F'f'K'_1\eta.$$

Again from the induction hypothesis if $\gamma \in b \cap \eta$

$$\begin{aligned} F'\gamma \in F'\eta &\iff F'\gamma \in E \wedge F'\gamma \in F'K'_1\eta \\ &\iff (\exists x, y) [F'\gamma = \langle x, y \rangle \wedge x \in y] \wedge F'f'\gamma \in F'f'K'_1\eta. \end{aligned}$$

From Lemma 8

$$\langle x, y \rangle = F'\gamma \in F''(b \cap \eta) \rightarrow x \in F''(b \cap \eta) \wedge y \in F''(b \cap \eta).$$

Then from Lemma 9

$$F'f'\gamma = H'F'\gamma = H'\langle x, y \rangle = \langle H'x, H'y \rangle.$$

Furthermore

$$x \in y \rightarrow H'x \in H'y.$$

Thus

$$F'f'\gamma \in E \cap F''f'K'_1\eta$$

i.e.

$$(\forall \gamma \in b \cap \eta) [F'\gamma \in F'\eta \rightarrow F'f'\gamma \in F'f'\eta].$$

If $K'_3\eta = 3$

$$F'\eta = F'K'_1\eta - F'K'_2\eta \wedge F'f'\eta = F'f'K'_1\eta - F'f'K'_2\eta.$$

From the induction hypothesis if $\gamma \in b \cap \eta$

$$\begin{aligned} F'\gamma \in F'\eta &\iff F'\gamma \in F'K'_1\eta \wedge F'\gamma \notin F'K'_2\eta \\ &\iff F'f'\gamma \in F'f'K'_1\eta \wedge F'f'\gamma \notin F'f'K'_2\eta \\ &\iff F'f'\gamma \in F'f'\eta. \end{aligned}$$

If $K'_3\eta = n, n = 4, 6, 7, 8$

$$F'\eta = F'K'_1\eta \cap Q_n''F'K'_2\eta \wedge F'f'\eta = F'f'K'_1\eta \cap Q_n''F'f'K'_2\eta.$$

Then if $\gamma \in b \cap \eta$

$$\begin{aligned} F'\gamma \in F'\eta &\iff F'\gamma \in F'K'_1\eta \wedge F'\gamma \in Q_n''F'K'_2\eta \\ &\iff F'f'\gamma \in F'f'K'_1\eta \wedge (\exists x \in F'K'_2\eta) [\langle x, F'\gamma \rangle \in Q_n]. \end{aligned}$$

But from Lemma 5

$$F'\gamma \in F''b \wedge \langle x, F'\gamma \rangle \in Q_n \rightarrow x \in F''b.$$

Then

$$F'K'_2\eta \in F''(b \cap \eta) \wedge x \in F'K'_2\eta \cap F''b.$$

Therefore by Lemma 6, $x \in F''(b \cap \eta)$. Thus

$$x \in F''(b \cap \eta) \wedge F'\gamma \in F''(b \cap \eta) \wedge \langle x, F'\gamma \rangle \in Q_n.$$

From Lemma 9

$$\langle H'x, H'F'\gamma \rangle \in Q_n.$$

But

$$H'F'\gamma = F'f'\gamma.$$

Therefore since $x \in F'K'_2\eta$

$$H'x \in H'F'K'_2\eta = F'f'K'_2\eta$$

i.e.

$$F'f'\gamma \in Q_n F'f'K'_2\eta.$$

Thus

$$(\forall \gamma \in b \cap \eta) [F'\gamma \in F'\eta \rightarrow F'f'\gamma \in F'f'\eta].$$

If $K'_3\eta = 5$

$$F'\eta = F'K'_1\eta \cap Q_5 F'K'_2\eta \wedge F'f'\eta = F'f'K'_1\eta \cap Q_5 F'f'K'_2\eta.$$

Thus if $\gamma \in b \cap \eta$ then since $Q_5 = Q_4^{-1}$

$$\begin{aligned} F'\gamma \in F'\eta &\iff F'\gamma \in F'K'_1\eta \wedge F'\gamma \in Q_5 F'K'_2\eta \\ &\iff F'f'\gamma \in F'f'K'_1\eta \wedge (\exists x \in F'K'_2\eta) [\langle F'\gamma, x \rangle \in Q_4]. \end{aligned}$$

Then

$$F'K'_2\eta \cap Q_4 \{F'\gamma\} \neq 0.$$

Furthermore since $F'\gamma \in F''b \wedge F'K'_2\eta \in F''b$ we have from Lemma 1

$$F'K'_2\eta \cap Q_4 \{F'\gamma\} \in F''b.$$

Therefore by Lemma 3

$$(\exists y \in F''b) [y \in F'K'_2\eta \cap Q_4 \{F'\gamma\}].$$

Thus

$$y \in F''b \wedge \langle F'\gamma, y \rangle \in Q_4.$$

Then

$$F'K'_2\eta \in F''(b \cap \eta) \wedge y \in F'K'_2\eta \cap F''b$$

and hence by Lemma 6

$$y \in F''(b \cap \eta).$$

Then since $\langle F'\gamma, y \rangle \in Q_4$ we have from Lemma 9

$$\langle H'F'\gamma, H'y \rangle \in Q_4,$$

$$\langle H'y, H'F'\gamma \rangle \in Q_5.$$

But $H'F'\gamma = F'f'\gamma$. Therefore since $y \in F'K'_2\eta$

$$H'y \in H'F'K'_2\eta = F'f'K'_2\eta$$

i.e.

$$F'f'\gamma \in Q_5'' F'f'K_2\eta.$$

Thus

$$(\forall \gamma \in b \cap \eta) [F'\gamma \in F'\eta \rightarrow F'f'\gamma \in F'f'\eta].$$

Having exhausted all cases we have proved

$$(\forall \gamma \in b \cap \eta) [F'\gamma \in F'\eta \rightarrow F'f'\gamma \in F'f'\eta].$$

The implication in the reverse direction follows from symmetry i.e.

$$f^{-1} \text{Isom}_{E,E}(c, b) \wedge H^{-1} \text{Isom}_{E,E}(F''(c \cap f'\eta), F''f^{-1}(c \cap f'\eta)).$$

Therefore, since

$$\begin{aligned} \gamma \in b \cap \eta &\rightarrow f'\gamma \in c \cap f'\eta, \\ F'f'\gamma \in F'f'\eta &\rightarrow F'\gamma \in F'\eta. \end{aligned}$$

This completes the proof of i).

From i) we next prove iii). If $F'\eta \neq F'\gamma$ then $F'\eta - F'\gamma \neq 0$ or $F'\gamma - F'\eta \neq 0$. Since $F'\gamma \in F''b \wedge F'\eta \in F''b$ we have from Lemma 1

$$F'\eta - F'\gamma \in F''b \wedge F'\gamma - F'\eta \in F''b.$$

If $F'\eta - F'\gamma \neq 0$ then by Lemma 3

$$(\exists x \in F''b) [x \in F'\eta - F'\gamma].$$

Thus

$$x \in F'\eta \cap F''b$$

and hence by Lemma 7

$$x \in F''(b \cap \eta)$$

i.e.

$$(\exists v \in b \cap \eta) [x = F'v].$$

But $x \in F'\eta - F'\gamma$. Therefore

$$F'v \in F'\eta \wedge F'v \notin F'\gamma.$$

From i)

$$F'f'v \in F'f'\eta \wedge F'f'v \notin F'f'\gamma.$$

Thus $F'f'v \in F'f'\eta - F'f'\gamma$ and hence $F'f'\gamma \neq F'f'\eta$.

If $F'\gamma - F'\eta \neq 0$ the argument is similar to the foregoing one and is left to the reader. We have then proved

$$(\forall \gamma \in b \cap \eta) [F'f'\eta = F'f'\gamma \rightarrow F'\eta = F'\gamma].$$

Again the implication in the reverse direction follows from symmetry.

We next prove ii) from i) and iii). If $F'\eta \in F'\gamma$ and if $v = \text{Od}'F'\eta$ then

$$v = \text{Od}'F'\eta < \text{Od}'F'\gamma \leq \gamma < \eta$$

i.e. $v < \eta$. Since $F'\eta \in F''b$ we have from Lemma 2

$$v = \text{Od}'F'\eta \in b.$$

Then $v \in b \cap \eta$ and hence by the induction hypothesis

$$F'v \in F'\gamma \rightarrow F'f'v \in F'f'\gamma.$$

But since $F'v = F'\eta$ we have from iii)

$$F'f'v = F'f'\eta.$$

Hence

$$F'\eta \in F'\gamma \rightarrow F'f'\eta \in F'f'\gamma.$$

Again the reverse implication follows from symmetry.

2) The proof is left to the reader.

Theorem 15.43.

1) If $\mathcal{Q} \subseteq b \subseteq \text{On}$, if b is closed with respect to C, K_1, K_2 , and J and if $f \text{ Isom}_{E,E}(b, \eta)$ then

$$(\forall \alpha, \beta \in b) [F'\alpha \in F'\beta \leftrightarrow F'f'\alpha \in F'f'\beta].$$

2) If $\mathcal{Q} \subseteq b \subseteq \text{On}$, if b is closed with respect to C_a, K_1, K_2 , and J and if $f \text{ Isom}_{E,E}(b, \eta)$ then

$$(\forall \alpha, \beta \in b) [F'_a\alpha \in F'_a\beta \leftrightarrow F'_af'\alpha \in F'_af'\beta].$$

Proof. 1) From Theorem 15.40, η is closed with respect to J . Since

$$K'_1\alpha \leq \alpha \wedge K'_2\alpha \leq \alpha \wedge C'\alpha \leq \alpha$$

η is also closed with respect to C, K_1 , and K_2 . Therefore by Theorem 15.42

$$(\forall \alpha, \beta \in b) [F'\alpha \in F'\beta \leftrightarrow F'f'\alpha \in F'f'\beta].$$

2) The proof is left to the reader.

Theorem 15.44.

1) $V = L \rightarrow \text{GCH}$.

2) $V = L_a \rightarrow \text{GCH}$.

Proof. Theorems 15.43, 15.39, and 15.38.

Theorem 15.45.

- 1) L is a model of the GCH.
- 2) L_a is a model of the GCH.

Proof. Theorems 15.44 and 15.30.

Remark. We have now shown how to select from V a subclass L that is a model of $ZF + AC + GCH + V = L$. This process can be relativized to any standard transitive model \mathcal{M} to produce a subclass of \mathcal{M} that is also a model of $ZF + AC + GCH + V = L$. Recall that

$$L^{\mathcal{M}} = \{x \in \mathcal{M} \mid (\exists \alpha \in \mathcal{M}) [x = F'\alpha]\} .$$

Theorem 15.46. *If \mathcal{M} is a standard transitive model of ZF then*

- 1) $(\forall \alpha \in \mathcal{M}) [F'\alpha \in \mathcal{M}]$,
- 2) $a \subseteq \omega \wedge a \in \mathcal{M} \rightarrow (\forall \alpha \in \mathcal{M}) [F'_a \alpha \in \mathcal{M}]$.

Proof. 1) (by induction). Since \mathcal{M} is a model of the Axiom Schema of Replacement it follows from the induction hypothesis that if $K'_3 \alpha = 0$ then

$$F'\alpha = F''\alpha = \{x \mid (\exists \beta \in \alpha) [x = F'\beta]\} = \{x \in \mathcal{M} \mid (\exists \beta \in \alpha) [x = F'\beta]^{\mathcal{M}}\} \in \mathcal{M} .$$

If $K'_3 \alpha = i \neq 0$ then $K'_1 \alpha < \alpha$, $K'_2 \alpha < \alpha$ and from the induction hypothesis and the fact that \mathcal{M} is closed under the eight fundamental operations

$$F'\alpha = \mathcal{F}_i(F'K'_1 \alpha, F'K'_2 \alpha) \in \mathcal{M} .$$

- 2) The proof is left to the reader.

Theorem 15.47. *If \mathcal{M} is a standard transitive model of ZF then*

- 1) $L^{\mathcal{M}} = \{x \mid (\exists \alpha \in \mathcal{M}) [x = F'\alpha]\}$,
- 2) $a \subseteq \omega \wedge a \in \mathcal{M} \rightarrow L_a^{\mathcal{M}} = \{x \mid (\exists \alpha \in \mathcal{M}) [x = F'_a \alpha]\}$.

Proof. Obvious from Theorem 15.46.

Remark. That $L^{\mathcal{M}}$ is a standard transitive model of $ZF + AC + GCH + V = L$ is immediate from the following theorem.

Theorem 15.48. *If \mathcal{M} is a standard transitive model of ZF and if φ is a wff of ZF then*

- 1) $(\varphi^L)^{\mathcal{M}} \longleftrightarrow \varphi^{L^{\mathcal{M}}}$
- 2) $(\varphi^{L_a})^{\mathcal{M}} \longleftrightarrow \varphi^{L_a^{\mathcal{M}}}$, if $a \in \mathcal{M}$.

Proof. 1) (By induction on the number of logical symbols in φ .) φ must be of the form 1) $a \in b$, 2) $\neg \psi$, 3) $\psi \wedge \eta$, or 4) $(\forall x) \psi$. The arguments

for cases 1)–3) we leave to the reader. If φ is of the form $(\forall x) \psi$ then as our induction hypothesis

$$(\psi^L)^{\mathcal{M}} \longleftrightarrow \psi^{L^{\mathcal{M}}}.$$

Then

$$\begin{aligned} [[(\forall x) \psi]^L]^{\mathcal{M}} &\longleftrightarrow [(\forall x) [x \in L \rightarrow \psi^L]]^{\mathcal{M}} \\ &\longleftrightarrow [(\forall x) [(\exists \alpha) [x = F'\alpha] \rightarrow \psi^L]]^{\mathcal{M}} \\ &\longleftrightarrow (\forall x \in \mathcal{M}) [(\exists \alpha \in \mathcal{M}) [x = F'\alpha] \rightarrow (\psi^L)^{\mathcal{M}}] \\ &\longleftrightarrow (\forall x) [x \in L^{\mathcal{M}} \rightarrow \psi^{L^{\mathcal{M}}}] \\ &\longleftrightarrow [(\forall x) \psi]^{L^{\mathcal{M}}}. \end{aligned}$$

2) The proof is left to the reader.

Theorem 15.49. *If \mathcal{M} is a standard transitive model of ZF then*

1) $L^{\mathcal{M}}$ is a standard transitive model of ZF + AC + GCH + $V = L$ and $\text{On}^{L^{\mathcal{M}}} = \text{On}^{\mathcal{M}}$.

2) $L_a^{\mathcal{M}}$ is a standard transitive model of ZF + AC + GCH + $V = L_a$ and $\text{On}^{L_a^{\mathcal{M}}} = \text{On}^{\mathcal{M}}$, if $a \in \mathcal{M}$.

Proof. 1) Since $0 \in \mathcal{M} \wedge F'0 = 0$ it follows that $0 \in L^{\mathcal{M}}$ and hence $L^{\mathcal{M}}$ is not empty. Furthermore if $y \in x \in L^{\mathcal{M}}$ then

$$(\exists \alpha \in \mathcal{M}) [y \in x = F'\alpha \subseteq F''\alpha].$$

Therefore $(\exists \beta < \alpha) [y = F'\beta]$. But $\alpha \in \mathcal{M}$ and \mathcal{M} is transitive. Thus $\beta \in \mathcal{M}$ and $y \in L^{\mathcal{M}}$, and hence $L^{\mathcal{M}}$ is transitive.

Since L is a model of ZF + AC + GCH + $V = L$ it follows that if φ is any axiom of ZF + AC + GCH + $V = L$ then φ^L is a theorem of ZF. Since \mathcal{M} is a model of ZF every theorem of ZF relativized to \mathcal{M} is a theorem of ZF. Therefore $(\varphi^L)^{\mathcal{M}}$ is a theorem of ZF. Then by Theorem 15.48 $\varphi^{L^{\mathcal{M}}}$ is a theorem of ZF. Hence $L^{\mathcal{M}}$ is a model of φ .

Consequently $L^{\mathcal{M}}$ is a standard transitive model of ZF + AC + GCH + $V = L$.

By Theorem 15.26 $\text{On}^L = \text{On}$. Relativizing to \mathcal{M} we have $\text{On}^{L^{\mathcal{M}}} = \text{On}^{\mathcal{M}}$. Details are left to the reader.

2) The proof is left to the reader.

16 The Arithmetization of Model Theory

In Section 12 we developed a theory of internal models and gave meaning to the idea of a class M being a standard transitive model of ZF. For \mathcal{M} to be a standard model of ZF means in particular that \mathcal{M} is a model of Axiom 5, the Axiom Schema of Replacement. Since Axiom 5 is a schema “ \mathcal{M} is a standard model of ZF” is a meta-statement asserting that a certain infinite collection of sentences of ZF hold. Can this meta-statement be formalized in ZF, that is, can “ \mathcal{M} is a standard model of ZF” be expressed as a single sentence in ZF? This question is as yet unresolved. It can be resolved however if \mathcal{M} is a set, that is, “ m is a standard model of ZF” can be expressed as a single sentence in ZF. The basic objective of this section is to produce such a sentence. Our approach is to assign Gödel numbers to the well formed formulas of our language. This assignment will be made by the mapping J of Definition 15.2.

For purposes that will become clear in the next section we choose to introduce Gödel numbers for a somewhat more general language than that of ZF. To this language we adjoin the axioms and rules of inference of ZF to obtain a slightly different formulation of Zermelo-Fraenkel theory that we will denote by $ZF\{\alpha_0\}$ where α_0 is a fixed ordinal. The primitive symbols of the language of $ZF\{\alpha_0\}$ are the following.

individual variables: $x_0, x_1, \dots, x_i, \dots i \in \omega,$

individual constants: $k_0, k_1, \dots, k_\alpha, \dots \alpha \in \alpha_0,$

a predicate constant: $\in,$

primitive logical symbols: $\neg, \wedge, \exists.$

The formation rules are the same as for ZF with the added requirement that if a and b are variables or individual constants then $a \in b$ is a well formed formula. $ZF\{\alpha_0\}$ is then an extension of ZF, indeed $ZF\{0\}$ is ZF.

With each wff φ of $ZF\{\alpha_0\}$ we associate an ordinal number ' φ ', called the Gödel number of φ . This number we define inductively in the following way.

Definition 16.1

- 1) $\ulcorner x_i \urcorner \triangleq J \langle i, 0, 0 \rangle, i \in \omega.$
- 2) $\ulcorner k_\alpha \urcorner \triangleq J \langle \alpha, 1, 1 \rangle, \alpha \in \alpha_0.$
- 3) $\ulcorner a \in b \urcorner \triangleq J \langle \ulcorner a \urcorner, \ulcorner b \urcorner, 2 \rangle, a$ and b variables or individual constants.
- 4) $\ulcorner (\exists x_i) \varphi \urcorner \triangleq J \langle \ulcorner x_i \urcorner, \ulcorner \varphi \urcorner, 3 \rangle.$
- 5) $\ulcorner \neg \varphi \urcorner \triangleq J \langle \ulcorner \varphi \urcorner, 4, 4 \rangle.$
- 6) $\ulcorner \varphi \wedge \eta \urcorner \triangleq J \langle \ulcorner \varphi \urcorner, \ulcorner \eta \urcorner, 5 \rangle.$

Remark. From Definition 16.1 it is clear that for each Gödel number α there is one and only one wff φ of $ZF \{\alpha_0\}$ for which

$$\alpha = \ulcorner \varphi \urcorner,$$

and for each φ there is one and only one Gödel number α such that

$$\ulcorner \varphi \urcorner = \alpha.$$

This one-to-one correspondence between wffs and Gödel numbers will be useful in the following way. In order to express Axiom 5, the Axiom Schema of Replacement, as a single sentence we must find some way of asserting a property of each wff. While we cannot quantify on wffs we can quantify on Gödel numbers. Using Gödel numbers we will code Axiom 5 into a single wff of the type

$$(\forall \alpha \in a) [\varphi(\alpha)],$$

where a is an appropriately chosen class of Gödel numbers and φ is a wff that asserts that the wff with Gödel number α is satisfied in the model.

Throughout this section it should be kept in mind that we will use the language ZF introduced in Section 2 to talk about the languages $ZF \{\alpha_0\}$. References to wffs will always be references to wff of $ZF \{\alpha_0\}$. The proofs of theorems about these wffs of $ZF \{\alpha_0\}$ will be carried out in the language ZF , or the meta-language.

We begin with a definition of “wff(α_0)” the class of Gödel numbers of wffs of $ZF \{\alpha_0\}$. We will frequently refer to wff(α_0) simply as the wffs of $ZF \{\alpha_0\}$. In defining “wff(α_0)” we will be guided by one particular use to which our theory will be put. Later we will be interested in conditions on $K \triangleq \{k_\alpha \mid \alpha \in \alpha_0\}$ that assure us that K is a standard transitive model of ZF . We will of course wish to say that K satisfies a wff $(\exists x_i) \varphi(x_i)$ iff $(\exists k_\alpha \in K) \varphi(k_\alpha)$. Here “ $\varphi(k_\alpha)$ ” is the wff obtained from $\varphi(x_i)$ by replacing each occurrence of x_i that is free for substitution, by k_α . We must therefore either define “occurrence of x_i that is free for substitution” or arrange our formulas so that in $(\exists x_i) \varphi(x_i)$ we do not have x_i occurring bound in $\varphi(x_i)$. We choose the second alternative.

Formulas such as

$$(\exists x_1) (\exists x_1) [x_1 = x_1],$$

are technically well formed by our formation rules. There is however no loss of generality if we exclude such formulas from consideration in defining “wff(α_0)”. We will therefore define “wff(α_0)” as the class of Gödel numbers of those wffs of ZF $\{\alpha_0\}$ that are built up from prime formulas by negation, conjunction and quantification taking care to see that we do not quantify twice on the same variable. This definition must be formulated in ZF. This will be done inductively using the class of Gödel numbers of variables “va”, the class of Gödel numbers of individual constants “ic(α_0)”, and the class of Gödel numbers of prime wffs “pf(α_0)”.

Throughout this chapter \mathcal{M} is a standard transitive model of ZF.

Theorem 16.2. $(\forall \alpha \in \mathcal{M}) (\forall \beta \in \mathcal{M}) (\forall n < 9) [J\langle \alpha, \beta, n \rangle \in \mathcal{M}]$.

Proof. Since $\text{On} \subseteq L$

$$(\forall \alpha) (\forall \beta) (\forall n < 9) [J\langle \alpha, \beta, n \rangle \in L]$$

i.e.

$$(\forall \alpha) (\forall \beta) (\forall n < 9) (\exists \gamma) (\exists \delta) [J\langle \alpha, \beta, n \rangle = \gamma \wedge F\delta = \gamma].$$

Since \mathcal{M} is a standard transitive model of ZF and $J\langle \alpha, \beta, n \rangle$ and $F\delta$ are each absolute with respect to \mathcal{M} we have

$$(\forall \alpha \in \mathcal{M}) (\forall \beta \in \mathcal{M}) (\forall n < 9) (\exists \gamma \in \mathcal{M}) (\exists \delta \in \mathcal{M}) [J\langle \alpha, \beta, n \rangle = \gamma \wedge F\delta = \gamma].$$

In particular

$$(\forall \alpha \in \mathcal{M}) (\forall \beta \in \mathcal{M}) (\forall n < 9) [J\langle \alpha, \beta, n \rangle \in \mathcal{M}].$$

Definition 16.3.

- 1) $\text{va} = \{ \ulcorner x_i \urcorner \mid i \in \omega \}$.
- 2) $\text{ic}(\alpha_0) = \{ \ulcorner k_\alpha \urcorner \mid \alpha \in \alpha_0 \}$.
- 3) $\text{vc}(\alpha_0) = \text{va} \cup \text{ic}(\alpha_0)$.

Theorem 16.4.

- 1) $\text{va} \subseteq \omega \wedge \text{va} \in \mathcal{M}$.
- 2) $[x \in \text{va}] \text{Abs } \mathcal{M}$.

Proof.

1) $\alpha < \omega \rightarrow J\langle \alpha, 0, 0 \rangle < \omega$. Hence $\text{va} \subseteq \omega$. Since \mathcal{M} is a model of the Axiom Schema of Replacement, and $\omega \in \mathcal{M}$

$$\text{va} = \{ \alpha \mid (\exists \beta \in \omega) [\alpha = J\langle \beta, 0, 0 \rangle] \} \in \mathcal{M}.$$

- 2) $x \in \text{va} \leftrightarrow (\exists \alpha) [\alpha \in \omega \wedge x = J\langle \alpha, 0, 0 \rangle]$.

Theorem 16.5.

- 1) $\text{ic}(\alpha_0) \subseteq \mathcal{M} \wedge \text{vc}(\alpha_0) \subseteq \mathcal{M}$, if $\alpha_0 \subseteq \mathcal{M}$.
- 2) $[x \in \text{ic}(\alpha_0)] \text{ Abs } \mathcal{M}$.
- 3) $[x \in \text{vc}(\alpha_0)] \text{ Abs } \mathcal{M}$.

Proof.

- 1) If $\alpha \in \alpha_0 \subseteq \mathcal{M}$ then since \mathcal{M} is closed with respect to J , $J' \langle \alpha, 1, 1 \rangle \in \mathcal{M}$. Hence $\text{ic}(\alpha_0) \subseteq \mathcal{M}$ and $\text{vc}(\alpha_0) = [\text{va} \cup \text{ic}(\alpha_0)] \subseteq \mathcal{M}$.
- 2) $x \in \text{ic}(\alpha_0) \longleftrightarrow (\exists \alpha) [\alpha \in \alpha_0 \wedge x = J' \langle \alpha, 1, 1 \rangle]$.
- 3) $\text{vc}(\alpha_0) = \text{va} \cup \text{ic}(\alpha_0)$.

Definition 16.6. $\text{pf}(\alpha_0) = \{ 'a \in b' \mid 'a', 'b' \in \text{vc}(\alpha_0) \}$.

Remark. The class $\text{pf}(\alpha_0)$ is the class of (Gödel numbers of) prime formulas of ZF $\{\alpha_0\}$. In Definition 16.6 we have deliberately abused the language in an effort to make clear the intended meaning. We will continue this practice and rely upon the reader to convince himself that such statements can be formalized in ZF. For example, Definition 16.6 should be

$$\text{pf}(\alpha_0) = \{ \gamma \mid (\exists \alpha, \beta \in \text{vc}(\alpha_0)) [\gamma = J' \langle \alpha, \beta, 2 \rangle] \}.$$

Theorem 16.7.

- 1) $\text{pf}(\alpha_0) \subseteq \mathcal{M}$, if $\alpha_0 \subseteq \mathcal{M}$.
- 2) $[x \in \text{pf}(\alpha_0)] \text{ Abs } \mathcal{M}$.

The proof is left to the reader.

Remark. In this section we will also derive certain results that will be needed in the next section where we take up Cohen's work. One idea that is basic for Cohen's results is the notion of a limited quantifier, " \exists^β ":

Definition 16.8.

$$(\exists^\beta x_i) \varphi(x_i) \stackrel{\Delta}{\longleftrightarrow} (\exists x_i) (\exists \alpha < \beta) [x_i = k_\alpha \wedge \varphi(k_\alpha)].$$

Remark. Although formulas that contain limited quantifiers are abbreviations for wffs that have been assigned Gödel numbers we find it convenient to introduce special Gödel numbers for such abbreviations. This means of course that certain formulas will have two Gödel numbers.

Definition 16.9. $'(\exists^\beta x_i) \varphi(x_i)' = J' \langle \beta, J' \langle 'x_i', ' \varphi ', 3 \rangle, 6 \rangle$.

Remark. For convenience in the discussions ahead we introduce the following more compact notation for the Gödel numbering function of Definition 16.9 and for the last three functions of Definition 16.1.

Definition 16.10.

- 1) $R_3' \langle \alpha, \beta \rangle \triangleq J' \langle \alpha, \beta, 3 \rangle,$
- 2) $R_4' \alpha \triangleq J' \langle \alpha, 4, 4 \rangle,$
- 3) $R_5' \langle \alpha, \beta \rangle \triangleq J' \langle \alpha, \beta, 5 \rangle,$
- 4) $R_6' \langle \beta, \gamma, \delta \rangle \triangleq J' \langle \beta, J' \langle \gamma, \delta, 3 \rangle, 6 \rangle.$

Definition 16.11. $G'b \triangleq b \cup R_3''(va \times b) \cup R_4''b \cup R_5''b^2,$

$$\begin{aligned} f'0 &\triangleq \text{pf}(\alpha_0), \\ f'(i+1) &\triangleq G'f'i, i \in \omega, \\ \text{qf}(\alpha_0) &\triangleq \bigcup_{i \in \omega} f'i. \end{aligned}$$

Definition 16.12. $G'b \triangleq b \cup R_4''b \cup R_5''b^2 \cup R_6''(\alpha_0 \times va \times b).$

$$\begin{aligned} f'0 &\triangleq \text{pf}(\alpha_0), \\ f(i+1) &\triangleq G'f'i, i \in \omega, \\ \text{qlf}(\alpha_0) &\triangleq \bigcup_{i \in \omega} f'i. \end{aligned}$$

Remark. We call $\text{qf}(\alpha_0)$ the class of quasi formulas and $\text{qlf}(\alpha_0)$ the class of quasi limited formulas. As we will prove below $\text{qf}(\alpha_0)$ is the class of Gödel numbers of the wffs of $\text{ZF} \{\alpha_0\}$. Although $\text{qlf}(\alpha_0)$ is not a subset of $\text{qf}(\alpha_0)$ it is the class of Gödel numbers of a subcollection of the wffs of $\text{ZF} \{\alpha_0\}$. This subcollection consists of all wffs that are quantifier free and all wffs that contain only limited quantification.

Theorem 16.13. *For each wff φ of $\text{ZF} \{\alpha_0\}$, $\ulcorner \varphi \urcorner \in \text{qf}(\alpha_0)$.*

Proof. (By induction on the number n of logical symbols in φ .) If $n=0$ then φ is a prime formula i.e. φ is of the form $a \in b$ where a and b are variables or individual constants. Then

$$\ulcorner \varphi \urcorner = \ulcorner a \in b \urcorner \in \text{pf}(\alpha_0) \subseteq \text{qf}(\alpha_0).$$

If $n > 0$ then φ is of the form $(\exists x_i) \psi$ or $\neg \psi$ or $\psi \wedge \eta$. As our induction hypothesis we have

$$\ulcorner \psi \urcorner \in \text{qf}(\alpha_0) \wedge \ulcorner \eta \urcorner \in \text{qf}(\alpha_0)$$

i.e.

$$(\exists m) (\exists n) [\ulcorner \psi \urcorner \in f'm \wedge \ulcorner \eta \urcorner \in f'n].$$

If $l = \max(m, n)$ then

$$\ulcorner \psi \urcorner \in f'l \wedge \ulcorner \eta \urcorner \in f'l$$

hence

$$\ulcorner (\exists x_i) \psi \urcorner \in R_3^{\ulcorner \text{va} \times f \urcorner l},$$

$$\ulcorner \neg \psi \urcorner \in R_4^{\ulcorner f \urcorner l},$$

$$\ulcorner \psi \wedge \eta \urcorner \in R_5^{\ulcorner f \urcorner l)^2}.$$

Therefore

$$\ulcorner \varphi \urcorner \in f(l+1) \subseteq \text{qf}(\alpha_0).$$

Theorem 16.14.

1) For each α in $\text{qf}(\alpha_0)$ there is a wff φ of ZF $\{\alpha_0\}$ for which $\alpha = \ulcorner \varphi \urcorner$.

2) For each α in $\text{qlf}(\alpha_0)$ there is a wff φ of ZF $\{\alpha_0\}$ for which $\alpha = \ulcorner \varphi \urcorner$.

Proof. 1) In view of the definition of $\text{qf}(\alpha_0)$ it is sufficient to prove the result for $\alpha \in f \urcorner n$. This we do by induction on n .

If $\alpha \in f \urcorner 0 = \text{pf}(\alpha_0)$ then there exist variables or individual constants a and b such that

$$\alpha = \ulcorner a \in b \urcorner.$$

Since

$$f(n+1) = f \urcorner n \cup R_3^{\ulcorner \text{va} \times f \urcorner n} \cup R_4^{\ulcorner f \urcorner n} \cup R_5^{\ulcorner f \urcorner n)^2}$$

it follows that if $\alpha \in f(n+1)$ then

$$\begin{aligned} \alpha \in f \urcorner n \vee (\exists \ulcorner x_i \urcorner \in \text{va}) (\exists \ulcorner \varphi \urcorner \in f \urcorner n) [\alpha = \ulcorner (\exists x_i) \varphi \urcorner] \\ \vee (\exists \ulcorner \varphi \urcorner \in f \urcorner n) [\alpha = \ulcorner \varphi \urcorner] \vee (\exists \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner \in f \urcorner n) [\alpha = \ulcorner \varphi \wedge \psi \urcorner]. \end{aligned}$$

2) The proof is left to the reader.

Theorem 16.15.

1) $(\forall \ulcorner \varphi \urcorner \in \text{qf}(\alpha_0)) [\ulcorner \varphi \urcorner \in \text{pf}(\alpha_0) \vee (\exists \ulcorner x_i \urcorner \in \text{va}) (\exists \ulcorner \psi \urcorner \in \text{qf}(\alpha_0))$
 $[\ulcorner \varphi \urcorner = \ulcorner (\exists x_i) \psi \urcorner] \vee (\exists \ulcorner \psi \urcorner \in \text{qf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner \neg \psi \urcorner]$
 $\vee (\exists \ulcorner \psi \urcorner, \ulcorner \eta \urcorner \in \text{qf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner \psi \wedge \eta \urcorner]].$

2) $(\forall \ulcorner \varphi \urcorner \in \text{qlf}(\alpha_0)) [[\ulcorner \varphi \urcorner \in \text{pf}(\alpha_0) \vee (\exists \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner \neg \psi \urcorner]]$
 $\vee (\exists \ulcorner \psi \urcorner, \ulcorner \eta \urcorner \in \text{qlf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner \psi \wedge \eta \urcorner]$
 $\vee (\exists \beta \in \alpha_0) (\exists \ulcorner x_i \urcorner \in \text{va}) (\exists \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner (\exists^\beta x_i) \psi \urcorner]].$

Theorem 16.16.

1) $(\forall \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner \in \text{qf}(\alpha_0)) (\forall \ulcorner x_i \urcorner \in \text{va}) [[\ulcorner (\exists x_i) \varphi \urcorner \in \text{qf}(\alpha_0)]$
 $\wedge [\ulcorner \neg \varphi \urcorner \in \text{qf}(\alpha_0)] \wedge [\ulcorner \varphi \wedge \psi \urcorner \in \text{qf}(\alpha_0)]]].$

2) $(\forall \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0)) (\forall \ulcorner x_i \urcorner \in \text{va}) [[\ulcorner \neg \varphi \urcorner \in \text{qlf}(\alpha_0)]$
 $\wedge [\ulcorner \varphi \wedge \psi \urcorner \in \text{qlf}(\alpha_0)] \wedge [\ulcorner (\exists^\beta x_i) \varphi \urcorner \in \text{qlf}(\alpha_0)]]].$

The proofs are left to the reader.

Theorem 16.17. *If $\alpha_0 \subseteq \mathcal{M}$ then*

- 1) $\text{qf}(\alpha_0) \subseteq \mathcal{M} \wedge [x \in \text{qf}(\alpha_0)] \text{ Abs } \mathcal{M}$,
- 2) $\text{qlf}(\alpha_0) \subseteq \mathcal{M} \wedge [x \in \text{qlf}(\alpha_0)] \text{ Abs } \mathcal{M}$.

Proof. 1) With f as in Definition 16.11 we see that $f'0 = \text{pf}(\alpha_0) \subseteq \mathcal{M}$.
If $f'n \subseteq \mathcal{M}$ then since

$$f(n+1) = f'n \cup R_3''(\text{va} \times f'n) \cup R_4''f'n \cup R_5''(f'n)^2$$

and since \mathcal{M} is closed with respect to J it follows that $f(n+1) \subseteq \mathcal{M}$.
Therefore

$$\text{qf}(\alpha_0) = \bigcup_{n < \omega} f'n \subseteq \mathcal{M}.$$

To prove $[x \in \text{qf}(\alpha_0)] \text{ Abs } \mathcal{M}$ we note that $x \in \text{qf}(\alpha_0)$ iff

$$\begin{aligned} & (\exists f) [f \mathcal{F}n \omega \wedge f'0 = \text{pf}(\alpha_0) \\ & \wedge (\forall n \in \omega) [f(n+1) = f'n \cup R_3''(\text{va} \times f'n) \cup R_4''f'n \cup R_5''(f'n)^2] \\ & \wedge (\exists n \in \omega) [x \in f'n]]. \end{aligned}$$

Also $x \in \text{qf}(\alpha_0)$ iff

$$\begin{aligned} & (\forall f) [f \mathcal{F}n \omega \wedge f'0 = \text{pf}(\alpha_0) \\ & \wedge (\forall n \in \omega) [f(n+1) = f'n \cup R_3''(\text{va} \times f'n) \cup R_4''f'n \cup R_5''(f'n)^2] \\ & \rightarrow (\exists n \in \omega) [x \in f'n]]. \end{aligned}$$

But from the definitions of R_3 , R_4 , and R_5 , $x \in f(n+1)$ iff

$$\begin{aligned} & [x \in f'n] \vee (\exists \gamma) (\exists \delta) [\gamma \in \text{va} \wedge \delta \in f'n \wedge x = J'\langle \gamma, \delta, 3 \rangle] \\ & \vee (\exists \gamma) [\gamma \in f'n \wedge x = J'\langle \gamma, 4, 4 \rangle] \vee (\exists \gamma) (\exists \delta) [\gamma \in f'n \wedge x = J'\langle \gamma, \delta, 5 \rangle]. \end{aligned}$$

Since $J'\langle \alpha, \beta, n \rangle$ is absolute with respect to \mathcal{M} it follows from Theorem 13.8 that

$$[x \in \text{qf}(\alpha_0)] \text{ Abs } \mathcal{M}.$$

- 2) The proof is left to the reader.

Exercises. *Prove the following*

- 1) $\alpha < J'\langle 1, \alpha, n \rangle$.
- 2) $\alpha \neq 0 \rightarrow \beta < J'\langle \alpha, \beta, n \rangle$.
- 3) $\max(\alpha, \beta) < R_3'\langle \alpha, \beta \rangle$.
- 4) $\alpha < R_4'\alpha$.
- 5) $\max(\alpha, \beta) < R_5'\langle \alpha, \beta \rangle$.
- 6) $\text{va} \subseteq \mathcal{M}$.
- 7) $\text{ic}(\alpha_0) \text{ Abs } \mathcal{M}$.

- 8) $0 \notin \text{pf}(\alpha_0) \wedge 1 \notin \text{pf}(\alpha_0)$.
- 9) $\text{pf}(\alpha_0) \text{ Abs } \mathcal{M}$.
- 10) $0 \notin \text{qf}(\alpha_0) \wedge 1 \notin \text{qf}(\alpha_0)$.
- 11) $\text{qf}(\alpha_0) \text{ Abs } \mathcal{M}$.

Remark. The class of wffs “wff(α_0)” we will define as a certain subclass of the class of quasi formulas qf(α_0). For its definition we introduce the notion of a string, or pyramid, of quasi-formulas.

Definition 16.18.

- 1) $\text{Sqf}(\alpha_0; f) \stackrel{\Delta}{\longleftrightarrow} (\exists n \in \omega) [f \mathcal{F}_n(n+1) \wedge \mathcal{W}(f) \subseteq \text{qf}(\alpha_0)$
 $\wedge f'0 \in \text{pf}(\alpha_0) \wedge (\forall i < n) [(\exists \alpha \in \text{va}) [f(i+1) = R'_3 \langle \alpha, f' i \rangle]$
 $\vee [f(i+1) = R'_4 f' i] \vee (\exists \alpha \in \text{qf}(\alpha_0)) [f(i+1) = R'_5 \langle \alpha, f' i \rangle]$
 $\vee f(i+1) = R'_5 \langle f' i, \alpha \rangle]]$.
- 2) $\text{Sqlf}(\alpha_0; f) \stackrel{\Delta}{\longleftrightarrow} (\exists n \in \omega) [f \mathcal{F}_n(n+1) \wedge \mathcal{W}(f) \subseteq \text{qlf}(\alpha_0)$
 $\wedge f'0 \in \text{pf}(\alpha_0) \wedge (\forall i < n) [f(i+1) = R'_4 f' i$
 $\vee (\exists \alpha \in \text{qlf}(\alpha_0)) [f(i+1) = R'_5 \langle \alpha, f' i \rangle \vee f(i+1) = R'_5 \langle f' i, \alpha \rangle]$
 $\vee (\exists \beta \in \alpha_0) (\exists \gamma \in \text{va}) [f(i+1) = R'_6 \langle \beta, \gamma, f' i \rangle]]$.

Remark. The idea in Definition 16.18 is that each formula of ZF $\{\alpha_0\}$ can be thought of as having been built up from prime formulas. Consider, for example,

$$(\exists x_1) (\exists x_2) \neg [x_1 \in x_2 \wedge \neg x_1 \in x_2].$$

This formula can be built up from prime formulas in accordance with Definition 16.18 in the following way.

$$\begin{aligned} & x_1 \in x_2, \\ & [x_1 \in x_2] \wedge \neg [x_1 \in x_2], \\ & \neg [[x_1 \in x_2] \wedge \neg [x_1 \in x_2]], \\ & (\exists x_2) \neg [[x_1 \in x_2] \wedge \neg [x_1 \in x_2]], \\ & (\exists x_1) (\exists x_2) \neg [[x_1 \in x_2] \wedge \neg [x_1 \in x_2]]. \end{aligned}$$

This sequence of formulas illustrates our intuitive notion of a string of quasi formulas. However to code this into ZF our formal definition of a string is a function from integers to Gödel numbers of quasi formulas. For the example at hand this is a function with domain 5 so defined that

$$\begin{aligned} f'0 &= \ulcorner x_1 \in x_2 \urcorner, \\ f'1 &= \ulcorner x_1 \in x_2 \wedge \neg x_1 \in x_2 \urcorner, \\ &\text{etc.} \end{aligned}$$

It should be noted that there may be more than one string of quasi formulas building up to a given formula. For the foregoing example we could have

$$\begin{aligned}
& x_1 \in x_2, \\
& \neg [x_1 \in x_2], \\
& [x_1 \in x_2] \wedge \neg [x_1 \in x_2], \\
& \neg [[x_1 \in x_2] \wedge \neg [x_1 \in x_2]], \\
& (\exists x_2) \neg [[x_1 \in x_2] \wedge \neg [x_1 \in x_2]], \\
& (\exists x_1) (\exists x_2) \neg [[x_1 \in x_2] \wedge \neg [x_1 \in x_2]].
\end{aligned}$$

Theorem 16.19.

- 1) $\text{Sqf}(\alpha_0; f) \wedge [i \in \mathcal{D}(f)] \wedge [f^* i \in \text{pf}(\alpha)] \rightarrow i = 0.$
- 2) $\text{Sqlf}(\alpha_0; f) \wedge [i \in \mathcal{D}(f)] \wedge [f^* i \in \text{pf}(\alpha_0)] \rightarrow i = 0.$

Proof. 1) $i \neq 0 \rightarrow (\exists j) [i = j + 1].$ Then from Definition 16.18

$$\begin{aligned}
& (\exists \alpha \in \text{var}) [f^* i = R_3^* \langle \alpha, f^* j \rangle] \vee [f^* i = R_4^* f^* j] \\
& \vee (\exists \alpha \in \text{qf}(\alpha_0)) [f^* i = R_5^* \langle \alpha, f^* j \rangle \vee f^* i = R_5^* \langle f^* j, \alpha \rangle].
\end{aligned}$$

In each case $f^* i \notin \text{pf}(\alpha_0).$

- 2) The proof is left to the reader.

Theorem 16.20.

- 1) $\text{Sqf}(\alpha_0; f) \wedge [i \in \mathcal{D}(f)] \rightarrow [i < f^* i].$
- 2) $\text{Sqlf}(\alpha_0; f) \wedge [i \in \mathcal{D}(f)] \rightarrow [i < f^* i].$

Proof. 1) (By induction on i .) If $i = 0$ then since

$$\begin{aligned}
& [f^* 0 \in \text{pf}(\alpha_0)] \wedge [0 = J^* \langle 0, 0, 0 \rangle \notin \text{pf}(\alpha_0)] \\
& 0 < f^* 0.
\end{aligned}$$

As our induction hypothesis we assume

$$i < f^* i.$$

If $i + 1 \in \mathcal{D}(f)$ then $i + 1 \leq f^* i$ and

$$\begin{aligned}
& (\exists \alpha \in \text{va}) [f(i + 1) = R_3^* \langle \alpha, f^* i \rangle] \vee [f(i + 1) = R_4^* f^* i] \\
& \vee (\exists \alpha \in \text{qf}(\alpha_0)) [f(i + 1) = R_5^* \langle \alpha, f^* i \rangle \vee f(i + 1) = R_5^* \langle f^* i, \alpha \rangle].
\end{aligned}$$

But

$$\begin{aligned}
R_3^* \langle \alpha, f^* i \rangle &= 9J_0^* \langle \alpha, f^* i \rangle + 3 > J_0^* \langle \alpha, f^* i \rangle \geq f^* i, \\
R_4^* f^* i &= 9J_0^* \langle f^* i, 4 \rangle + 4 > J_0^* \langle f^* i, 4 \rangle \geq f^* i, \\
R_5^* \langle \alpha, f^* i \rangle &= 9J_0^* \langle \alpha, f^* i \rangle + 5 > J_0^* \langle \alpha, f^* i \rangle \geq f^* i, \\
R_5^* \langle f^* i, \alpha \rangle &= 9J_0^* \langle f^* i, \alpha \rangle + 5 > J_0^* \langle f^* i, \alpha \rangle \geq f^* i.
\end{aligned}$$

Therefore, in each case,

$$f(i) < f(i+1).$$

2) The proof is left to the reader.

Theorem 16.21.

1) $\text{Sqf}(\alpha_0; f) \wedge 0 < i \in \mathcal{D}(f) \rightarrow \text{Sqf}(\alpha_0; f \uparrow i)$.

2) $\text{Sqlf}(\alpha_0; f) \wedge 0 < i \in \mathcal{D}(f) \rightarrow \text{Sqlf}(\alpha_0; f \uparrow i)$.

Proof. 1) Since $i > 0$, $(\exists j) [i = j + 1]$; hence $(f \uparrow i) \mathcal{F}_n(j + 1)$.

$$\mathcal{W}(f \uparrow i) \subseteq \mathcal{W}(f) \subseteq \text{qf}(\alpha_0).$$

$$(f \uparrow i)'0 = f'0 \in \text{pf}(\alpha_0).$$

$$(\forall k < j) [(f \uparrow i)'(k + 1) = f'(k + 1)].$$

Therefore

$$(\exists \alpha \in \text{va}) [(f \uparrow i)'(k + 1) = R_3' \langle \alpha, (f \uparrow i)'k \rangle]$$

or

$$(f \uparrow i)'(k + 1) = R_4'(f \uparrow i)'k$$

or

$$(\exists \alpha \in \text{qf}(\alpha_0)) [(f \uparrow i)'(k + 1) = R_5' \langle \alpha, (f \uparrow i)'k \rangle]$$

$$\vee (f \uparrow i)'(k + 1) = R_5' \langle (f \uparrow i)'k, \alpha \rangle].$$

2) The proof is left to the reader.

Remark. The domain of a string of quasi formulas is simply the number of steps required to build the last formula in the string from prime formulas. We will refer to the domain of a string as the length of the string and the last formulas as the end of the string.

Definition 16.22. $\text{end}(f) = f'(\cup(\mathcal{D}(f)))$.

Theorem 16.23.

1) $'\varphi' \in \text{qf}(\alpha_0) \rightarrow (\exists f) [\text{Sqf}(\alpha_0; f) \wedge '\varphi' = \text{end}(f)]$.

2) $'\varphi' \in \text{qlf}(\alpha_0) \rightarrow (\exists f) [\text{Sqlf}(\alpha_0; f) \wedge '\varphi' = \text{end}(f)]$.

Proof. 1) (By induction on $'\varphi'$.) From Theorem 16.15

$$' \varphi ' \in \text{pf}(\alpha_0) \vee (\exists 'x_i' \in \text{va}) (\exists ' \psi ' \in \text{qf}(\alpha_0)) [' \varphi ' = '(\exists x_i) \psi ']$$

$$\vee (\exists ' \psi ' \in \text{qf}(\alpha_0)) [' \varphi ' = ' \neg \psi ']$$

$$\vee (\exists ' \psi ' \in \text{qf}(\alpha_0)) (\exists ' \eta ' \in \text{qf}(\alpha_0)) [' \varphi ' = ' \psi \wedge \eta '].$$

If $\ulcorner\varphi\urcorner \in \text{pf}(\alpha_0)$ and if $f = \{\langle 0, \ulcorner\varphi\urcorner\rangle\}$ then $\text{Sqf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f)$. In all other cases $\ulcorner\psi\urcorner \in \text{qf}(\alpha_0)$ and $\ulcorner\psi\urcorner < \ulcorner\varphi\urcorner$. Therefore by the induction hypothesis

$$(\exists g) [\text{Sqf}(\alpha_0; g) \wedge \ulcorner\psi\urcorner = \text{end}(g)].$$

Thus if

$$f = g \cup \{\langle \mathcal{D}(g), \ulcorner\varphi\urcorner\rangle\}$$

then

$$\text{Sqf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f).$$

2) The proof is left to the reader.

Theorem 16.24.

$$1) \quad \alpha_0 \subseteq \mathcal{M} \wedge \text{Sqf}(\alpha_0; f) \rightarrow f \in \mathcal{M}.$$

$$2) \quad \alpha_0 \subseteq \mathcal{M} \wedge \text{Sqlf}(\alpha_0; f) \rightarrow f \in \mathcal{M}.$$

Proof. f is a finite subset of $\omega \times \text{qf}(\alpha_0)$ or $\omega \times \text{qlf}(\alpha_0)$.

Definition 16.25.

$$1) \quad (\forall \ulcorner\varphi\urcorner \in \text{qf}(\alpha_0)) [\text{deg}(\alpha_0; \ulcorner\varphi\urcorner) = \cup \{\mathcal{D}(f) \mid \text{Sqf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f)\}].$$

$$2) \quad (\forall \ulcorner\varphi\urcorner \in \text{qlf}(\alpha_0)) [\text{deg}_l(\alpha_0; \ulcorner\varphi\urcorner) = \cup \{\mathcal{D}(f) \mid \text{Sqlf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f)\}].$$

Remark. Earlier we pointed out that our intuitive notion of a string, or pyramid, of quasi formulas stems from the observation that each formula can be built from prime formulas in a manner specified by Definition 16.18. We observed that a given formula can be the end of strings of different length. By the degree of a formula we mean simply the length of the longest string ending with that formula. That there is a longest string of quasi formulas ending in a given formula φ is intuitively clear. Indeed the longest string cannot be longer than the number of symbols in φ .

Theorem 16.26.

$$1) \quad (\forall \ulcorner\varphi\urcorner \in \text{qf}(\alpha_0)) [\text{deg}(\alpha_0; \ulcorner\varphi\urcorner) = \beta \\ \iff (\forall f) [\text{Sqf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f) \rightarrow \mathcal{D}(f) \leq \beta] \\ \wedge (\exists f) [\text{Sqf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f) \wedge \mathcal{D}(f) = \beta]].$$

$$2) \quad (\forall \ulcorner\varphi\urcorner \in \text{qlf}(\alpha_0)) [\text{deg}_l(\alpha_0; \ulcorner\varphi\urcorner) = \beta \\ \iff (\forall f) [\text{Sqlf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f) \rightarrow \mathcal{D}(f) \leq \beta] \\ \wedge (\exists f) [\text{Sqlf}(\alpha_0; f) \wedge \ulcorner\varphi\urcorner = \text{end}(f) \wedge \mathcal{D}(f) = \beta]].$$

Proof. Obvious from Definition 16.25.

Theorem 16.27.

- 1) $\ulcorner \varphi \urcorner \in \text{pf}(\alpha_0) \rightarrow \text{deg}(\alpha_0; \ulcorner \varphi \urcorner) = 1.$
- 2) $\ulcorner x_i \urcorner \in \text{va} \wedge \ulcorner \varphi \urcorner \in \text{qf}(\alpha_0) \rightarrow \text{deg}(\alpha_0; \ulcorner (\exists x_i) \varphi \urcorner) = 1 + \text{deg}(\alpha_0; \ulcorner \varphi \urcorner).$
- 3) $\ulcorner \varphi \urcorner \in \text{qf}(\alpha_0) \rightarrow \text{deg}(\alpha_0; \ulcorner \neg \varphi \urcorner) = 1 + \text{deg}(\alpha_0; \ulcorner \varphi \urcorner).$
- 4) $\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner \in \text{qf}(\alpha_0) \rightarrow \text{deg}(\alpha_0; \ulcorner \varphi \wedge \psi \urcorner) = 1$
 $+ \max(\text{deg}(\alpha_0; \ulcorner \varphi \urcorner), \text{deg}(\alpha_0; \ulcorner \psi \urcorner)).$

Proof. 1) $\text{Sqf}(\alpha_0; f) \wedge \ulcorner \varphi \urcorner = \text{end}(f) \wedge f \mathcal{F}_n(i+1) \rightarrow f^i = \ulcorner \varphi \urcorner.$

But

$$\ulcorner \varphi \urcorner \in \text{pf}(\alpha_0) \rightarrow i = 0.$$

Thus every string of quasi formulas ending in the prime formula φ has length 1. Furthermore for each prime formula φ there is a string of quasi formulas ending in that formula. Hence by Theorem 16.26

$$\text{deg}(\alpha_0; \ulcorner \varphi \urcorner) = 1.$$

- 2) $\text{Sqf}(\alpha_0; f) \wedge \ulcorner (\exists x_i) \varphi \urcorner = \text{end}(f) = f^{i+1} \rightarrow \ulcorner \varphi \urcorner = f^i.$

Thus every string of quasi formulas ending in $\ulcorner (\exists x_i) \varphi \urcorner$ is one longer than some string of quasi formulas ending in $\ulcorner \varphi \urcorner$. Consequently every string of quasi formulas ending in $\ulcorner (\exists x_i) \varphi \urcorner$ is not longer than $1 + \text{deg}(\alpha_0; \ulcorner \varphi \urcorner)$.

Since there is a string ending in $\ulcorner \varphi \urcorner$ and of length $\text{deg}(\alpha_0; \ulcorner \varphi \urcorner)$ there is a string ending in $\ulcorner (\exists x_i) \varphi \urcorner$ and of length $1 + \text{deg}(\alpha_0; \ulcorner \varphi \urcorner)$. Therefore

$$\text{deg}(\alpha_0; \ulcorner (\exists x_i) \varphi \urcorner) = 1 + \text{deg}(\alpha_0; \ulcorner \varphi \urcorner).$$

- 3) $\text{Sqf}(\alpha_0; f) \wedge \ulcorner \neg \varphi \urcorner = \text{end}(f) = f^{i+1} \rightarrow \ulcorner \varphi \urcorner = f^i.$

Then as before

$$\text{deg}(\alpha_0; \ulcorner \neg \varphi \urcorner) = 1 + \text{deg}(\alpha_0; \ulcorner \varphi \urcorner).$$

- 4) The proof is left to the reader.

Theorem 16.28.

- 1) $\ulcorner \varphi \urcorner \in \text{pf}(\alpha_0) \rightarrow \text{deg}_l(\alpha_0; \ulcorner \varphi \urcorner) = 1.$
- 2) $\ulcorner \varphi \urcorner \in \text{qlf}(\alpha_0) \rightarrow \text{deg}_l(\alpha_0; \ulcorner \neg \varphi \urcorner) = 1 + \text{deg}_l(\alpha_0; \ulcorner \varphi \urcorner).$
- 3) $\ulcorner \varphi \urcorner, \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0) \rightarrow \text{deg}_l(\alpha_0; \ulcorner \varphi \wedge \psi \urcorner) = 1$
 $+ \max(\text{deg}_l(\alpha_0; \ulcorner \varphi \urcorner), \text{deg}_l(\alpha_0; \ulcorner \psi \urcorner)).$
- 4) $\beta \in \alpha_0 \wedge \ulcorner x_i \urcorner \in \text{va} \wedge \ulcorner \varphi \urcorner \in \text{qlf}(\alpha_0) \rightarrow \text{deg}_l(\alpha_0; \ulcorner (\exists^\beta x_i) \varphi \urcorner) = 1$
 $+ \text{deg}^l(\alpha_0; \ulcorner \varphi \urcorner).$
- 5) The proof is left to the reader.

Theorem 16.29.

- 1) $(\forall \ulcorner \varphi \urcorner \in \text{qf}(\alpha_0)) [\text{deg}(\alpha_0; \ulcorner \varphi \urcorner) < \min(\omega, \ulcorner \varphi \urcorner)].$
- 2) $(\forall \ulcorner \varphi \urcorner \in \text{qlf}(\alpha_0)) [\text{deg}_l(\alpha_0; \ulcorner \varphi \urcorner) < \min(\omega, \ulcorner \varphi \urcorner)].$

Proof. 1) (By induction on $\ulcorner \varphi \urcorner$.) Since $\ulcorner \varphi \urcorner \in \text{qf}(\alpha_0)$
 $\ulcorner \varphi \urcorner \in \text{pf}(\alpha_0) \vee (\exists \ulcorner x_i \urcorner \in \text{va}) (\exists \ulcorner \psi \urcorner \in \text{qf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner (\exists x_i) \psi \urcorner]$
 $\vee (\exists \ulcorner \psi \urcorner \in \text{qf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner \neg \psi \urcorner]$
 $\vee (\exists \ulcorner \eta \urcorner, \ulcorner \psi \urcorner \in \text{qf}(\alpha_0)) [\ulcorner \varphi \urcorner = \ulcorner \psi \wedge \eta \urcorner].$

Then by Theorem 16.27

$$\text{deg}(\alpha_0; \ulcorner \varphi \urcorner) = 1 \vee [\text{deg}(\alpha_0; \ulcorner \varphi \urcorner) = 1 + \text{deg}(\alpha_0, \ulcorner \psi \urcorner)] \\ \vee \text{deg}(\alpha_0; \ulcorner \varphi \urcorner) = 1 + \max(\text{deg}(\alpha_0, \ulcorner \psi \urcorner), \text{deg}(\alpha_0; \ulcorner \eta \urcorner)).$$

If $\text{deg}(\alpha_0, \ulcorner \psi \urcorner) = 1$ then since $\ulcorner \varphi \urcorner \in \text{qf}(\alpha_0)$ implies $\ulcorner \varphi \urcorner > 1$

$$\text{deg}(\alpha_0, \ulcorner \varphi \urcorner) < \min(\omega, \ulcorner \varphi \urcorner).$$

In all other cases $\ulcorner \eta \urcorner < \ulcorner \varphi \urcorner$, $\ulcorner \psi \urcorner < \ulcorner \varphi \urcorner$ and hence by the induction hypothesis

$$\text{deg}(\alpha_0; \ulcorner \eta \urcorner) < \min(\omega, \ulcorner \eta \urcorner) \wedge \text{deg}(\alpha_0, \ulcorner \psi \urcorner) < \min(\omega, \ulcorner \psi \urcorner).$$

Therefore

$$\text{deg}(\alpha_0, \ulcorner \varphi \urcorner) = 1 + \text{deg}(\alpha_0, \ulcorner \psi \urcorner) < \min(\omega, \ulcorner \psi \urcorner + 1) \leq \min(\omega, \ulcorner \varphi \urcorner)$$

or

$$\text{deg}(\alpha_0; \ulcorner \varphi \urcorner) = 1 + \max(\text{deg}(\alpha_0; \ulcorner \psi \urcorner), \text{deg}(\alpha_0; \ulcorner \eta \urcorner)) < \min(\omega, \ulcorner \varphi \urcorner).$$

Remark. We next introduce the notion of φ being a subquasi formula of ψ .

Definition 16.30.

- 1) $\text{Subqf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \stackrel{\Delta}{\leftrightarrow} [\ulcorner \varphi \urcorner \in \text{qf}(\alpha_0)] \wedge [\ulcorner \psi \urcorner \in \text{qf}(\alpha_0)] \\ \wedge (\exists f)[\text{Sqf}(\alpha_0; f) \wedge \ulcorner \psi \urcorner = \text{end}(f) \wedge (\exists i \in \mathcal{D}(f)) [\ulcorner \varphi \urcorner = f \ulcorner i \urcorner]].$
- 2) $\text{Subqlf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \stackrel{\Delta}{\leftrightarrow} [\ulcorner \varphi \urcorner \in \text{qlf}(\alpha_0)] \wedge [\ulcorner \psi \urcorner \in \text{qlf}(\alpha_0)] \\ \wedge (\exists f)[\text{Sqlf}(\alpha_0; f) \wedge \ulcorner \psi \urcorner = \text{end}(f) \wedge (\exists i \in \mathcal{D}(f)) [\ulcorner \psi \urcorner = f \ulcorner i \urcorner]].$

Theorem 16.31.

- 1) $\text{Subqf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \wedge \text{Subqf}(\alpha_0; \ulcorner \psi \urcorner, \ulcorner \eta \urcorner) \rightarrow \text{Subqf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \eta \urcorner).$
- 2) $\text{Subqlf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \wedge \text{Subqlf}(\alpha_0; \ulcorner \psi \urcorner, \ulcorner \eta \urcorner) \rightarrow \text{Subqlf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \eta \urcorner).$

Proof. 1) (By induction on $\text{deg}(\alpha_0; \ulcorner \eta \urcorner$.) Since $\ulcorner \eta \urcorner \in \text{qf}(\alpha_0)$

$$(\exists \ulcorner x_i \urcorner \in \text{va}) (\exists \ulcorner \zeta_0 \urcorner \in \text{qf}(\alpha_0)) [\ulcorner \eta \urcorner = \ulcorner (\exists x_i) \zeta_0 \urcorner]$$

or

$$(\exists ' \zeta_0 ' \in \text{qf}(\alpha_0)) [' \eta ' = ' \neg \zeta_0 ']$$

or

$$(\exists ' \zeta_0 ' \in \text{qf}(\alpha_0)) (\exists ' \zeta_1 ' \in \text{qf}(\alpha_0)) [' \eta ' = ' \zeta_0 \wedge \zeta_1 '] .$$

In each case we have

$$\text{deg}(\alpha_0, ' \zeta_0 ') < \text{deg}(\alpha_0, ' \eta ') \wedge \text{deg}(\alpha_0, ' \zeta_1 ') < \text{deg}(\alpha_0, ' \eta ') .$$

Since ' ψ ' is a subformula of ' η ' either ' $\psi = \eta$ ', and hence the conclusion, or $\text{Subqf}(\alpha_0; ' \psi ', ' \zeta_0 ') \vee \text{Subqf}(\alpha_0; ' \psi ', ' \zeta_1 ') .$ With the aid of the induction hypothesis it follows, in each case, that

$$\text{Subqf}(\alpha_0; ' \varphi ', ' \zeta_1 ') \vee \text{Subqf}(\alpha_0; ' \varphi ', ' \zeta_0 ') .$$

But this means that there is a string f of quasi formulas ending in ' ζ_1 ' or in ' ζ_0 ' that assumes the value ' φ ' somewhere in its domain. Adjoining to f the ordered pair $\langle \mathcal{D}(f), ' \eta ' \rangle$ we obtain a string of quasi formulas ending in ' η ' that assumes the value ' φ ' at some point in its domain. Therefore

$$\text{Subqf}(\alpha_0; ' \varphi ', ' \eta ') .$$

2) The proof is left to the reader.

Remark. Intuitively if φ is a subformula of ψ then φ is the end of some string of formulas that can be extended to a string ending in ψ .

We next formalize the ideas of a variable occurring bound in a formula, and of a variable occurring in a formula.

Definition 16.32.

- 1) $\text{Ob}(\alpha_0; ' x_i ', ' \varphi ') \stackrel{\Delta}{\longleftrightarrow} ' x_i ' \in \text{va} \wedge ' \varphi ' \in \text{qf}(\alpha_0) \wedge (\exists ' \psi ' \in \text{qf}(\alpha_0)) [\text{Subqf}(\alpha_0; ' (\exists x_i) \psi ', ' \varphi ')] .$
- 2) $\text{Ob}_l(\alpha_0; ' x_i ', ' \varphi ') \longleftrightarrow ' x_i ' \in \text{va} \wedge ' \varphi ' \in \text{qlf}(\alpha_0) \wedge (\exists \beta \in \alpha_0) (\exists ' \psi ' \in \text{qlf}(\alpha_0)) [\text{Subqlf}(\alpha_0; ' (\exists^\beta x_i) \psi ', ' \varphi ')] .$

Definition 16.33.

$$\text{O}(\alpha_0; ' x_i ', ' \varphi ') \stackrel{\Delta}{\longleftrightarrow} (\exists ' a ' \in \text{vc}(\alpha_0)) [\text{Subqf}(\alpha_0; ' x_i \in a ', ' \varphi ') \vee \text{Subqf}(\alpha_0; ' a \in x_i ', ' \varphi ')] .$$

Remark. We arrive finally at our formal notion of a wff, namely, a wff is a quasi formula in which the scope of each quantifier $(\exists x_i)$ does not have a bound occurrence of the quantifier variable x_i . Thus

$$(\exists x_1) (\exists x_1) [x_1 = x_1]$$

is a quasi formula that is well formed by the formation rules of our language, but for our purposes it is excluded from the formal wffs.

Definition 16.34.

- 1) $\text{wff}(\alpha_0) = \{ \ulcorner \varphi \urcorner \in \text{qf}(\alpha_0) \mid (\forall \ulcorner x_i \urcorner \in \text{va}) (\forall \ulcorner \psi \urcorner \in \text{qf}(\alpha_0))$
 $[\text{Subqf}(\alpha_0; \ulcorner (\exists x_i) \psi \urcorner, \ulcorner \varphi \urcorner) \rightarrow \neg \text{Ob}(\alpha_0; \ulcorner x_i \urcorner, \ulcorner \psi \urcorner)] \}$.
- 2) $\text{wflf}(\alpha_0) = \{ \ulcorner \varphi \urcorner \in \text{qlf}(\alpha_0) \mid (\forall \beta \in \alpha_0) (\forall \ulcorner x_i \urcorner \in \text{va}) (\forall \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0))$
 $[\text{Subqlf}(\alpha_0; \ulcorner (\exists^\beta x_i) \psi \urcorner, \ulcorner \varphi \urcorner) \rightarrow \neg \text{Ob}_l(\alpha_0; \ulcorner x_i \urcorner, \ulcorner \psi \urcorner)] \}$.

Theorem 16.35.

- 1) $\text{Sqf}(\alpha_0; f) \text{ Abs } \mathcal{M}$,
- 2) $\text{Subqf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \text{ Abs } \mathcal{M}$,
- 3) $\text{Ob}(\alpha_0; \ulcorner x_i \urcorner, \ulcorner \varphi \urcorner) \text{ Abs } \mathcal{M}$.

Proof.

- 1) $\text{Sqf}(\alpha_0; f) \longleftrightarrow (\exists i \in \omega) [f \mathcal{F}_n(i+1) \wedge \mathcal{W}(f) \subseteq \text{qf}(\alpha_0)$
 $\wedge f^*0 \in \text{pf}(\alpha_0) \wedge (\forall j \in i) [(\exists \alpha \in \text{va}) [f^*(j+1) = J^* \langle \alpha, f^*j, 3 \rangle]$
 $\vee f^*(j+1) = J^* \langle f^*j, 4, 4 \rangle$
 $\vee (\exists \alpha \in \text{qf}(\alpha_0)) [f^*(j+1) = J^* \langle \alpha, f^*j, 5 \rangle$
 $\vee f^*(j+1) = J^* \langle f^*j, \alpha, 5 \rangle]]$.

From Theorems 16.4, 16.7, and 16.17 it follows that

$$\text{Sqf}(\alpha_0; f) \text{ Abs } \mathcal{M}.$$

- 2) $\text{Subqf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \longleftrightarrow \ulcorner \varphi \urcorner \in \text{qf}(\alpha_0) \wedge \ulcorner \psi \urcorner \in \text{qf}(\alpha_0)$
 $\wedge (\exists f) [\text{Sqf}(\alpha_0; f) \wedge \ulcorner \psi \urcorner = f^*(\cup(\mathcal{D}(f)))$
 $\wedge (\exists i) [i \in \mathcal{D}(f) \wedge \ulcorner \varphi \urcorner = f^*i]]$.

From 1) above and Theorem 16.24 we conclude that

$$\text{Subqf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \text{ Abs } \mathcal{M}.$$

- 3) $\text{Ob}(\alpha_0; \ulcorner x_i \urcorner, \ulcorner \varphi \urcorner) \longleftrightarrow \ulcorner x_i \urcorner \in \text{va} \wedge \ulcorner \varphi \urcorner \in \text{qf}(\alpha_0)$
 $\wedge (\exists \ulcorner \psi \urcorner) [\ulcorner \psi \urcorner \in \text{qf}(\alpha_0) \wedge \text{Subqf}(\alpha_0; \ulcorner (\exists x_i) \psi \urcorner, \ulcorner \varphi \urcorner)]$.

Theorem 16.36.

- 1) $\text{Sqlf}(\alpha_0; f) \text{ Abs } \mathcal{M}$,
- 2) $\text{Subqlf}(\alpha_0; \ulcorner \varphi \urcorner, \ulcorner \psi \urcorner) \text{ Abs } \mathcal{M}$,
- 3) $\text{Ob}_l(\alpha_0; \ulcorner x_i \urcorner, \ulcorner \varphi \urcorner) \text{ Abs } \mathcal{M}$.

The proof is left to the reader.

Theorem 16.37. If $\alpha_0 \subseteq \mathcal{M}$ then

- 1) $\text{wff}(\alpha_0) \subseteq \mathcal{M} \wedge [x \in \text{wff}(\alpha_0)] \text{ Abs } \mathcal{M}$,
- 2) $\text{wflf}(\alpha_0) \subseteq \mathcal{M} \wedge [x \in \text{wflf}(\alpha_0)] \text{ Abs } \mathcal{M}$.

Proof. 1) $\text{wff}(\alpha_0) \subseteq \text{qf}(\alpha_0) \subseteq \mathcal{M}$.

To prove that $x \in \text{wff}(\alpha_0)$ is absolute with respect to \mathcal{M} we need only note that $x \in \text{wff}(\alpha_0)$ iff

$$x \in \text{qf}(\alpha_0) \wedge (\forall 'x_i' \in \text{va}) (\forall ' \psi ' \in \text{qf}(\alpha_0)) \\ [\text{Subqf}(\alpha_0; '(\exists x_i) \psi', x) \rightarrow \neg \text{Ob}(\alpha_0; 'x_i', ' \psi ')] .$$

From Theorem 16.35 we then conclude that

$$[x \in \text{wff}(\alpha_0)] \text{ Abs } \mathcal{M} .$$

- 2) The proof is left to the reader.

Remark. We next define the class of subformulas of ' φ '.

Definition 16.38.

- 1) $\text{Sub}(\alpha_0; ' \varphi ') \triangleq \{ ' \psi ' \mid \text{Subqf}(\alpha_0; ' \psi ', ' \varphi ') \}$.
- 2) $\text{Sub}_l(\alpha_0; ' \varphi ') \triangleq \{ ' \psi ' \mid \text{Subqlf}(\alpha_0; ' \psi ', ' \varphi ') \}$.

Theorem 16.39.

- 1) $\text{Sqf}(\alpha_0; f) \wedge \text{end}(f) \in \text{wff}(\alpha_0) \rightarrow (\forall i \in \mathcal{D}(f)) [f'i \in \text{wff}(\alpha_0)]$.
- 2) $\text{Sqlf}(\alpha_0; f) \wedge \text{end}(f) \in \text{wflf}(\alpha_0) \rightarrow (\forall i \in \mathcal{D}(f)) [f'i \in \text{wflf}(\alpha_0)]$.

Proof. 1) Since $f'i$ is a subformula of $\text{end}(f)$ it follows from Theorem 16.31 that any subformula of $f'i$ is a subformula of $\text{end}(f)$. If $f'i \notin \text{wff}(\alpha_0)$ then $f'i$ contains a subformula ' $(\exists x_i) \psi$ ' with $\text{Ob}(\alpha_0; 'x_i', ' \psi ')$. But since $\text{end}(f)$ must contain this same formula we arrive at the contradiction

$$\text{end}(f) \notin \text{wff}(\alpha_0) .$$

- 2) The proof is left to the reader.

Theorem 16.40.

- 1) $' \varphi ' \in \text{wff}(\alpha_0) \rightarrow \text{Sub}(\alpha_0; ' \varphi ') \subseteq \text{wff}(\alpha_0)$.
- 2) $' \varphi ' \in \text{wflf}(\alpha_0) \rightarrow \text{Sub}_l(\alpha_0; ' \varphi ') \subseteq \text{wflf}(\alpha_0)$.

Proof. Theorem 16.39.

Remark. We next define the class of closed wffs and the class of closed limited formulas.

Definition 16.41.

- 1) $\text{cwf}(\alpha_0) = \{ \ulcorner \varphi \urcorner \in \text{wff}(\alpha_0) \mid (\forall \ulcorner x_i \urcorner, \ulcorner x_j \urcorner \in \text{va}) (\forall \beta \in \alpha_0) (\forall f) [\text{Sqf}(\alpha_0; f) \wedge \ulcorner \varphi \urcorner = \text{end}(f) \wedge [f \cdot 0 = \ulcorner x_i \in x_j \urcorner \vee f \cdot 0 = \ulcorner x_j \in x_i \urcorner \vee f \cdot 0 = \ulcorner x_i \in k_\beta \urcorner \vee f \cdot 0 = \ulcorner k_\beta \in x_i \urcorner] \rightarrow (\exists k \in \mathcal{D}(f)) (\exists \ulcorner \psi \urcorner \in \text{qf}(\alpha_0)) [f \cdot k = \ulcorner (\exists x_i) \psi \urcorner]] \}$.
- 2) $\text{clf}(\alpha_0) = \{ \ulcorner \varphi \urcorner \in \text{wflf}(\alpha_0) \mid (\forall \ulcorner x_i \urcorner, \ulcorner x_j \urcorner \in \text{va}) (\forall \beta \in \alpha_0) (\forall f) [\text{Sqlf}(\alpha; f) \wedge \ulcorner \varphi \urcorner = \text{end}(f) \wedge [f \cdot 0 = \ulcorner x_i \in x_j \urcorner \vee f \cdot 0 = \ulcorner x_j \in x_i \urcorner \vee f \cdot 0 = \ulcorner x_i \in k_\beta \urcorner \vee f \cdot 0 = \ulcorner k_\beta \in x_i \urcorner] \rightarrow (\exists k \in \mathcal{D}(f)) (\exists \beta \in \alpha_0) (\exists \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0)) [f \cdot k = \ulcorner (\exists^\beta x_i) \psi \urcorner]] \}$.

Theorem 16.42. *If $\alpha_0 \subseteq \mathcal{M}$ then*

- 1) $\text{cwf}(\alpha_0) \subseteq \mathcal{M} \wedge [x \in \text{cf}(\alpha_0)] \text{ Abs } \mathcal{M}$.
- 2) $\text{clf}(\alpha_0) \subseteq \mathcal{M} \wedge [x \in \text{clf}(\alpha_0)] \text{ Abs } \mathcal{M}$.

The proof is left to the reader.

Remark. Having defined the class of wffs we turn now to the problem of formalizing the idea of a wff being satisfied. As a first step in this direction we define the notion of a formula being obtained from another by a change of variable.

If φ is a wff then there is an effective procedure for identifying each occurrence of x_i in φ . If at each such occurrence we replace x_i by x_j the result is a wff ψ . We wish to code this into ZF. The result of substituting the j^{th} variable for the i^{th} variable in the formula with Gödel number γ we will denote by $S_j^i \gamma$ which we define inductively on $\text{deg}(\alpha_0; \gamma)$.

Definition 16.43. $(\forall i \in \omega) (\forall j \in \omega)$

- 1) $S_j^i \ulcorner x_i \urcorner = \ulcorner x_j \urcorner, \quad S_j^i \ulcorner x_k \urcorner = \ulcorner x_k \urcorner, \quad k \neq i.$
- 2) $S_j^i \ulcorner k_\alpha \urcorner = \ulcorner k_\alpha \urcorner, \quad \alpha \in \alpha_0.$
- 3) $S_j^i \ulcorner a \in b \urcorner = J \langle S_j^i \ulcorner a \urcorner, S_j^i \ulcorner b \urcorner, 2 \rangle, \quad \ulcorner a \urcorner, \ulcorner b \urcorner \in \text{vc}(\alpha_0).$
- 4) $S_j^i \ulcorner (\exists x_i) \psi \urcorner = J \langle \ulcorner x_j \urcorner, S_j^i \ulcorner \psi \urcorner, 3 \rangle, \quad \ulcorner x_i \urcorner \in \text{va} \wedge \ulcorner \psi \urcorner \in \text{qf}(\alpha_0).$
- 5) $S_j^i \ulcorner \neg \psi \urcorner = J \langle S_j^i \ulcorner \psi \urcorner, 4, 4 \rangle, \quad \ulcorner \psi \urcorner \in \text{qf}(\alpha_0).$
- 6) $S_j^i \ulcorner \psi \wedge \eta \urcorner = J \langle S_j^i \ulcorner \psi \urcorner, S_j^i \ulcorner \eta \urcorner, 5 \rangle, \quad \ulcorner \psi \urcorner, \ulcorner \eta \urcorner \in \text{qf}(\alpha_0).$

Remark. That the conditions of Definition 16.43 uniquely determine $S_j^i \gamma$ we leave as an exercise for the reader. It should be noted that $S_j^i \gamma$ may fail to be in $\text{wff}(\alpha_0)$ even though γ is. For example if

$$\gamma = \ulcorner (\exists x_0) (\exists x_1) [x_0 \in x_1] \urcorner$$

then

$$S_1^0 \gamma = '(\exists x_1) (\exists x_1) [x_1 \in x_1]'.$$

We next introduce the class of formulas of degree smaller than (less than or equal to) n . From this in turn we define satisfaction functions.

Definition 16.44. $\text{Sm}(\alpha_0; n) = \{ '\varphi' \in \text{wff}(\alpha_0) \mid \text{deg}(\alpha_0; '\varphi') \leq n \}$.

Definition 16.45.

$$\begin{aligned} \text{Sat}(g, b, n) &\stackrel{\Delta}{\longleftrightarrow} n \geq 1 \wedge g : (b^\omega \times \text{Sm}(\alpha_0; n)) \rightarrow 2 \\ &\wedge (\forall f \in b^\omega) (\forall i \in \omega) (\forall j \in \omega) [g' \langle f, 'x_i \in x_j' \rangle = 1 \longleftrightarrow f' i \in f' j] \\ &\wedge (\forall f \in b^\omega) (\forall i \in \omega) (\forall \alpha \in \alpha_0) [g' \langle f, 'x_i \in k_\alpha' \rangle = 1 \longleftrightarrow f' i \in F'_a \alpha] \\ &\wedge (\forall f \in b^\omega) (\forall i \in \omega) (\forall \alpha \in \alpha_0) [g' \langle f, 'k_\alpha \in x_i' \rangle = 1 \longleftrightarrow F'_a \alpha \in f' i] \\ &\wedge (\forall f \in b^\omega) (\forall \alpha \in \alpha_0) (\forall \beta \in \alpha_0) [g' \langle f, 'k_\alpha \in k_\beta' \rangle = 1 \longleftrightarrow F'_a \alpha \in F'_a \beta] \\ &\wedge (\forall f \in b^\omega) (\forall '\varphi' \in \text{wff}(\alpha_0)) ['(\exists x_i) \varphi' \in \text{Sm}(\alpha_0; n) \\ &\quad \rightarrow g' \langle f, '(\exists x_i) \varphi' \rangle = 1 \longleftrightarrow (\exists h \in b^\omega) [(\forall j) [j \neq i \rightarrow f' j = h' j] \\ &\quad \wedge g' \langle h, '\varphi' \rangle = 1] \\ &\wedge (\forall f \in b^\omega) (\forall '\varphi' \in \text{wff}(\alpha_0)) [' \neg \varphi' \in \text{Sm}(\alpha_0; n) \rightarrow [g' \langle f, ' \neg \varphi' \rangle = 1 \\ &\quad \longleftrightarrow g' \langle f, '\varphi' \rangle = 0]] \\ &\wedge (\forall f \in b^\omega) (\forall '\varphi' \in \text{wff}(\alpha_0)) (\forall '\psi' \in \text{wff}(\alpha_0)) ['\varphi \wedge \psi' \in \text{Sm}(\alpha_0; n) \\ &\quad \rightarrow [g' \langle f, '\varphi \wedge \psi' \rangle = 1 \longleftrightarrow g' \langle f, '\varphi' \rangle = 1 \wedge g' \langle f, '\psi' \rangle = 1]]. \end{aligned}$$

Theorem 16.46. $(\forall b) (\forall n \geq 1) (\exists! g) \text{Sat}(g, b, n)$.

Proof. (By induction on n) $\text{Sm}(\alpha_0; 1) = \text{pf}(\alpha_0)$. Therefore if

$$\begin{aligned} g = &\{ \langle \langle f, 'x_i \in x_j' \rangle, 1 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge j \in \omega \wedge f' i \in f' j \} \\ &\cup \{ \langle \langle f, 'x_i \in x_j' \rangle, 0 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge j \in \omega \wedge f' i \notin f' j \} \\ &\cup \{ \langle \langle f, 'x_i \in k_\alpha' \rangle, 1 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge \alpha \in \alpha_0 \wedge f' i \in F'_a \alpha \} \\ &\cup \{ \langle \langle f, 'x_i \in k_\alpha' \rangle, 0 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge \alpha \in \alpha_0 \wedge f' i \notin F'_a \alpha \} \\ &\cup \{ \langle \langle f, 'k_\alpha \in x_i' \rangle, 1 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge \alpha \in \alpha_0 \wedge F'_a \alpha \in f' i \} \\ &\cup \{ \langle \langle f, 'k_\alpha \in x_i' \rangle, 0 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge \alpha \in \alpha_0 \wedge F'_a \alpha \notin f' i \} \\ &\cup \{ \langle \langle f, 'k_\alpha \in k_\beta' \rangle, 1 \rangle \mid f \in b^\omega \wedge F'_a \alpha \in F'_a \beta \} \\ &\cup \{ \langle \langle f, 'k_\alpha \in k_\beta' \rangle, 0 \rangle \mid f \in b^\omega \wedge F'_a \alpha \notin F'_a \beta \} \end{aligned}$$

then $\text{Sat}(g, b, n)$.

As our induction hypothesis we assume that $(\exists g_1) \text{Sat}(g_1, b, n)$. If $\ulcorner \varphi \urcorner \in \text{Sm}(\alpha_0; n+1)$ then $\ulcorner \varphi \urcorner \in \text{Sm}(\alpha_0; n)$ or

$$\begin{aligned} & (\exists i \in \omega) (\exists \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n)) [\text{deg}(\alpha_0; \ulcorner \psi \urcorner) = n \wedge \ulcorner \varphi \urcorner = \ulcorner (\exists x_i) \psi \urcorner] \\ & \vee (\exists \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n)) [\ulcorner \varphi \urcorner = \ulcorner \neg \psi \urcorner] \\ & \vee (\exists \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n) (\exists \ulcorner \eta \urcorner \in \text{Sm}(\alpha_0; n)) [\ulcorner \varphi \urcorner = \ulcorner \psi \wedge \eta \urcorner]. \end{aligned}$$

Therefore if

$$\begin{aligned} g = g_1 \cup \{ \langle \langle f, \ulcorner (\exists x_i) \psi \urcorner \rangle, 1 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n) \\ \wedge (\exists h \in b^\omega) [(\forall j) [j \neq i \rightarrow h^j = f^j] \wedge g_1^i \langle h, \ulcorner \psi \urcorner \rangle = 1] \} \\ \cup \{ \langle \langle f, \ulcorner (\exists x_i) \psi \urcorner \rangle, 0 \rangle \mid f \in b^\omega \wedge i \in \omega \wedge \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n) \\ \wedge (\exists h \in b^\omega) [(\forall j) [j \neq i \rightarrow h^j = f^j] \wedge g_1^i \langle h, \ulcorner \psi \urcorner \rangle = 0] \} \\ \cup \{ \langle \langle f, \ulcorner \neg \psi \urcorner \rangle, 1 \rangle \mid f \in b^\omega \wedge \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n) \wedge g_1^i \langle f, \ulcorner \psi \urcorner \rangle = 0 \} \\ \cup \{ \langle \langle f, \ulcorner \neg \psi \urcorner \rangle, 0 \rangle \mid f \in b^\omega \wedge \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n) \wedge g_1^i \langle f, \ulcorner \psi \urcorner \rangle = 1 \} \\ \cup \{ \langle \langle f, \ulcorner \psi \wedge \eta \urcorner \rangle, 1 \rangle \mid f \in b^\omega \wedge \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n) \wedge \ulcorner \eta \urcorner \in \text{Sm}(\alpha_0; n) \\ \wedge g_1^i \langle f, \ulcorner \psi \urcorner \rangle = 1 \wedge g_1^i \langle f, \ulcorner \eta \urcorner \rangle = 1 \} \\ \cup \{ \langle \langle f, \ulcorner \psi \wedge \eta \urcorner \rangle, 0 \rangle \mid f \in b^\omega \wedge \ulcorner \psi \urcorner \in \text{Sm}(\alpha_0; n) \wedge \ulcorner \eta \urcorner \in \text{Sm}(\alpha_0; n) \\ \wedge [g_1^i \langle f, \ulcorner \psi \urcorner \rangle = 0 \vee g_1^i \langle f, \ulcorner \eta \urcorner \rangle = 0] \}. \end{aligned}$$

Then $\text{Sat}(g, b, n+1)$.

It is easily proved by induction that

$$\text{Sat}(g_1, b, n) \wedge \text{Sat}(g_2, b, n) \rightarrow g_1 = g_2.$$

Details are left to the reader.

Definition 16.47

- 1) If $f \in b^\omega \wedge \ulcorner \varphi \urcorner \in \text{wff}(\alpha_0)$ then
 $b, f \models \ulcorner \varphi \urcorner \iff (\exists g) [\text{Sat}(g, b, \text{deg}(\alpha_0; \ulcorner \varphi \urcorner)) \wedge g^i \langle f, \ulcorner \varphi \urcorner \rangle = 1].$
- 2) If $\ulcorner \varphi \urcorner \in \text{cf}(\alpha_0)$ then
 $b \models \ulcorner \varphi \urcorner \iff (\exists f \in b^\omega) [b, f \models \ulcorner \varphi \urcorner].$

Theorem 16.48. If $f \in b^\omega \wedge \ulcorner \psi \urcorner, \ulcorner \eta \urcorner \in \text{wff}(\alpha_0)$ then

- 1) $b, f \models \ulcorner x_i \in x_j \urcorner \iff f^i \in f^j.$
- 2) $b, f \models \ulcorner x_i \in k_\alpha \urcorner \iff f^i \in F_\alpha^i.$
- 3) $b, f \models \ulcorner k_\alpha \in x_i \urcorner \iff F_\alpha^i \in f^i.$
- 4) $b, f \models \ulcorner k_\alpha \in k_\beta \urcorner \iff F_\alpha^i \in F_\beta^i.$
- 5) $b, f \models \ulcorner (\exists x_i) \psi \urcorner \iff (\exists h \in b^\omega) [(\forall j) [j \neq i \rightarrow h^j = f^j] \wedge b, h \models \ulcorner \psi \urcorner].$

$$6) \quad b, f \models \ulcorner \neg \psi \urcorner \longleftrightarrow \neg [b, f \models \ulcorner \psi \urcorner].$$

$$7) \quad b, f \models \ulcorner \psi \wedge \eta \urcorner \longleftrightarrow [b, f \models \ulcorner \psi \urcorner] \wedge [b, f \models \ulcorner \eta \urcorner].$$

Proof. Left to the reader.

Theorem 16.49. *If $\varphi(x_{i_1}, \dots, x_{i_n})$ has no free variables other than x_{i_1}, \dots, x_{i_n} and if $f \in b^\omega$ then*

$$b, f \models \varphi(x_{i_1}, \dots, x_{i_n}) \longleftrightarrow \varphi^b(f^{i_1}, \dots, f^{i_n}).$$

Proof. (By induction on m the number of logical symbols in φ .) If $m = 0$ then φ is a prime formula and the result follows from Theorem 16.48. If $m > 0$ then φ is of the form $(\exists x_{i_0}) \psi(x_{i_0}, x_{i_1}, \dots, x_{i_n})$ or $\psi(x_{i_1}, \dots, x_{i_n})$ or $\psi(x_{i_1}, \dots, x_{i_n}) \wedge \eta(x_{i_1}, \dots, x_{i_n})$ where ψ and η have no free variables other than those indicated.

If φ is $(\exists x_{i_0}) \psi(x_{i_0}, \dots, x_{i_n})$ then by Definitions 16.45 and 16.47

$$b, f \models \ulcorner \varphi \urcorner \longleftrightarrow (\exists h \in b^\omega) [(\forall j) [j \neq i_0 \rightarrow h^j = f^j] \wedge [b, h \models \ulcorner \psi \urcorner]].$$

Since ψ has one less logical symbol than φ , we have from the induction hypothesis

$$b, h \models \ulcorner \psi(x_{i_0}, x_{i_1}, \dots, x_{i_n}) \urcorner \longleftrightarrow \psi^b(h^{i_0}, h^{i_1}, \dots, h^{i_n}).$$

Since $h^{i_0} \in b$ and $h^j = f^j$ for $j \neq i_0$

$$b, f \models \ulcorner \varphi \urcorner \longleftrightarrow (\exists x_{i_0} \in b) \psi^b(x_{i_0}, f^{i_1}, \dots, f^{i_n}).$$

If φ is $\neg \psi(x_{i_1}, \dots, x_{i_n})$ then $b, f \models \ulcorner \psi(x_{i_1}, \dots, x_{i_n}) \urcorner \longleftrightarrow \psi^b(f^{i_1}, \dots, f^{i_n})$ and hence

$$b, f \models \ulcorner \varphi \urcorner \longleftrightarrow \neg [b, f \models \ulcorner \psi(x_{i_1}, \dots, x_{i_n}) \urcorner] \longleftrightarrow \neg \psi^b(f^{i_1}, \dots, f^{i_n}).$$

If φ is $\psi(x_{i_1}, \dots, x_{i_n}) \wedge \eta(x_{i_1}, \dots, x_{i_n})$ then

$$b, f \models \ulcorner \psi(x_{i_1}, \dots, x_{i_n}) \urcorner \longleftrightarrow \psi^b(f^{i_1}, \dots, f^{i_n})$$

$$b, f \models \ulcorner \eta(x_{i_1}, \dots, x_{i_n}) \urcorner \longleftrightarrow \eta^b(f^{i_1}, \dots, f^{i_n})$$

and hence

$$\begin{aligned} b, f \models \ulcorner \varphi \urcorner &\longleftrightarrow b, f \models \ulcorner \psi \urcorner \wedge b, f \models \ulcorner \eta \urcorner \\ &\longleftrightarrow \psi^b(f^{i_1}, \dots, f^{i_n}) \wedge \eta^b(f^{i_1}, \dots, f^{i_n}). \end{aligned}$$

Corollary 16.50. $\ulcorner \varphi \urcorner \in \text{cf}(\alpha_0) \rightarrow [b \models \ulcorner \varphi \urcorner \longleftrightarrow \varphi^b]$.

Remark. With Definition 16.47 we can express “ b is a model of ZF” as a single sentence in ZF. This we do by saying essentially

$$(\forall \alpha) [\alpha \text{ is an axiom} \rightarrow b \models \alpha].$$

It only remains to formalize “ α is an axiom”. The problem is of course Axiom 5 which is an axiom schema.

In order to formalize “ α is an instance of Axiom 5” we must define the class of Gödel numbers of instances of Axiom 5. This in turn requires that we describe the class of input formulas $\varphi(x_1, x_2)$.

Definition 16.51.

$$\begin{aligned} \text{wff}(1, 2; 0, 3) \stackrel{\Delta}{=} & \{ \varphi \in \text{wff}(\alpha_0) \mid \neg O(\alpha_0; \ulcorner x_0 \urcorner, \ulcorner \varphi \urcorner) \wedge \neg \text{Ob}(\alpha_0; \ulcorner x_0 \urcorner, \ulcorner \varphi \urcorner) \\ & \wedge \neg \text{Ob}(\alpha_0; \ulcorner x_1 \urcorner, \ulcorner \varphi \urcorner) \wedge \neg \text{Ob}(\alpha_0; \ulcorner x_2 \urcorner, \ulcorner \varphi \urcorner) \wedge \neg \text{Ob}(\alpha_0; \ulcorner x_3 \urcorner, \ulcorner \varphi \urcorner) \\ & \wedge \neg O(\alpha_0; \ulcorner x_3 \urcorner, \ulcorner \varphi \urcorner) \}. \end{aligned}$$

Remark. Definition 16.51 describe the admissible φ 's for Axiom 5. In order to code Axiom 5 into ZF we must express the axiom in primitive terms. It is clear that, using Definition 16.43, this can be done and that a formula for Gödel numbers of such formulas can be defined.

We next define the class of Gödel numbers of instances of Axiom 5.

Definition 16.52.

$$\begin{aligned} \text{ax}_5 \stackrel{\Delta}{=} & \{ \ulcorner \varphi \urcorner \in \text{cf}(\alpha_0) \mid (\exists \ulcorner \psi(x_1, x_2) \urcorner \in \text{wff}(1, 2; 0, 3)) \\ & [\ulcorner \varphi \urcorner = \ulcorner (\forall x_0) [(\forall x_1) (\forall x_2) (\forall x_3) \psi(x_1, x_2) \wedge \psi(x_1, x_3) \rightarrow x_2 = x_3] \\ & \rightarrow (\exists x_1) (\forall x_2) [x_2 \in x_1 \leftrightarrow (\exists x_3) [x_3 \in x_0 \wedge \psi(x_3, x_2)]]] \}. \end{aligned}$$

Definition 16.53.

- 1) $\text{SM}(b, \text{ZF}) \stackrel{\Delta}{\leftrightarrow} b \neq 0 \wedge (\text{Ax. 1})^b \wedge (\text{Ax. 2})^b \wedge (\text{Ax. 3})^b \wedge (\text{Ax. 4})^b$
 $\wedge (\text{Ax. 6})^b \wedge (\text{Ax. 7})^b \wedge (\forall \ulcorner \varphi \urcorner \in \text{ax}_5) [b \models \ulcorner \varphi \urcorner]$.
- 2) $\text{STM}(b, \text{ZF}) \stackrel{\Delta}{\leftrightarrow} \text{SM}(b, \text{ZF}) \wedge \text{Tr}(b)$.

17 Cohen's Method

In proving that the AC and the GCH are consistent with ZF Gödel used the so called method of internal models. From the assumption that the universe V is a model of ZF Gödel prescribed a method for producing a submodel L that is also a model of $V = L$, AC and GCH. This submodel is defined as the class of all sets having a certain property i.e.

$$L = \{a | (\exists \alpha) [a = F^{\alpha}]\} .$$

Indeed since $a = F^{\alpha}$ is absolute w.r.t. every standard transitive model \mathcal{M} it follows that if

$$L^{\mathcal{M}} = \{a | (\exists \alpha \in \mathcal{M}) [a = F^{\alpha}]\}$$

then $L^{\mathcal{M}}$ is a submodel of \mathcal{M} that is also a model of $V = L$.

If $V = L$ is valid in every model then $V = L$ must be provable in ZF and conversely if $V = L$ is not provable in ZF then $V = L$ is not valid in some model. Can we hope to find such a model by the method of internal models? That is, can we hope to produce a property $\varphi(a)$ such that

$$\{a | \varphi(a)\}$$

is a model of $ZF + V \neq L$? There are compelling reasons for believing that this method cannot succeed. The arguments turn upon the assumption that there is a set that is a standard model of ZF.

Theorem 17.1. *If there exists a set that is a standard model of ZF then there exists one and only one set \mathcal{M}_0 such that*

- 1) \mathcal{M}_0 is a countable standard transitive model of $ZF + V = L$ and
- 2) \mathcal{M}_0 is a submodel of every standard transitive model of ZF.

Proof. From Mostowski's theorem (Theorem 12.8) every standard model is ϵ -isomorphic to a standard transitive model. Therefore the existence of a set that is a standard model of ZF implies the existence of a set that is a standard transitive model. For transitive models the property of being an ordinal is absolute. That is, those sets in a transitive model that play the role of ordinals are ordinals. Furthermore from transitivity if α is in the model then all smaller ordinals are in the model. But a standard transitive model that is a set cannot contain all ordinals.

If α is the smallest ordinal not contained in such a model then α is the class of ordinals for that model. But the existence of such an ordinal implies the existence of a smallest such ordinal, α_0 , that is the set of all ordinals in some standard transitive model N_0 .

Since N_0 is a model of ZF it follows that if

$$\mathcal{M}_0 \triangleq L^{N_0} = \{a \mid (\exists \alpha \in N_0) [a = F'\alpha]\} = \{F'\alpha \mid \alpha < \alpha_0\}$$

then \mathcal{M}_0 is a model of $ZF + V = L$.

If N is any standard transitive model of ZF then N is closed w.r.t. the fundamental operations. Therefore since $\alpha_0 \subseteq N$ it follows that $\mathcal{M}_0 \subseteq N$. From this we see that \mathcal{M}_0 is unique, for if \mathcal{M}_0 and \mathcal{M}' are each standard transitive models with the prescribed properties then

$$\mathcal{M}_0 \subseteq \mathcal{M}' \quad \text{and} \quad \mathcal{M}' \subseteq \mathcal{M}_0.$$

Finally, from the Lowenheim-Skolem Theorem, \mathcal{M}_0 contains a countable standard submodel. But this submodel must be ϵ -isomorphic to a countable standard transitive model that must contain \mathcal{M}_0 as a submodel. Hence \mathcal{M}_0 is countable.

Remark. The unique model \mathcal{M}_0 described in Theorem 17.1 is called the *minimal model*. Its existence follows from the existence of a set that is a standard model. It should be observed that the minimal model contains no proper transitive submodel. Thus the existence of a model of ZF does not imply the existence of a standard model. (See Appendix.) We clearly cannot prove the existence of a standard model for that would prove the consistency of ZF. We therefore postulate the existence of such a model

Standard Model Hypothesis: $(\exists m) \text{SM}(m, \text{ZF})$.

From this assumption and Theorem 17.1 we are assured of the existence of the minimal model \mathcal{M}_0 that is 1) countable, hence contains a countable collection of ordinals α_0 , 2) a model of $ZF + V = L$, and 3) a submodel of every standard transitive model. Indeed from Mostowski's theorem every standard model contains a submodel ϵ -isomorphic to \mathcal{M}_0 .

From the existence of the minimal model it follows that an attempt to prove $V = L$ and ZF independent by the method of internal models is doomed to failure. Suppose that we could produce a wff $\varphi(a)$ for which it is provable in ZF that

$$\{a \mid \varphi(a)\}$$

is a model of $ZF + V \neq L$. It then follows that this theorem relativized to \mathcal{M}_0 is also a theorem. That is

$$\{a \in \mathcal{M}_0 \mid \varphi^{\mathcal{M}_0}(a)\}$$

is a submodel of \mathcal{M}_0 that is also a model of $ZF + V \neq L$. Since \mathcal{M}_0 is a model of $V = L$ this submodel is a proper submodel of \mathcal{M}_0 . But such a submodel of \mathcal{M}_0 must be isomorphic to a standard transitive proper submodel of \mathcal{M}_0 that must in turn contain \mathcal{M}_0 as a submodel. This is impossible.

The independence of $V = L$ must then be established by some method other than that of internal models. Cohen's approach is to extend the minimal model \mathcal{M}_0 by adjoining a subset of the integers that is not in \mathcal{M}_0 and then closing w.r.t. the fundamental operations. For this purpose we introduced the notion of relative constructibility.

In Section 15 we defined the functions $F_a, a \subseteq \omega$ in such a way that

$$\begin{aligned} F_a' \alpha &= \alpha & \alpha \leq \omega \\ &= a & \alpha = \omega + 1 \\ &= F_a'' \alpha & \alpha > \omega + 1 \wedge K_3' \alpha = 0 \\ &= \mathcal{F}_i(F_a' K_1' \alpha, F_a' K_2' \alpha), & \alpha > \omega + 1 \wedge 0 < i = K_3' \alpha. \end{aligned}$$

From each F_a we defined a class L_a that we proved to be a model of $ZF + AC + GCH + V = L_a$. We also proved that $L \subseteq L_a$.

L_a is the class obtained from L by adjoining a , in the sense that

$$L_a = \{x \mid (\exists \alpha \in L) [x = F_a' \alpha]\}.$$

We wish to consider the class obtained from the minimal model \mathcal{M}_0 by adjoining a .

Definition 17.2. $N_a \triangleq \{x \mid (\exists \alpha \in \mathcal{M}_0) [x = F_a' \alpha]\}, \quad a \subseteq \omega.$

Since α_0 is the set of ordinals in \mathcal{M}_0 it follows that $N_a = F_a'' \alpha_0$. But by Theorem 15.21, $F_a'' \alpha_0$ is transitive and, since $\alpha_0 > 0$, $F_a'' \alpha_0$ is nonempty. We have therefore proved.

Theorem 17.3. $\forall a \subseteq \omega, N_a$ is nonempty and transitive.

Theorem 17.4. $\forall a \subseteq \omega, N_a$ is closed with respect to the fundamental operations.

Proof.

$$x \in N_a \wedge y \in N_a \rightarrow (\exists \alpha, \beta \in \alpha_0) [\alpha > \omega \wedge \beta > \omega \wedge x = F_a' \alpha \wedge y = F_a' \beta].$$

If $\gamma = J' \langle \alpha, \beta, i \rangle$ then $\omega + 1 < \gamma$ and, since α_0 is closed w.r.t. J, K_1, K_2 , and K_3 , $\gamma \in \alpha_0$. Hence

$$\mathcal{F}_i(x, y) = \mathcal{F}_i(F_a' \alpha, F_a' \beta) = F_a' \gamma \in N_a.$$

Definition 17.5. $\text{cov}(\alpha) = \mu_\beta(\beta > \alpha \wedge K_3' \beta = 0).$

Theorem 17.6.

- 1) $\text{cov}(\alpha) > \alpha \wedge K_3 \text{cov}(\alpha) = 0.$
- 2) $\alpha \in \alpha_0 \rightarrow \text{cov}(\alpha) \in \alpha_0.$

The proofs are left to the reader.

Theorem 17.7. *For each $a \subseteq \omega$, if N_a is a standard transitive model of the Axiom of Powers and the Axiom Schema of Replacement then N_a is a standard transitive model of ZF.*

Proof. Since N_a is nonempty and transitive it follows from Theorems 13.24 and 13.35 that N_a is a standard transitive model of the Axiom of Extensionality (Axiom 1) and the Axiom of Regularity (Axiom 6).

From Theorem 17.4

$$x \in N_a \wedge y \in N_a \rightarrow \mathcal{F}_1(x, y) = \{x, y\} \in N_a.$$

Therefore, by Theorem 13.25, N_a is a model of the Axiom of Pairing (Axiom 2).

$$(\forall x \in N_a) (\exists \alpha \in \alpha_0) [x = F_a^* \alpha].$$

If $\gamma = \text{cov}(\alpha)$ then $\gamma > \alpha \wedge K_3 \gamma = 0$. Consequently

$$F_a^* \alpha \subseteq F_a^{**} \alpha \subseteq F_a^{**} \gamma = F_a^* \gamma.$$

Then

$$z \in y \wedge y \in x \rightarrow z \in F_a^* \gamma.$$

Since by hypothesis N_a is a model of the Axiom Schema of Replacement

$$\{z \in F_a^* \gamma \mid (\exists y \in x) [z \in y]\} \in N_a.$$

Therefore N_a is a model of the Axiom of Unions (Axiom 3).

Finally since

$$F_a^* \omega = \omega \in N_a$$

we have from Theorem 13.38 that N_a is a model of the Axiom of Infinity (Axiom 7).

Remark. Suppose that it could be shown that for some $a \subseteq \omega$, $a \notin \mathcal{M}_0$ and N_a is a standard transitive model of the Axiom of Powers and the Axiom Schema of Replacement. It would then follow from Theorem 17.7 that N_a is a standard transitive model of ZF. Suppose, in addition, it can be shown that

$$\alpha \in N_a \rightarrow \alpha \in \alpha_0.$$

It would then follow, since the minimal model \mathcal{M}_0 is a submodel of every standard transitive model of ZF, that α_0 is the class of ordinals in N_a . Then

$$\begin{aligned} N_a &= \{x \mid (\exists \alpha \in \alpha_0) [x = F_a^* \alpha]\} = L_a^{N_a}, \\ \mathcal{M}_0 &= \{x \mid (\exists \alpha \in \alpha_0) [x = F^* \alpha]\} = L^{N_a}. \end{aligned}$$

Consequently N_a is a standard transitive model of $ZF + AC + GCH$. Furthermore since $a \in N_a \wedge a \notin \mathcal{M}_0$, N_a is also a model of $V \neq L$. This would establish the independence of the Axiom of Constructibility and the axioms of $ZF + AC + GCH$.

To prove that $\alpha \in N_a \rightarrow \alpha \in \alpha_0$ is easy. We do that immediately. To prove that for some $a \subseteq \omega$, $a \notin \mathcal{M}_0 \wedge N_a$ is a model of the Axiom of Powers and the Axiom Schema of Replacement requires work that will not be completed until near the end of the next section.

Theorem 17.8. $\text{rank}(F_a^{\epsilon}\alpha) \leq \alpha$.

Proof (by induction). Since $F_a^{\epsilon}\alpha \subseteq F_a^{\epsilon\epsilon}\alpha$

$$\text{rank}(F_a^{\epsilon}\alpha) \leq \text{rank}(F_a^{\epsilon\epsilon}\alpha).$$

But from properties of rank and the induction hypothesis

$$\text{rank}(F_a^{\epsilon\epsilon}\alpha) = \mu_{\gamma}((\forall \beta < \alpha) [\text{rank}(F_a^{\epsilon}\beta) < \gamma]) \leq \mu_{\gamma}((\forall \beta < \alpha) [\beta < \gamma]) = \alpha.$$

Theorem 17.9. $(\forall a \subseteq \omega) [\alpha \in N_a \rightarrow \alpha \in \mathcal{M}_0]$.

Proof. $\alpha \in N_a \rightarrow (\exists \beta \in \alpha_0) [\alpha = F_a^{\epsilon}\beta]$.

But since $\alpha = \text{rank}(\alpha)$ we have from Theorem 17.8

$$\alpha = \text{rank}(F_a^{\epsilon}\beta) \leq \beta < \alpha_0.$$

To prove the existence of a set a for which N_a is a model of ZF we will develop a theory of N_a -models using the language $ZF\{\alpha_0\}$. Our principal interpretation for the constants in $ZF\{\alpha_0\}$ will be that

$$(\forall \alpha \in \alpha_0) [k_{\alpha} = F_a^{\epsilon}\alpha].$$

In view of this we will not always distinguish between k_{α} and $F_a^{\epsilon}\alpha$ in the discussions to follow.

In order for N_a to be a model of ZF it is necessary that N_a satisfy the axioms of ZF. We wish to know what restricts on a will insure this. From the definition of satisfaction (§ 16) and our principal interpretation we wish the following to hold:

$$N_a \models \text{'}\psi \wedge \eta\text{' } \longleftrightarrow [N_a \models \text{'}\psi\text{'}] \wedge [N_a \models \text{'}\eta\text{'}],$$

$$N_a \models \text{'}\neg \psi\text{' } \longleftrightarrow \neg [N_a \models \text{'}\psi\text{'}],$$

$$N_a \models \text{'}(\exists x) \psi(x)\text{' } \longleftrightarrow (\exists \alpha \in \alpha_0) [N_a \models \text{'}\psi(k_{\alpha})\text{' }].$$

By repeated application of these formulas we see that the question of whether or not N_a satisfies a given wff φ reduces to the question of whether or not N_a satisfies a certain collection of formulas of the type $k_{\alpha} \in k_{\beta}$. The basic task of our theory of N_a -models is, therefore, to determine the \in -connections between the constants in the language $ZF\{\alpha_0\}$ that will assure us that the class of all constants is a model of ZF. Since

in our principal interpretation $k_{\omega+1}$ is a , this will in turn impose restrictions on a .

The proper ε -connections we will determine in the following way. Note that

$$N_a \models ' \varphi ' .$$

is a predicate involving a . Any set is determined by its extent. It may however be the case that the truth or falsehood of the predicate $N_a \models ' \varphi '$ can be decided without a full knowledge of the extent of a . That is, if

$$a = \{m_0, m_1, \dots\} \quad \text{and} \quad \omega - a = \{n_0, n_1, \dots\}$$

it may be that $N_a \models \varphi$ can be expressed as a predicate

$$P(m_0, \dots, m_i; n_0, \dots, n_j; ' \varphi ')$$

where i and j depend on $' \varphi '$. Since $a \subseteq \mathcal{M}_0$ and $' \varphi ' \in \mathcal{M}_0$ it may also be the case that the truth or falsehood of the predicate $P(m_0, \dots, m_i; n_0, \dots, n_j; ' \varphi ')$ can be decided from our knowledge that \mathcal{M}_0 is a standard transitive model of ZF. For this it is essential that the predicate $P(m_0, \dots, m_i; n_0, \dots, n_j; ' \varphi ')$ be \mathcal{M}_0 -definable in the following sense.

Definition 17.10. A wff $\varphi(x_1, \dots, x_n)$ having no free variables other than x_1, \dots, x_n is \mathcal{M}_0 -definable iff there exists a wff $\psi(x_1, \dots, x_n)$ for which

$$x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow [\varphi(x_1, \dots, x_n) \longleftrightarrow \psi^{\mathcal{M}_0}(x_1, \dots, x_n)] .$$

Theorem 17.11. If φ and ψ are each \mathcal{M}_0 -definable then $\varphi \wedge \psi$, $\varphi \vee \psi$, $\neg \varphi$, $\varphi \rightarrow \psi$, and $\varphi \longleftrightarrow \psi$ are each \mathcal{M}_0 -definable.

Proof. If $\varphi(x_1, \dots, x_n)$ and $\psi(x_1, \dots, x_n)$ have no free variables other than those listed and if φ and ψ are each \mathcal{M}_0 -definable then there are wffs $\eta_1(x_1, \dots, x_n)$ and $\eta_2(x_1, \dots, x_n)$ for which

$$x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow [\varphi(x_1, \dots, x_n) \longleftrightarrow \eta_1^{\mathcal{M}_0}(x_1, \dots, x_n)] ,$$

$$x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow [\psi(x_1, \dots, x_n) \longleftrightarrow \eta_2^{\mathcal{M}_0}(x_1, \dots, x_n)] .$$

Then

$$\begin{aligned} x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow & [\varphi(x_1, \dots, x_n) \wedge \psi(x_1, \dots, x_n) \\ & \longleftrightarrow \eta_1^{\mathcal{M}_0}(x_1, \dots, x_n) \wedge \eta_2^{\mathcal{M}_0}(x_1, \dots, x_n)] , \end{aligned}$$

$$\begin{aligned} x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow & [\varphi(x_1, \dots, x_n) \vee \psi(x_1, \dots, x_n) \\ & \longleftrightarrow \eta_1^{\mathcal{M}_0}(x_1, \dots, x_n) \vee \eta_2^{\mathcal{M}_0}(x_1, \dots, x_n)] , \end{aligned}$$

$$x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow [\neg \varphi(x_1, \dots, x_n) \longleftrightarrow \neg \eta_1^{\mathcal{M}_0}(x_1, \dots, x_n)] ,$$

$$\begin{aligned} x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow & [[\varphi(x_1, \dots, x_n) \rightarrow \psi(x_1, \dots, x_n)] \\ & \longleftrightarrow [\eta_1^{\mathcal{M}_0}(x_1, \dots, x_n) \rightarrow \eta_2^{\mathcal{M}_0}(x_1, \dots, x_n)]] , \end{aligned}$$

$$\begin{aligned} x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow & [[\varphi(x_1, \dots, x_n) \longleftrightarrow \psi(x_1, \dots, x_n)] \\ & \longleftrightarrow [\eta_1^{\mathcal{M}_0}(x_1, \dots, x_n) \longleftrightarrow \eta_2^{\mathcal{M}_0}(x_1, \dots, x_n)]] . \end{aligned}$$

Theorem 17.12. $\varphi \text{ Abs } \mathcal{M}_0$ implies that φ is \mathcal{M}_0 -definable.

Proof. If $\varphi(x_1, \dots, x_n)$ has no free variables other than those indicated then since φ is absolute w.r.t. \mathcal{M}_0

$$x_1 \in \mathcal{M}_0 \wedge \dots \wedge x_n \in \mathcal{M}_0 \rightarrow [\varphi(x_1, \dots, x_n) \leftrightarrow \varphi^{\mathcal{M}_0}(x_1, \dots, x_n)].$$

Theorem 17.13. $\beta = 'k_\alpha'$ is \mathcal{M}_0 -definable.

Proof. $\beta = 'k_\alpha' \leftrightarrow \beta = J\langle \alpha, 1, 1 \rangle$. But $x = J\langle \alpha, 1, 1 \rangle \text{ Abs } \mathcal{M}_0$.

Remark. We next wish to show that $x \in ic(\alpha_0)$, $x \in pf(\alpha_0)$, $x \in qf(\alpha_0)$, etc. are each expressible in the minimal model. To do this we re-formulate certain definitions of § 16 replacing α_0 by On. For example

$$ic \triangleq \{\gamma \mid (\exists \alpha) [\gamma = J\langle \alpha, 1, 1 \rangle]\}$$

$$vc \triangleq va \cup ic$$

$$pf \triangleq \{\gamma \mid (\exists \alpha, \beta \in vc) [\gamma = J\langle \alpha, \beta, 2 \rangle]\}$$

Similarly, we define qf , qlf , wff , $wflf$, cf , and clf as we defined $qf(\alpha_0)$, $qlf(\alpha_0)$, $wff(\alpha_0)$, $wflf(\alpha_0)$, $cf(\alpha_0)$, and $clf(\alpha_0)$ except that α_0 is replaced by On. From these definitions the following theorems are immediate.

Theorem 17.14. $x \in vc(\alpha_0) \leftrightarrow [x \in vc]^{\mathcal{M}_0}$.

Theorem 17.15. 1) $x \in pf(\alpha_0) \leftrightarrow [x \in pf]^{\mathcal{M}_0}$

2) $x \in qf(\alpha_0) \leftrightarrow [x \in qf]^{\mathcal{M}_0}$ 3) $x \in qlf(\alpha_0) \leftrightarrow [x \in qlf]^{\mathcal{M}_0}$

4) $x \in wff(\alpha_0) \leftrightarrow [x \in wff]^{\mathcal{M}_0}$ 5) $x \in wflf(\alpha_0) \leftrightarrow [x \in wflf]^{\mathcal{M}_0}$

6) $x \in cf(\alpha_0) \leftrightarrow [x \in cf]^{\mathcal{M}_0}$ 7) $x \in clf(\alpha_0) \leftrightarrow [x \in clf]^{\mathcal{M}_0}$

18 Forcing

We turn now to the task of expressing

$$N_a \models \ulcorner \varphi \urcorner$$

as an \mathcal{M}_0 -definable predicate

$$P(m_0, \dots, m_i; n_0, \dots, n_j; \ulcorner \varphi \urcorner)$$

where i and j depend upon $\ulcorner \varphi \urcorner$, $a = \{m_0, m_1, \dots\}$ and $\omega - a = \{n_0, n_1, \dots\}$. Such a predicate will be defined in this section. When this predicate holds we say that $\langle \{m_0, \dots, m_i\}, \{n_0, \dots, n_j\} \rangle$ forces $\ulcorner \varphi \urcorner$. The ordered pair $\langle \{m_0, \dots, m_i\}, \{n_0, \dots, n_j\} \rangle$ is called a forcing condition.

Forcing will be defined inductively. For its definition we must distinguish between the limited formulas of ZF $\{\alpha_0\}$, defined in § 16, and the unlimited formula that we will now define. Here α_0 is the smallest ordinal not in the minimal model \mathcal{M}_0 .

Definition 18.1.

- 1) $\text{ulf}(\alpha_0) = \text{wff}(\alpha_0) - \text{wflf}(\alpha_0)$.
- 2) $\text{culf}(\alpha_0) = \text{cwf}(\alpha_0) - \text{clf}(\alpha_0)$.
- 3) $s(\alpha_0) = \text{culf}(\alpha_0) \cup \text{clf}(\alpha_0)$.

Remark. $s(\alpha_0)$ is the class of statements of ZF $\{\alpha_0\}$. It is partitioned into the class of closed unlimited formulas, $\text{culf}(\alpha_0)$, and the class of closed limited formulas, $\text{clf}(\alpha_0)$. On $s(\alpha_0)$ we will define a well founded relation for which we need the notion of type and parity:

Definition 18.2. $(\forall \ulcorner \varphi \urcorner \in \text{clf}(\alpha_0)) [\text{typ}(\alpha_0; \ulcorner \varphi \urcorner)$

$$= \mu_\beta((\forall \gamma \in \alpha_0) (\forall \ulcorner x_i \urcorner \in \text{va})) (\forall \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0)) [\text{Subqlf}(\alpha_0; \ulcorner (\exists^\gamma x_i) \psi \urcorner, \ulcorner \varphi \urcorner) \rightarrow \gamma \leq \beta] \\ \wedge (\forall \ulcorner x_i \urcorner \in \text{va}) (\forall \gamma, \delta \in \alpha_0) [\text{Subqlf}(\alpha_0; \ulcorner x_i \in k_\delta \urcorner, \ulcorner \varphi \urcorner) \vee \text{Subqlf}(\alpha_0; \ulcorner k_\gamma \in x_i \urcorner, \ulcorner \varphi \urcorner) \\ \vee \text{Subqlf}(\alpha_0; \ulcorner k_\gamma \in k_\delta \urcorner, \ulcorner \varphi \urcorner) \rightarrow \gamma < \beta \wedge \delta < \beta]]].$$

Remark. To determine the type of a given closed limited formula φ we examine φ for occurrences of bound quantifiers, \exists^{β_1} , \exists^{β_2} , etc., and for occurrences of individual constants k_{α_1} , k_{α_2} , etc. The type of φ is then the maximum ordinal in the finite collection $\beta_1, \beta_2, \dots, \alpha_1 + 1, \alpha_2 + 1, \dots$.

Definition 18.3. $\forall \ulcorner \varphi \urcorner \in \text{clf}(\alpha_0)$

$$\text{par}(\alpha_0; \ulcorner \varphi \urcorner) = 0 \iff (\forall \beta \in \alpha_0) (\forall \ulcorner x_i \urcorner \in \text{va}) (\forall \ulcorner \psi \urcorner \in \text{qlf}(\alpha_0))$$

$$[\text{Subqlf}(\alpha_0; \ulcorner (\exists^\beta x_i) \psi \urcorner, \ulcorner \varphi \urcorner) \rightarrow \beta < \text{typ}(\alpha_0; \ulcorner \varphi \urcorner)]$$

$$\wedge (\forall \beta \in \alpha_0) (\forall \ulcorner a \urcorner \in \text{vc}(\alpha_0)) [\text{Subqlf}(\alpha_0, \ulcorner k_\beta \in a \urcorner, \ulcorner \varphi \urcorner) \rightarrow \beta + 1 < \text{typ}(\alpha_0; \ulcorner \varphi \urcorner)].$$

$$\text{par}(\alpha_0; \ulcorner \varphi \urcorner) = 1 \iff \neg [\text{par}(\alpha_0; \ulcorner \varphi \urcorner) = 0].$$

Remark. We next define a partial ordering on $s(\alpha_0)$.

Definition 18.4. $\alpha < \beta \stackrel{\Delta}{\iff} [\alpha, \beta \in \text{culf}(\alpha_0) \wedge \text{deg}(\alpha_0, \alpha) < \text{deg}(\alpha_0, \beta)]$

$$\vee [\alpha \in \text{clf}(\alpha_0) \wedge \beta \in \text{culf}(\alpha_0)] \vee [\alpha, \beta \in \text{clf}(\alpha_0) \wedge [[\text{typ}(\alpha_0, \alpha) < \text{typ}(\alpha_0; \beta)]$$

$$\vee [[\text{typ}(\alpha_0; \alpha) = \text{typ}(\alpha_0; \beta)] \wedge [[\text{par}(\alpha_0, \alpha) < \text{par}(\alpha_0, \beta)]$$

$$\vee [[\text{par}(\alpha_0, \alpha) = \text{par}(\alpha_0, \beta)] \wedge [\text{deg}_l(\alpha_0; \alpha) < \text{deg}_l(\alpha_0, \beta)]]]]].$$

Theorem 18.5.

$$1) \quad s(\alpha_0) \subseteq \mathcal{M}_0.$$

$$2) \quad < \text{ is a well founded relation on } s(\alpha_0).$$

The proof is left to the reader.

Remark. We will define forcing inductively on the relation $<$. For the definition of forcing we first define the class of forcing conditions.

Definition 18.6.

$$\text{cond} = \{ \langle p_1, p_2 \rangle \mid p_1 \subset \omega \wedge p_2 \subset \omega \wedge \text{Fin}(p_1) \wedge \text{Fin}(p_2) \wedge p_1 \cap p_2 = 0 \}.$$

Remark. We will use $\mathcal{P}, \mathcal{P}', \mathcal{P}''$ as variables on forcing conditions.

Definition 18.7. Let $\mathcal{P} = \langle p_1, p_2 \rangle \wedge \mathcal{P}' = \langle p'_1, p'_2 \rangle$ then

$$\mathcal{P} \subseteq \mathcal{P}' \iff p_1 \subseteq p'_1 \wedge p_2 \subseteq p'_2.$$

Theorem 18.8.

$$1) \quad [x \in \text{cond}] \text{ Abs } \mathcal{M}_0.$$

$$2) \quad [\mathcal{P} \subseteq \mathcal{P}'] \text{ Abs } \mathcal{M}_0.$$

$$3) \quad \text{cond} \subseteq \mathcal{M}_0.$$

Proof.

$$1) \quad x \in \text{cond} \iff (\exists p_1) (\exists p_2) [p_1 \subset \omega \wedge p_2 \subset \omega \wedge \text{Fin}(p_1) \wedge \text{Fin}(p_2) \wedge p_1 \cap p_2 = 0 \wedge x = \langle p_1, p_2 \rangle].$$

$$2) \quad \mathcal{P} \subseteq \mathcal{P}' \iff \mathcal{P}, \mathcal{P}' \in \text{cond} \wedge (\exists p_1, p_2, p'_1, p'_2) [\mathcal{P} = \langle p_1, p_2 \rangle \wedge \mathcal{P}' = \langle p'_1, p'_2 \rangle \wedge p_1 \subseteq p'_1 \wedge p_2 \subseteq p'_2].$$

$$3) \quad (\forall \mathcal{P} \in \text{cond}) (\exists p_1, p_2 \in \mathcal{M}_0) [\mathcal{P} = \langle p_1, p_2 \rangle].$$

Corollary 18.9.

- 1) $x \in \text{cond}$ is \mathcal{M}_0 -definable.
- 2) $\mathcal{P} \subseteq \mathcal{P}'$ is \mathcal{M}_0 -definable.

Proof. Theorems 18.8 and 17.12.

Definition 18.10 (forcing for unlimited statements).

- 1) $\mathcal{P} \Vdash '(\exists x_i) \varphi(x_i)' \leftrightarrow (\exists \beta \in \alpha_0) [\mathcal{P} \Vdash '\varphi(k_\beta)']$.
- 2) $\mathcal{P} \Vdash '\neg \varphi' \leftrightarrow (\forall \mathcal{P}' \supseteq \mathcal{P}) \neg [\mathcal{P}' \Vdash '\varphi']$.
- 3) $\mathcal{P} \Vdash '\varphi \wedge \psi' \leftrightarrow [\mathcal{P} \Vdash '\varphi'] \wedge [\mathcal{P} \Vdash '\psi']$.

Remark. By $\varphi(k_\beta)$ we of course mean the formula obtained from $\varphi(x_i)$ by replacing x_i at each occurrence by k_β . By definition $'(\exists x_i) \varphi(x_i)' \in \text{culf}(\alpha_0)$ means that x_i does not occur bound in $\varphi(x_i)$. Hence $'\varphi(k_\beta)'$ is also a statement either limited or unlimited. Furthermore $'\varphi(k_\beta)'$ is of smaller degree than $'(\exists x_i) \varphi(x_i)'$. Also $'\varphi'$ is of lower degree than $'\neg \varphi'$ and $'\varphi'$ and $'\psi'$ are each of lower degree than $'\varphi \wedge \psi'$. Thus 1)–3) reduce the question of what $\mathcal{P} \Vdash '\varphi'$ means to the same question for formulas of degree lower than $\text{deg}(\alpha_0; '\varphi')$. After a finite number of steps the question is reduced to what $\mathcal{P} \Vdash '\varphi'$ means for limited statements.

Definition 18.11 (forcing for limited statements).

- 1) $\mathcal{P} \Vdash '\neg \varphi' \leftrightarrow (\forall \mathcal{P}' \supseteq \mathcal{P}) \neg [\mathcal{P}' \Vdash '\varphi']$.
- 2) $\mathcal{P} \Vdash '\varphi \wedge \psi' \leftrightarrow [\mathcal{P} \Vdash '\varphi'] \wedge [\mathcal{P} \Vdash '\psi']$.
- 3) $\mathcal{P} \Vdash '(\exists^\beta x_i) \varphi(x_i)' \leftrightarrow (\exists \alpha < \beta) \mathcal{P} \Vdash '\varphi(k_\alpha)'$.
- 4) $\beta < \alpha \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) [k_\alpha = x_i \wedge x_i \in k_\beta]']$.
- 5) If $\alpha < \beta \wedge \omega + 1 < \beta \wedge K'_1 \beta = \gamma \wedge K'_2 \beta = \delta$ then

$$K'_3 \beta = 0 \rightarrow \mathcal{P} \Vdash 'k_\alpha \in k_\beta'$$

$$K'_3 \beta = 1 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash 'k_\alpha = k_\gamma \vee k_\alpha = k_\delta']$$

$$K'_3 \beta = 2 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) (\exists^\beta x_j) [k_\alpha = \langle x_i, x_j \rangle \wedge x_i \in x_j \wedge k_\alpha \in k_\gamma]']$$

$$K'_3 \beta = 3 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash 'k_\alpha \in k_\gamma \wedge k_\alpha \notin k_\delta']$$

$$K'_3 \beta = 4 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) (\exists^\beta x_j) [k_\alpha = \langle x_i, x_j \rangle \wedge k_\alpha \in k_\gamma \wedge x_i \in k_\delta]']$$

$$K'_3 \beta = 5 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) [k_\alpha \in k_\gamma \wedge \langle k_\alpha, x_i \rangle \in k_\delta]']$$

$$K'_3 \beta = 6 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) (\exists^\beta x_j) [k_\alpha \in k_\gamma \wedge k_\alpha = \langle x_i, x_j \rangle \wedge \langle x_j, x_i \rangle \in k_\delta]']$$

$$K_3 \beta = 7 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) (\exists^\beta x_j) (\exists^\beta x_p) [k_\alpha \in k_\gamma \wedge k_\alpha = \langle x_i, x_j, x_p \rangle \wedge \langle x_p, x_i, x_j \rangle \in k_\delta]'].$$

$$K_3 \beta = 8 \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) (\exists^\beta x_j) (\exists^\beta x_p) [k_\alpha \in k_\gamma \wedge k_\alpha = \langle x_i, x_j, x_p \rangle \wedge \langle x_i, x_p, x_j \rangle \in k_\delta]'].$$

- 6) $[\mathcal{P} = \langle p_1, p_2 \rangle] \wedge [\alpha \in p_1] \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_{\omega+1}']$.
 $[\mathcal{P} = \langle p_1, p_2 \rangle] \wedge [\alpha \in p_2] \rightarrow \neg [\mathcal{P} \Vdash 'k_\alpha \in k_{\omega+1}']$.
- 7) $\neg [\mathcal{P} \Vdash 'k_\omega \in k_{\omega+1}']$.
- 8) $\neg [\mathcal{P} \Vdash 'k_\alpha \in k_\alpha']$.
- 9) $\alpha \leq \omega \wedge \beta \leq \omega \rightarrow [\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \leftrightarrow \alpha \in \beta]$.

Remark. We leave to the reader the verification that, for ' φ ' \in $\text{clf}(\alpha_0)$, each application of 1)–9) of Definition 18.11 either answers the question of the meaning of $\mathcal{P} \Vdash '\varphi'$ or reduces the problem to the question of the meaning of $\mathcal{P} \Vdash '\psi'$ for some ' ψ ' $<$ ' φ '. Before this can be done however we must clarify the role of equality in 4) and 5).

Normally we interpret $k_\alpha = x_i$ as an abbreviation for the unlimited formula

$$(\forall x_j) [x_j \in k_\alpha \leftrightarrow x_j \in x_i]$$

or in primitive symbols

$$\neg (\exists x_j) [x_j \in k_\alpha \wedge x_j \notin x_i] \wedge \neg (\exists x_j) [x_j \in x_i \wedge x_j \notin k_\alpha].$$

However in our principle interpretation $k_\alpha = F_a^* \alpha$. Since $F_a^* \alpha \subseteq F_a^* \alpha$ and since in 4) $k_\alpha = x_i$ is in the scope of a limited quantifier $(\exists^\beta x_j)$ with $\beta < \alpha$ we can and do interpret $k_\alpha = x_i$ in 4) as an abbreviation for the limited formula

$$\neg (\exists^\alpha x_j) [x_j \in k_\alpha \wedge x_j \notin x_i] \wedge \neg (\exists^\alpha x_j) [x_j \in x_i \wedge x_j \notin k_\alpha].$$

Similarly in 5) we interpret $k_\alpha = k_\gamma$ as an abbreviation for the limited formula

$$\neg (\exists^\delta x_j) [x_j \in k_\alpha \wedge x_j \notin k_\gamma] \wedge \neg (\exists^\delta x_j) [x_j \in k_\gamma \wedge x_j \notin k_\alpha].$$

where $\delta = \max(\alpha, \beta)$.

Theorem 18.12. *If ' φ ' \in $s(\alpha_0)$ then $\mathcal{P} \Vdash '\varphi'$ is \mathcal{M}_0 -definable.*

Proof (by $<$ -induction). If ' φ ' \in $s(\alpha_0)$ then φ is $(\exists x_i) \psi(x_i)$ or $\neg \psi$ or $\psi \wedge \eta$ or $(\exists^\beta x_i) \psi(x_i)$ or $k_\alpha \in k_\beta$. From Definition 18.11

$$\mathcal{P} \Vdash '(\exists x_i) \psi(x_i)' \leftrightarrow (\exists \alpha \in \alpha_0) [\mathcal{P} \Vdash '\psi(k_\alpha)']$$

$$\mathcal{P} \Vdash '\neg \psi' \leftrightarrow (\forall \mathcal{P}' \supseteq \mathcal{P}) \neg [\mathcal{P}' \Vdash '\psi']$$

$$\mathcal{P} \Vdash '\psi \wedge \eta' \leftrightarrow [\mathcal{P} \Vdash '\psi'] \wedge [\mathcal{P} \Vdash '\eta']$$

$$\mathcal{P} \Vdash '(\exists^\beta x_i) \psi(x_i)' \leftrightarrow (\exists \alpha < \beta) [\mathcal{P} \Vdash '\psi(k_\alpha)'].$$

In each case ' $\psi(k_\alpha)$ ' < ' φ ', ' ψ ' < ' φ ', ' η ' < ' φ '. From Corollary 18.9 and the induction hypothesis it follows that $\mathcal{P} \Vdash \varphi$ is \mathcal{M}_0 -definable.

If $\alpha > \beta$ then

$$\begin{aligned} \mathcal{P} \Vdash 'k_\alpha \in k_\beta' &\longleftrightarrow \mathcal{P} \Vdash '(\exists^\beta x_i) [k_\alpha = x_i \wedge x_i \in k_\beta]' \\ &\longleftrightarrow (\exists \gamma < \beta) [\mathcal{P} \Vdash 'k_\alpha = k_\gamma' \wedge \mathcal{P} \Vdash 'k_\gamma \in k_\beta']. \end{aligned}$$

Since ' $k_\alpha = k_\gamma$ ' < ' $k_\alpha \in k_\beta$ ' and $\gamma < \beta$ it follows from the induction hypothesis that $\mathcal{P} \Vdash 'k_\alpha \in k_\beta'$ is \mathcal{M}_0 -definable for $\alpha > \beta$ if, in general, $\mathcal{P} \Vdash 'k_\alpha \in k_\beta'$ is \mathcal{M}_0 -definable for $\alpha \leq \beta$.

If $\alpha = \beta$

$$\neg \mathcal{P} \Vdash 'k_\alpha \in k_\beta'$$

and hence

$$\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \longleftrightarrow \alpha \in \beta.$$

Thus $\mathcal{P} \Vdash 'k_\alpha \in k_\beta'$ is \mathcal{M}_0 -definable.

If $[\alpha < \beta] \wedge [\beta > \omega + 1]$ then

$$K_3 \beta = 0 \rightarrow \mathcal{P} \Vdash 'k_\alpha \in k_\beta'.$$

Hence

$$\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \longleftrightarrow \alpha = \alpha.$$

Thus $\mathcal{P} \Vdash 'k_\alpha \in k_\beta'$ is \mathcal{M}_0 -definable. In each other case i.e. $K_3 \beta = 1, \dots, K_3 \beta = 8$ it follows from Definition 18.11 and the induction hypothesis that $\mathcal{P} \Vdash 'k_\alpha \in k_\beta'$ is \mathcal{M}_0 -definable.

If $\alpha < \beta = \omega + 1$ then

$$\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \longleftrightarrow [\alpha < \omega] \wedge [\mathcal{P} = \langle p_1, p_2 \rangle] \wedge [\alpha \in p_1].$$

Thus $\mathcal{P} \Vdash 'k_\alpha \in k_\beta'$ is \mathcal{M}_0 -definable.

If $\alpha < \beta < \omega + 1$ then

$$\mathcal{P} \Vdash 'k_\alpha \in k_\beta'.$$

Hence

$$\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \longleftrightarrow \alpha \in \beta.$$

Thus $\mathcal{P} \Vdash 'k_\alpha \in k_\beta'$ is \mathcal{M}_0 -definable.

Remark. We next wish to establish that only true statements are forced and that every true statement is forced. This we prove by means of a forcing function for whose definition we need the class of statements that are shorter than a given statement. For this we introduce clf defined as $\text{clf}(\alpha_0)$ is defined but with α_0 replaced by On .

Definition 18.13. $\text{shs}(\varphi) \triangleq \{\beta \mid \beta \leq \varphi\}$.

Definition 18.14.

$$\begin{aligned}
& \text{forc}(g, \ulcorner \varphi \urcorner) \longleftrightarrow \ulcorner \varphi \urcorner \in \text{clf} \wedge g \text{ Fn cond} \times \text{shs}(\ulcorner \varphi \urcorner) \wedge \mathcal{W}(g) = 2 \\
& \wedge (\forall \mathcal{P}) [\ulcorner \neg \psi \urcorner \in \text{shs}(\ulcorner \varphi \urcorner) \\
& \quad \rightarrow [g' \langle \mathcal{P}, \ulcorner \neg \psi \urcorner \rangle = 1 \longleftrightarrow (\forall \mathcal{P}' \supseteq \mathcal{P}) g' \langle \mathcal{P}', \ulcorner \psi \urcorner \rangle = 0]] \\
& \wedge (\forall \mathcal{P}) [\ulcorner \psi \wedge \eta \urcorner \in \text{shs}(\ulcorner \varphi \urcorner) \\
& \quad \rightarrow [g' \langle \mathcal{P}, \ulcorner \psi \wedge \eta \urcorner \rangle = 1 \longleftrightarrow [g' \langle \mathcal{P}, \ulcorner \psi \urcorner \rangle = 1 \wedge g' \langle \mathcal{P}, \ulcorner \eta \urcorner \rangle = 1]]] \\
& \wedge (\forall \mathcal{P}) [\ulcorner (\exists^\beta x_i) \psi(x_i) \urcorner \in \text{shs}(\ulcorner \varphi \urcorner) \\
& \quad \rightarrow [g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) \psi(x_i) \urcorner \rangle = 1 \longleftrightarrow (\exists \alpha < \beta) g' \langle \mathcal{P}, \ulcorner \psi(k_\alpha) \urcorner \rangle = 1]] \\
& \wedge (\forall \mathcal{P}) (\forall \alpha) (\forall \beta) [\ulcorner k_\alpha \in k_\beta \urcorner \in \text{shs}(\ulcorner \varphi \urcorner) \\
& \quad \rightarrow [g' \langle \mathcal{P}, \ulcorner k_\alpha \in k_\beta \urcorner \rangle = 1 \longleftrightarrow [\alpha > \beta \wedge g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) [k_\alpha = x_i \wedge x_i \in k_\beta] \urcorner \rangle = 1] \\
& \quad \vee [\alpha < \beta \wedge \beta > \omega + 1 \wedge K'_1 \beta = \gamma \wedge K'_2 \beta = \delta \wedge [K'_3 \beta = 0 \\
& \quad \vee [K'_3 \beta = 1 \wedge g' \langle \mathcal{P}, \ulcorner k_\alpha = k_\gamma \vee k_\alpha = k_\delta \urcorner \rangle = 1] \\
& \quad \vee [K'_3 \beta = 2 \wedge g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) (\exists^\beta x_j) [k_\alpha = \langle x_i, x_j \rangle \wedge x_i \in x_j \wedge k_\alpha \in k_\gamma] \urcorner \rangle = 1] \\
& \quad \vee [K'_3 \beta = 3 \wedge g' \langle \mathcal{P}, \ulcorner k_\alpha \in k_\gamma \wedge k_\alpha \in k_\delta \urcorner \rangle = 1] \\
& \quad \vee [K'_3 \beta = 4 \wedge g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) (\exists^\beta x_j) [k_\alpha = \langle x_i, x_j \rangle \wedge k_\alpha \in k_\gamma \wedge x_i \in k_\delta] \urcorner \rangle = 1] \\
& \quad \vee [K'_3 \beta = 5 \wedge g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) [k_\alpha \in k_\gamma \wedge \langle k_\alpha, x_i \rangle \in k_\delta] \urcorner \rangle = 1] \\
& \quad \vee [K'_3 \beta = 6 \wedge g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) (\exists^\beta x_j) [k_\alpha \in k_\gamma \wedge k_\alpha = \langle x_i, x_j \rangle \wedge \langle x_j, x_i \rangle \in k_\delta] \urcorner \rangle = 1] \\
& \quad \vee [K'_3 \beta = 7 \wedge g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) (\exists^\beta x_j) (\exists^\beta x_p) [k_\alpha = \langle x_i, x_j, x_p \rangle \\
& \quad \quad \quad \wedge k_\alpha \in k_\gamma \wedge \langle x_p, x_i, x_j \rangle \in k_\delta] \urcorner \rangle = 1] \\
& \quad \vee [K'_3 \beta = 8 \wedge g' \langle \mathcal{P}, \ulcorner (\exists^\beta x_i) (\exists^\beta x_j) (\exists^\beta x_p) [k_\alpha = \langle x_i, x_j, x_p \rangle \\
& \quad \quad \quad \wedge k_\alpha \in k_\gamma \wedge \langle x_i, x_p, x_j \rangle \in k_\delta] \urcorner \rangle = 1]]] \\
& \vee (\exists p_1, p_2) [\alpha < \beta = \omega + 1 \wedge \mathcal{P} = \langle p_1, p_2 \rangle \wedge \alpha \in p_1] \vee [\alpha < \beta \leq \omega]].
\end{aligned}$$

Remark. From Definition 18.14 we then prove the following lemmas.

Lemma 1. $\text{forc}(g, \ulcorner \varphi \urcorner)$ is absolute w.r.t. every standard transitive model of ZF.

Lemma 2. $\ulcorner \varphi \urcorner \in \text{clf} \rightarrow (\exists ! g) \text{forc}(g, \ulcorner \varphi \urcorner)$.

Definition 3. $\text{Forc}(\mathcal{P}, \ulcorner \varphi \urcorner) \longleftrightarrow (\exists g) [\text{forc}(g, \ulcorner \varphi \urcorner) \wedge g' \langle \mathcal{P}, \ulcorner \varphi \urcorner \rangle = 1]$.

Lemma 4. $\text{Forc}(\mathcal{P}, \ulcorner \varphi \urcorner)$ is absolute w.r.t. every standard transitive model of ZF.

Theorem 18.15. $\ulcorner \varphi \urcorner \in \text{clf}(\alpha_0) \rightarrow [\mathcal{P} \Vdash \ulcorner \varphi \urcorner \longleftrightarrow [\text{Forc}(\mathcal{P}, \ulcorner \varphi \urcorner)]^{\mathcal{M}_0}]$.

Remark. From Theorem 18.15 it is clear that $\mathcal{P} \Vdash \ulcorner \varphi \urcorner$ is \mathcal{M}_0 -definable.

Theorem 18.16. $\ulcorner \varphi \urcorner \in \text{clf}(\alpha_0) \wedge \mathcal{P} \Vdash \ulcorner \varphi \urcorner$ is \mathcal{M}_0 -definable.

Theorem 18.17. $\neg[\mathcal{P} \Vdash \varphi \wedge \mathcal{P} \Vdash \neg\varphi]$.

Proof. $\mathcal{P} \Vdash \neg\varphi \leftrightarrow (\forall \mathcal{P}' \supseteq \mathcal{P}) \neg[\mathcal{P}' \Vdash \varphi]$. In particular $\mathcal{P} \Vdash \neg\varphi \rightarrow \neg[\mathcal{P} \Vdash \varphi]$ and hence

$$\mathcal{P} \Vdash \varphi \rightarrow \neg[\mathcal{P} \Vdash \neg\varphi].$$

Theorem 18.18. $[\mathcal{P}' \supseteq \mathcal{P}] \wedge [\mathcal{P} \Vdash \varphi] \rightarrow \mathcal{P}' \Vdash \varphi$.

Proof (by $<$ -induction). φ must be 1) $(\exists x_i) \psi(x_i)$ or 2) $\neg\psi$ or 3) $\psi \wedge \eta$ or 4) $(\exists^\beta x_i) \psi(x_i)$ or 5) $k_\alpha \in k_\beta$. In cases 1)–4) we have as our induction hypothesis

$$\mathcal{P}' \supseteq \mathcal{P} \wedge \mathcal{P} \Vdash \psi \rightarrow \mathcal{P}' \Vdash \psi \quad \text{and} \quad \mathcal{P}' \supseteq \mathcal{P} \wedge \mathcal{P} \Vdash \eta \rightarrow \mathcal{P}' \Vdash \eta.$$

Then since $\mathcal{P}' \supseteq \mathcal{P}$,

$$\begin{aligned} 1) \quad \mathcal{P} \Vdash (\exists x_i) \psi(x_i) &\rightarrow (\exists \beta \in \alpha_0) \mathcal{P} \Vdash \psi(k_\beta) \\ &\rightarrow (\exists \beta \in \alpha_0) \mathcal{P}' \Vdash \psi(k_\beta) \\ &\rightarrow \mathcal{P}' \Vdash (\exists x_i) \psi(x_i). \end{aligned}$$

$$\begin{aligned} 2) \quad \mathcal{P} \Vdash \neg\psi &\rightarrow (\forall \mathcal{P}'' \supseteq \mathcal{P}) \neg[\mathcal{P}'' \Vdash \psi] \\ &\rightarrow (\forall \mathcal{P}'' \supseteq \mathcal{P}') \neg[\mathcal{P}'' \Vdash \psi] \\ &\rightarrow \mathcal{P}' \Vdash \neg\psi. \end{aligned}$$

$$\begin{aligned} 3) \quad \mathcal{P} \Vdash \psi \wedge \eta &\rightarrow [\mathcal{P} \Vdash \psi] \wedge [\mathcal{P} \Vdash \eta] \\ &\rightarrow [\mathcal{P}' \Vdash \psi] \wedge [\mathcal{P}' \Vdash \eta] \\ &\rightarrow \mathcal{P}' \Vdash \psi \wedge \eta. \end{aligned}$$

$$\begin{aligned} 4) \quad \mathcal{P} \Vdash (\exists^\beta x_i) \psi(x_i) &\rightarrow (\exists \alpha < \beta) \mathcal{P} \Vdash \psi(k_\alpha) \\ &\rightarrow (\exists \alpha < \beta) \mathcal{P}' \Vdash \psi(k_\alpha) \\ &\rightarrow \mathcal{P}' \Vdash (\exists^\beta x_i) \psi(x_i). \end{aligned}$$

5) If $\alpha > \beta$ then from Definition 18.11 and the induction hypothesis

$$\begin{aligned} \mathcal{P} \Vdash k_\alpha \in k_\beta &\rightarrow \mathcal{P} \Vdash (\exists^\beta x_i) [k_\alpha = x_i \wedge x_i \in k_\beta] \\ &\rightarrow (\exists \gamma < \beta) \mathcal{P} \Vdash k_\alpha = k_\gamma \wedge k_\gamma \in k_\beta \\ &\rightarrow (\exists \gamma < \beta) [\mathcal{P} \Vdash k_\alpha = k_\gamma \wedge \mathcal{P} \Vdash k_\gamma \in k_\beta]. \end{aligned}$$

If $\mathcal{P} \Vdash k_\gamma \in k_\beta \rightarrow \mathcal{P}' \Vdash k_\gamma \in k_\beta$ whenever $\gamma < \beta$ it then follows from the induction hypothesis that

$$\begin{aligned} \mathcal{P} \Vdash k_\alpha \in k_\beta &\rightarrow (\exists \gamma < \beta) [\mathcal{P}' \Vdash k_\alpha = k_\gamma \wedge \mathcal{P}' \Vdash k_\gamma \in k_\beta] \\ &\rightarrow \mathcal{P}' \Vdash k_\alpha \in k_\beta. \end{aligned}$$

If $\alpha = \beta$ then

$$\neg[\mathcal{P} \Vdash k_\alpha \in k_\beta]$$

and hence

$$\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \rightarrow \mathcal{P}' \Vdash 'k_\alpha \in k_\beta'.$$

If $[\alpha < \beta] \wedge [\beta > \omega + 1]$ then from Definition 18.11 and the induction hypothesis it follows that

$$[\mathcal{P}' \supseteq \mathcal{P}] \wedge [\mathcal{P} \Vdash 'k_\alpha \in k_\beta'] \rightarrow \mathcal{P}' \Vdash 'k_\alpha \in k_\beta'.$$

If $[\alpha < \beta = \omega + 1] \wedge [\mathcal{P} = \langle p_1, p_2 \rangle] \wedge [\mathcal{P}' = \langle p'_1, p'_2 \rangle]$ then $\mathcal{P}' \supseteq \mathcal{P}$ implies $p_1 \subseteq p'_1$ and

$$\begin{aligned} \mathcal{P} \Vdash 'k_\alpha \in k_\beta' &\rightarrow \alpha \in p_1 \\ &\rightarrow \alpha \in p'_1 \\ &\rightarrow \mathcal{P}' \Vdash 'k_\alpha \in k_\beta'. \end{aligned}$$

If $\alpha < \beta \leq \omega$ then $\mathcal{P}' \Vdash 'k_\alpha \in k_\beta'$. Hence

$$\mathcal{P} \Vdash 'k_\alpha \in k_\beta' \rightarrow \mathcal{P}' \Vdash 'k_\alpha \in k_\beta'.$$

Theorem 18.19.

$$(\forall ' \varphi ' \in s(\alpha_0)) (\forall \mathcal{P}) (\exists \mathcal{P}' \supseteq \mathcal{P}) [\mathcal{P}' \Vdash ' \varphi ' \vee \mathcal{P}' \Vdash ' \neg \varphi '].$$

Proof. Either $\mathcal{P} \Vdash ' \neg \varphi '$ or $\neg [\mathcal{P} \Vdash ' \neg \varphi ']$.

But $[\mathcal{P} \Vdash ' \neg \varphi '] \leftrightarrow (\forall \mathcal{P}' \supseteq \mathcal{P}) \neg [\mathcal{P}' \Vdash ' \varphi ']$

$$\neg [\mathcal{P} \Vdash ' \neg \varphi '] \leftrightarrow (\exists \mathcal{P}' \supseteq \mathcal{P}) [\mathcal{P}' \Vdash ' \varphi '].$$

Remark. Theorem 18.19 is a key result for our theory. It assures us of success in our search for a subset a of ω such that every statement of ZF $\{\alpha_0\}$ can be resolved with only a knowledge of some finite subset of a and some finite subset of $\omega - a$.

For each statement φ of ZF $\{\alpha_0\}$ there is a forcing condition \mathcal{P} such that $\mathcal{P} \Vdash ' \varphi '$ or $\mathcal{P} \Vdash ' \neg \varphi '$. Since $s(\alpha_0)$ is countable we can enumerate the statements of ZF $\{\alpha_0\}$ and, using Theorem 18.19, form a complete sequence of forcing conditions:

Definition 18.20. A sequence $\{\mathcal{P}_n\}$ of forcing conditions is a *complete sequence* iff $(\forall n) [\mathcal{P}_n \subseteq \mathcal{P}_{n+1}]$ and

$$(\forall ' \varphi ' \in s(\alpha_0)) (\exists n) [\mathcal{P}_n \Vdash ' \varphi ' \vee \mathcal{P}_n \Vdash ' \neg \varphi '].$$

Theorem 18.21. *There exists a complete sequence of forcing conditions.*

Proof. Since $s(\alpha_0) \subseteq \mathcal{M}_0$ and \mathcal{M}_0 is countable $s(\alpha_0)$ is countable, let

$$' \varphi_0 ', ' \varphi_1 ', \dots$$

be an enumeration of $s(\alpha_0)$. Since $\text{cond} \subseteq \mathcal{M}_0$ we can enumerate forcing conditions. Given any such enumeration of forcing conditions we define

$\{\mathcal{P}_n\}$ inductively in the following manner,

$$\mathcal{P}_0 = \langle 0, 0 \rangle$$

By Theorem 18.19

$$(\exists \mathcal{P} \supseteq \mathcal{P}_0) [\mathcal{P} \Vdash \text{'}\varphi_0\text{' } \vee \mathcal{P} \Vdash \text{'}\neg \varphi_0\text{' }].$$

Let \mathcal{P}_1 be the first such \mathcal{P} in our given enumeration. Continuing inductively we take \mathcal{P}_n as the first forcing condition in our given enumeration for which

$$\mathcal{P}_n \supseteq \mathcal{P}_{n-1} \wedge [\mathcal{P}_n \Vdash \text{'}\varphi_{n-1}\text{' } \vee \mathcal{P}_n \Vdash \text{'}\neg \varphi_{n-1}\text{' }].$$

The sequence $\{\mathcal{P}_n\}$ is then complete since

$$(\forall n) [\mathcal{P}_{n+1} \supseteq \mathcal{P}_n \wedge \mathcal{P}_{n+1} \Vdash \text{'}\varphi_n\text{' } \vee \mathcal{P}_{n+1} \Vdash \text{'}\neg \varphi_n\text{' }].$$

Definition 18.22. A set a is a *forcing set* iff $a \subseteq \omega$ and there exists a complete sequence of forcing conditions $\{\mathcal{P}_n\}$ for which

$$a = \bigcup_{n=0}^{\infty} \mathcal{D}(\mathcal{P}_n).$$

Remark. The existence of a forcing set is established by Theorem 18.21. We now wish to prove two things: 1) for each forcing set a , N_a is a standard transitive model of $ZF + AC + GCH$; 2) There exists a forcing set a that is not in \mathcal{M}_0 . From 1) and 2) it then follows that there exists a forcing set a for which N_a is a standard transitive model of $ZF + AC + GCH + V \neq L$.

Theorem 18.23. *There exists a forcing set a that is not in \mathcal{M}_0 .*

Proof. As in the proof of Theorem 18.21 we enumerate $s(\alpha_0)$

$$\text{'}\varphi_0\text{'}, \text{'}\varphi_1\text{'}, \dots$$

and cond. In addition we enumerate the ordinals in \mathcal{M}_0

$$\beta_0, \beta_1, \dots$$

We then define a complete sequence of forcing conditions $\{\mathcal{P}_n\}$ inductively.

$$\begin{aligned} \mathcal{P}_0 &= \{ \langle 0, \{0\} \rangle \} && \text{if } 0 \in F^* \beta_0 \\ &= \{ \langle \{0\}, 0 \rangle \} && \text{if } 0 \notin F^* \beta_0. \end{aligned}$$

From our enumeration of cond we choose \mathcal{P}_1 as the first extension of \mathcal{P}_0 for which

$$\mathcal{P}_1 \Vdash \text{'}\varphi_0\text{' } \vee \mathcal{P}_1 \Vdash \text{'}\neg \varphi_0\text{' }.$$

If $\mathcal{P}_{2n-1} = \langle p_1, p_2 \rangle$ and if k is the smallest natural number not in $p_1 \cup p_2$ then

$$\begin{aligned} \mathcal{P}_{2m} &= \langle p_1, p_2 \cup \{k\} \rangle && \text{if } k \in F^* \beta_m \\ &= \langle p_1 \cup \{k\}, p_2 \rangle && \text{if } k \notin F^* \beta_m. \end{aligned}$$

\mathcal{P}_{2m+1} is the first extension of \mathcal{P}_{2m} for which

$$\mathcal{P}_{2m+1} \Vdash \ulcorner \varphi_m \urcorner \vee \mathcal{P}_{2m+1} \Vdash \ulcorner \neg \varphi_m \urcorner.$$

Thus $\{\mathcal{P}_n\}$ is a complete sequence of forcing conditions. If

$$a = \bigcup_{n=0}^{\infty} \mathcal{D}(\mathcal{P}_n)$$

then a is a forcing set. However from the definition of the forcing conditions \mathcal{P}_n we see that

$$(\forall n) (\exists k) [[k \in F^* \beta_n \wedge k \notin a] \vee [k \in a \wedge k \notin F^* \beta_n]]$$

i.e.

$$(\forall n) [a \neq F^* \beta_n].$$

Thus

$$a \notin \mathcal{M}_0.$$

Remark. We turn now to the problem of proving that if a is a forcing set then N_a is a model of ZF + AC + GCH.

Theorem 18.24. *If a is a forcing set corresponding to the complete sequence of forcing conditions $\{\mathcal{P}_n\}$ then for each $\ulcorner \varphi \urcorner \in s(\alpha_0)$*

$$(\exists n) [\mathcal{P}_n \Vdash \ulcorner \varphi \urcorner] \longleftrightarrow \varphi^{N_a}.$$

Proof (by $<$ -induction). If $\ulcorner \varphi \urcorner \in s(\alpha_0)$ then φ is 1) $(\exists x_i) \psi(x_i)$ or 2) $\neg \psi$ or 3) $\psi \wedge \eta$ or 4) $(\exists^\beta x_i) \psi(x_i)$ or 5) $k_\alpha \in k_\beta$. In cases 1)–4) we have as our induction hypothesis

$$(\exists i) [\mathcal{P}_i \Vdash \ulcorner \psi \urcorner] \longleftrightarrow \psi^{N_a} \quad \text{and} \quad (\exists i) [\mathcal{P}_i \Vdash \ulcorner \eta \urcorner] \longleftrightarrow \eta^{N_a}.$$

Then 1)

$$\begin{aligned} (\exists i) [\mathcal{P}_i \Vdash \ulcorner (\exists x_i) \psi(x_i) \urcorner] &\longleftrightarrow (\exists i) (\exists \alpha \in \alpha_0) [\mathcal{P}_i \Vdash \ulcorner \psi(k_\alpha) \urcorner] \\ &\longleftrightarrow (\exists \alpha \in \alpha_0) [\psi(k_\alpha)]^{N_a} \\ &\longleftrightarrow (\exists x_i \in N) [\psi(x_i)]^{N_a} \\ &\longleftrightarrow [(\exists x_i) \psi(x_i)]^{N_a}. \end{aligned}$$

2) Since $\{\mathcal{P}_n\}$ is a complete sequence it follows from Theorems 18.17 and 18.18 that

$$\begin{aligned} \mathcal{P}_i \Vdash \ulcorner \neg \psi \urcorner &\rightarrow (\forall n \geq i) [\mathcal{P}_n \Vdash \ulcorner \neg \psi \urcorner] \\ &\rightarrow (\forall n \geq i) \neg [\mathcal{P}_n \Vdash \ulcorner \psi \urcorner] \end{aligned}$$

Also

$$(\forall n < i) [\mathcal{P}_n \Vdash \ulcorner \psi \urcorner \rightarrow \mathcal{P}_i \Vdash \ulcorner \psi \urcorner].$$

Thus

$$\begin{aligned} (\exists i) [\mathcal{P}_i \Vdash \neg \psi] &\longleftrightarrow (\forall n) \neg [\mathcal{P}_n \Vdash \psi] \\ &\longleftrightarrow \neg (\exists i) [\mathcal{P}_i \Vdash \psi] \\ &\longleftrightarrow \neg \psi^{N_a}. \end{aligned}$$

$$\begin{aligned} 3) \quad (\exists i) [\mathcal{P}_i \Vdash \psi \wedge \eta] &\longleftrightarrow (\exists i) [\mathcal{P}_i \Vdash \psi \wedge \mathcal{P}_i \Vdash \eta] \\ &\longleftrightarrow (\exists i) [\mathcal{P}_i \Vdash \psi] \wedge (\exists i) [\mathcal{P}_i \Vdash \eta] \\ &\longleftrightarrow \psi^{N_a} \wedge \eta^{N_a}. \end{aligned}$$

$$\begin{aligned} 4) \quad (\exists i) [\mathcal{P}_i \Vdash (\exists^\beta x_j) \psi(x_j)] &\longleftrightarrow (\exists i) (\exists \alpha < \beta) [\mathcal{P}_i \Vdash \psi(k_\alpha)] \\ &\longleftrightarrow (\exists \alpha < \beta) [\psi(k_\alpha)]^{N_a} \\ &\longleftrightarrow (\exists x_j) (\exists \alpha) [\alpha \in \beta \wedge x_j = k_\alpha \wedge [\psi(x_j)]^{N_a}] \\ &\longleftrightarrow [(\exists^\beta x_j) \psi(x_j)]^{N_a}. \end{aligned}$$

5) If $\alpha > \beta$

$$(\exists i) [\mathcal{P}_i \Vdash k_\alpha \in k_\beta] \longleftrightarrow (\exists i) [\mathcal{P}_i \Vdash (\exists^\beta x_j) [k_\alpha = x_j \wedge x_j \in k_\beta]].$$

Then from the induction hypothesis

$$\begin{aligned} (\exists i) [\mathcal{P}_i \Vdash k_\alpha \in k_\beta] &\longleftrightarrow [(\exists^\beta x_j) [k_\alpha = x_j \wedge x_j \in k_\beta]]^{N_a} \\ &\longleftrightarrow [k_\alpha \in k_\beta]^{N_a}. \end{aligned}$$

If $\alpha = \beta$ then $(\forall i) \neg [\mathcal{P}_i \Vdash k_\alpha \in k_\beta]$ and $\neg [k_\alpha \in k_\beta]$, hence

$$(\exists i) [\mathcal{P}_i \Vdash k_\alpha \in k_\beta] \longleftrightarrow [k_\alpha \in k_\beta]^{N_a}.$$

If $\alpha < \beta \wedge \beta > \omega + 1$ then from Definition 18.10 and the induction hypothesis it is clear that

$$(\exists i) [\mathcal{P}_i \Vdash k_\alpha \in k_\beta] \longleftrightarrow [k_\alpha \in k_\beta]^{N_a}.$$

If $\alpha < \beta = \omega + 1$ then

$$(\exists i) [\mathcal{P}_i \Vdash k_\alpha \in k_\beta] \longleftrightarrow (\exists i) (\exists p_1) (\exists p_2) [\mathcal{P}_i = \langle p_1, p_2 \rangle \wedge k_\alpha \in p_1].$$

From our interpretation i.e.

$$k_\alpha = F_a^\alpha$$

we have that $k_\beta = a$ and hence

$$k_\alpha \in k_\beta \longleftrightarrow (\exists i) (\exists p_1) (\exists p_2) [\mathcal{P}_i = \langle p_1, p_2 \rangle \wedge k_\alpha \in p_1].$$

Thus $(\exists i) [\mathcal{P}_i \Vdash k_\alpha \in k_\beta] \longleftrightarrow [k_\alpha \in k_\beta]^{N_a}$.

If $\alpha < \beta < \omega + 1$ then

$$(\exists i) [\mathcal{P}_i \Vdash k_\alpha \in k_\beta] \longleftrightarrow [\alpha \in \beta]^{N_a}.$$

Remark. Theorem 18.24 says that every true statement in $s(\alpha_0)$ is forced and only true statements in $s(\alpha_0)$ are forced. Thus to prove that N_a is a standard transitive model of ZF we must prove that each axiom of ZF is forced. Indeed in view of Theorem 17.7 it is sufficient to prove that the Axiom of Powers is forced and each instance of the Axiom Schema of Replacement is forced. For this we need the following properties of fundamental functions. By a fundamental function we mean a function on n -tuples that is generated from fundamental operations by composition. The definition is inductive.

Definition 18.25. G is a *fundamental function* on n -tuples of type zero iff $G \mathcal{F}_n V^n$ and $(\exists i \leq n) [(\forall a_1) \dots (\forall a_n) [G(a_1, \dots, a_n) = a_i] \wedge i \geq 1]$. G is a fundamental function on n -tuples of type $m+1$ iff G is a fundamental function on n -tuples of type m or there exist fundamental functions on n -tuples G_1 , and G_2 of type m and a fundamental operation \mathcal{F}_i such that

$$(\forall a_1) \dots (\forall a_n) [G(a_1, \dots, a_n) = \mathcal{F}_i(G_1(a_1, \dots, a_n), G_2(a_1, \dots, a_n))].$$

Theorem 18.26. *If G is a fundamental function on n -tuples and \mathcal{M} is a standard transitive model of ZF then*

$$a_1 \in \mathcal{M} \wedge \dots \wedge a_n \in \mathcal{M} \rightarrow G(a_1, \dots, a_n) \in \mathcal{M}.$$

Proof. Since every standard transitive model \mathcal{M} is closed with respect to the fundamental operations the result follows by induction on the type of G . Details are left to the reader.

Theorem 18.27. *There exists a fundamental function G on $(n+1)$ -tuples, such that for $0 \leq i \leq n$ and $0 \leq j \leq n$*

$$[a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \wedge [a_j \subseteq a_0] \rightarrow G(a_0, a_1, \dots, a_n) = a_i \times a_j.$$

Proof. If $[a_i \subseteq a_0] \wedge [a_j \subseteq a_0] \wedge [a_0 \times a_0 \subseteq a_0]$ then

$$a_i \times a_j \subseteq a_0 \times a_0 \subseteq a_0.$$

Then

$$a_i \times a_j \subseteq [a_0 \cap (a_i \times V)].$$

Furthermore, $a_0 \times a_0 \subseteq a_0 \rightarrow a_0 \times a_0 \subseteq a_0^{-1}$, hence

$$a_i \times a_j \subseteq [a_0 \cap (a_j \times V)]^{-1}.$$

Thus

$$a_i \times a_j \subseteq [a_0 \cap (a_i \times V)] \cap [a_0 \cap (a_j \times V)]^{-1}.$$

Furthermore if

$$\langle x, y \rangle \in [a_0 \cap (a_i \times V)] \cap [a_0 \cap (a_j \times V)]^{-1}$$

then

$$\langle x, y \rangle \in a_i \times V \quad \text{and} \quad \langle x, y \rangle \in V \times a_j.$$

Consequently

$$\langle x, y \rangle \in a_i \times a_j.$$

Therefore

$$a_i \times a_j = [a_0 \cap (a_i \times V)] \cap [a_0 \cap (a_j \times V)]^{-1} = \mathcal{F}_6(\mathcal{F}_4(a_0, a_i), \mathcal{F}_4(a_0, a_j)).$$

Theorem 18.28. 1) For each $k > 1$ there exists a fundamental function G on $(n + 1)$ -tuples such that for $0 \leq i \leq n$

$$a_i \times a_i \subseteq a_i \rightarrow G(a_0, \dots, a_n) = a_i^k.$$

2) There exists a fundamental function G on $(n + 1)$ -tuples such that

$$G(a_0, \dots, a_n) = 0.$$

Proof 1) (By induction on k .) If $k = 2$ then from Theorem 18.27 there is a fundamental function G_1 on $(n + 2)$ -tuples such that

$$b \times b \subseteq b \rightarrow G_1(b, a_0, a_1, \dots, a_n) = b \times b.$$

If $G(a_0, \dots, a_n) = G_1(a_i, a_0, a_1, \dots, a_n)$ then G is an $(n + 1)$ -ary fundamental function (Why?) and

$$a_i \times a_i \subseteq a_i \rightarrow G(a_0, \dots, a_n) = G_1(a_i, a_0, a_1, \dots, a_n) = a_i \times a_i.$$

As our induction hypothesis we assume that there exists an $(n + 1)$ -ary fundamental function G_2 such that

$$a_i \times a_i \subseteq a_i \rightarrow G_2(a_0, \dots, a_n) = a_i^k.$$

If $G(a_0, \dots, a_n) = G_1(G_2(a_0, \dots, a_n), a_0, \dots, a_n)$ then since $a_i^k \subseteq a_i$ (proof by induction)

$$a_i \times a_i \subseteq a_i \rightarrow G(a_0, \dots, a_n) = G_1(G_2(a_0, \dots, a_n), a_0, \dots, a_n) = a_i^k \times a_i = a_i^{k+1}.$$

2) The proof is left to the reader.

Theorem 18.29. There exists a fundamental function G on $(n + 1)$ -tuples such that for $0 \leq i \leq n$

$$\text{Tr}(a_0) \wedge [a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \rightarrow G(a_0, a_1, \dots, a_n) = \mathcal{D}(a_i).$$

Proof. $\text{Tr}(a_0) \wedge [a_i \subseteq a_0] \rightarrow [\mathcal{D}(a_i) \subseteq a_0]$. Then

$$\mathcal{D}(a_i) = a_0 \cap \mathcal{D}(a_i) = \mathcal{F}_5(a_0, a_i).$$

Theorem 18.30. There exists a fundamental function G on $(n + 1)$ -tuples such that for $0 \leq i \leq n$

$$\text{Tr}(a_0) \wedge [a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \rightarrow G(a_0, a_1, \dots, a_n) = \mathcal{W}(a_i).$$

Proof. $[a_i \subseteq a_0] \wedge [a_0 \times a_0 \subseteq a_0] \wedge \text{Tr}(a_0) \rightarrow [a_i^{-1} \subseteq a_0]$. Then $\mathcal{W}(a_i) = \mathcal{D}(a_i^{-1}) = a_0 \cap \mathcal{D}(a_0 \cap a_i^{-1}) = \mathcal{F}_5(a_0, \mathcal{F}_6(a_0, a_i))$.

Theorem 18.31. For each permutation (i_1, i_2, i_3) of $\{1, 2, 3\}$ there exists a fundamental function G on $(n+1)$ -tuples such that for $0 \leq i \leq n$

$$\begin{aligned} & \text{Tr}(a_0) \wedge [a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \\ & \rightarrow G(a_0, a_1, \dots, a_n) = \{\langle x_1, x_2, x_3 \rangle \mid \langle x_{i_1}, x_{i_2}, x_{i_3} \rangle \in a_i\}. \end{aligned}$$

Proof. $\text{Tr}(a_0) \wedge [a_0 \times a_0 \subseteq a_0]$ implies that

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_{i_1}, x_{i_2}, x_{i_3} \rangle \in a_i\} \subseteq a_0.$$

Therefore

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_1, x_3, x_2 \rangle \in a_i\} = \text{Cnv}_3(a_i) = a_0 \cap \text{Cnv}_3(a_i) = \mathcal{F}_8(a_0, a_i).$$

$$\{\langle x_1, x_2, x_3 \rangle \mid \langle x_3, x_1, x_2 \rangle \in a_i\} = \text{Cnv}_2(a_i) = a_0 \cap \text{Cnv}_2(a_i) = \mathcal{F}_7(a_0, a_i).$$

$$\begin{aligned} \{\langle x_1, x_2, x_3 \rangle \mid \langle x_3, x_2, x_1 \rangle \in a_i\} &= \text{Cnv}_2(\text{Cnv}_3(a_i)) \\ &= a_0 \cap \text{Cnv}_2(a_0 \cap \text{Cnv}_3(a_i)) = \mathcal{F}_7(a_0, \mathcal{F}_8(a_0, a_i)). \end{aligned}$$

$$\begin{aligned} \{\langle x_1, x_2, x_3 \rangle \mid \langle x_1, x_2, x_3 \rangle \in a_i\} &= \text{Cnv}_3(\text{Cnv}_3(a_i)) \\ &= a_0 \cap \text{Cnv}_3(a_0 \cap \text{Cnv}_3(a_i)) = \mathcal{F}_8(a_0, \mathcal{F}_8(a_0, a_i)). \end{aligned}$$

$$\begin{aligned} \{\langle x_1, x_2, x_3 \rangle \mid \langle x_2, x_1, x_3 \rangle \in a_i\} &= \text{Cnv}_3(\text{Cnv}_2(a_i)) \\ &= a_0 \cap \text{Cnv}_3(a_0 \cap \text{Cnv}_2(a_i)) = \mathcal{F}_8(a_0, \mathcal{F}_7(a_0, a_i)). \end{aligned}$$

$$\begin{aligned} \{\langle x_1, x_2, x_3 \rangle \mid \langle x_2, x_3, x_1 \rangle \in a_i\} &= \text{Cnv}_2(\text{Cnv}_2(a_i)) \\ &= a_0 \cap \text{Cnv}_2(a_0 \cap \text{Cnv}_2(a_i)) = \mathcal{F}_7(a_0, \mathcal{F}_7(a_0, a_i)). \end{aligned}$$

Theorem 18.32. There exists a fundamental function G on $(n+1)$ -tuples such that for $0 \leq i \leq n$

$$\text{Tr}(a_i) \wedge [a_i \times a_i \subseteq a_i] \rightarrow G(a_0, \dots, a_n) = \{\langle x, y \rangle \mid [x \in a_i] \wedge [y \in a_i] \wedge [x \in y]\}.$$

Proof. $\text{Tr}(a_i) \wedge [a_i \times a_i \subseteq a_i]$ implies

$$[x \in a_i] \wedge [y \in a_i] \wedge [x \in y] \leftrightarrow [\langle x, y \rangle \in a_i \cap E].$$

Then $\{\langle x, y \rangle \mid [x \in a_i] \wedge [y \in a_i] \wedge [x \in y]\} = a_i \cap E = \mathcal{F}_2(a_i, a_i)$.

Theorem 18.33. There exists a fundamental function G on $(n+1)$ -tuples, such that for $0 \leq i \leq n$ and $0 \leq j \leq n$

$$\begin{aligned} & \text{Tr}(a_0) \wedge [a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \wedge [a_j \subseteq a_0] \\ & \rightarrow (G(a_0, a_1, \dots, a_n) = \{\langle x, z, y \rangle \mid [\langle x, y \rangle \in a_i] \wedge [z \in a_j]\}). \end{aligned}$$

Proof. $\text{Tr}(a_0) \wedge [a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \wedge [a_j \subseteq a_0]$ implies that

$$\{\langle x, z, y \rangle \mid [\langle x, y \rangle \in a_i] \wedge [z \in a_j]\} = a_0 \cap \text{Cnv}_3(a_i \times a_j) = \mathcal{F}_8(a_0, a_i \times a_j).$$

By Theorem 18.27 there exists a fundamental function G_1 such that

$$[a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \wedge [a_j \subseteq a_0] \rightarrow G_1(a_0, a_1, \dots, a_n) = a_i \times a_j.$$

Therefore if

$$G(a_0, a_1, \dots, a_n) = \mathcal{F}_8(a_0, G_1(a_0, a_1, \dots, a_n))$$

then

$$\begin{aligned} & \text{Tr}(a_0) \wedge [a_0 \times a_0 \subseteq a_0] \wedge [a_i \subseteq a_0] \wedge [a_j \subseteq a_0] \\ & \rightarrow G(a_0, a_1, \dots, a_n) = \{\langle x, z, y \rangle \mid [\langle x, y \rangle \in a_i] \wedge [z \in a_j]\}. \end{aligned}$$

Theorem 18.34. *For each wff $\varphi(x_1, \dots, x_m, b_1, \dots, b_n)$ having no free variables other than those listed and for which each quantifier occurrence is of the form $(\forall x \in b_i)$ or $(\exists x \in b_i)$ for $1 \leq i \leq n$ there exists a fundamental function G on $(n+1)$ -tuples such that*

$$\begin{aligned} & \text{Tr}(b) \wedge [b \times b \subseteq b] \wedge [b_1 \subseteq b] \wedge \dots \wedge [b_n \subseteq b] \rightarrow G(b, b_1, \dots, b_n) \\ & = \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \varphi(x_1, \dots, x_m, b_1, \dots, b_n)\}. \end{aligned}$$

Proof. (By induction on k the number of logical symbols in φ .) If $k=0$ then φ is 1) $b_i \in b_j$, 2) $x_i \in b_j$, 3) $b_i \in x_j$, or 4) $x_i \in x_j$.

1) If φ is $b_i \in b_j$ then

$$\begin{aligned} \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge [b_i \in b_j]\} &= b^m \quad \text{if } b_i \in b_j \\ &= 0 \quad \text{if } b_i \notin b_j. \end{aligned}$$

The existence of G is then assured by Theorem 18.28.

2) If φ is $x_i \in b_j$ then

$$\{\langle x_1, \dots, x_i \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_i \in b] \wedge [x_i \in b_j]\} = b^{i-1} \times b_j.$$

Since $b \times b \subseteq b \rightarrow b^{i-1} \subseteq b$ it follows from Theorems 18.27 and 18.28 that there exists a fundamental function G_1 on $(n+1)$ -tuples such that

$$G_1(b, b_1, \dots, b_n) = b^{i-1} \times b_j.$$

But $b^{i-1} \times b_j \subseteq b$ and hence by Theorem 18.27 there is a fundamental function G_{j+1} on $(n+1)$ -tuples for which

$$G_{j+1}(b, b_1, \dots, b_n) = (b^{i-1} \times b_j) \times b.$$

With $m - (j+1)$ further applications of Theorem 18.27 it follows that there is a fundamental function G on $(n+1)$ -tuples such that

$$G(b, b_1, \dots, b_n) = \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge [x_i \in b_j]\}.$$

3) If φ is $b_i \in x_j$ then $[b \times b \subseteq b] \wedge \text{Tr}(b)$ implies

$$[x_j \in b] \wedge [b_i \in x_j] \leftrightarrow [\langle b_i, x_j \rangle \in b \cap E].$$

Therefore

$$\{x_j \mid [x_j \in b] \wedge [b_i \in x_j]\} = \mathcal{W}(b \cap E) = \mathcal{W}(\mathcal{F}_2(b, b)).$$

Since $\mathcal{F}_2(b, b) \subseteq b$ it then follows by Theorem 18.30, that there is a fundamental function G_2 on $(n+1)$ -tuples for which

$$G_2(b, b_1, \dots, b_n) = \mathcal{W}(\mathcal{F}_2(b, b)) = \{x_j \mid [x_j \in b] \wedge [b_i \in x_j]\}.$$

But

$$\begin{aligned} & \{\langle x_1, \dots, x_j \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_j \in b] \wedge [b_i \in x_j]\} \\ &= b^{j-1} \times \{x_j \mid [x_j \in b] \wedge [b_i \in x_j]\}. \end{aligned}$$

Again since $b^{j-1} \subseteq b \wedge \{x_j \mid [x_j \in b] \wedge [b_i \in x_j]\} \subseteq b$ it follows from Theorems 18.27 and 18.28 that there is a fundamental function G_3 on $(n+1)$ -tuples such that

$$G_3(b, b_1, \dots, b_n) = b^{j-1} \times \{x_j \mid [x_j \in b] \wedge [b_i \in x_j]\}.$$

Furthermore $G_3(b, b_1, \dots, b_n) \subseteq b \times b$. Therefore by Theorem 18.27 there is a fundamental function G_{j+1} on $(n+1)$ -tuples for which

$$G_{j+1}(b, b_1, \dots, b_n) = (b^{j-1} \times \{x_j \mid [x_j \in b] \wedge [b_i \in x_j]\}) \times b.$$

As before, with $m - (j+1)$ additional applications of Theorem 18.27, there exists a fundamental function G on $(n+1)$ -tuples for which

$$G(b, b_1, \dots, b_n) = \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge [b_i \in x_j]\}.$$

4) If φ is $x_i \in x_j$ with $i < j$ then

$$\begin{aligned} & \{\langle x_1, \dots, x_i, x_j \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_i \in b] \wedge [x_j \in b] \wedge [x_i \in x_j]\} \\ &= \{\langle x_1, \dots, x_i, x_j \rangle \mid \langle x_1, \dots, x_{i-1}, x_j \rangle \in b^i \\ & \quad \wedge x_j \in \mathcal{D}\{\langle x_i, x_j \rangle \mid [x_i \in b] \wedge [x_j \in b] \wedge [x_i \in x_j]\}\}. \end{aligned}$$

From Theorems 18.28, 18.29, 18.32, and 18.33 it follows that there is a fundamental function G on $(n+1)$ -tuples such that

$$\begin{aligned} & G_1(b, b_1, \dots, b_n) \\ &= \{\langle x_1, \dots, x_i, x_j \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_i \in b] \wedge [x_j \in b] \wedge [x_i \in x_j]\}. \end{aligned}$$

With $j - i - 1$ applications of Theorem 18.33 it then follows that there is a fundamental function G_2 on $(n+1)$ -tuples such that

$$G_2(b, b_1, \dots, b_n) = \{\langle x_1, \dots, x_j \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_j \in b] \wedge [x_i \in x_j]\}.$$

From this $m - j$ applications of Theorem 18.27 establishes the existence of a fundamental function G on $(n+1)$ -tuples for which

$$G(b, b_1, \dots, b_n) = \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge [x_i \in x_j]\}.$$

If φ is $x_i \in x_j$ with $j < i$ then

$$\begin{aligned} & \{\langle x_i, x_j \rangle \mid [x_i \in b] \wedge [x_j \in b] \wedge [x_j \in x_i]\} \\ &= b \cap \{\langle x_j, x_i \rangle \mid [x_i \in b] \wedge [x_j \in b] \wedge [x_j \in x_i]\}^{-1} \\ &= \mathcal{F}_6(b, \{\langle x_j, x_i \rangle \mid [x_i \in b] \wedge [x_j \in b] \wedge [x_j \in x_i]\}). \end{aligned}$$

We then proceed as in the case $i < j$.

If $k > 0$ then φ is

$$\begin{aligned} & \neg \psi(x_1, \dots, x_m, b_1, \dots, b_n) \quad \text{or} \quad (\exists x_0 \in b_i) \psi(x_1, \dots, x_m, x_0, b_1, \dots, b_n) \\ & \text{or} \\ & \psi(x_1, \dots, x_m, b_1, \dots, b_n) \wedge \eta(x_1, \dots, x_m, b_1, \dots, b_n) \end{aligned}$$

where ψ and η have no free variables other than those indicated.

If φ is $\neg \psi(x_1, \dots, x_m, b_1, \dots, b_n)$ then

$$\begin{aligned} & \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \varphi(x_1, \dots, x_m, b_1, \dots, b_n)\} \\ &= b^m - \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \psi(x_1, \dots, x_m, b_1, \dots, b_n)\}. \end{aligned}$$

If φ is $(\exists x_0 \in b_i) \psi(x_1, \dots, x_m, x_0, b_1, \dots, b_n)$ then

$$\begin{aligned} & \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \varphi(x_1, \dots, x_m, b_1, \dots, b_n)\} \\ &= \mathcal{D}\{\langle x_1, \dots, x_m, x_0 \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge [x_0 \in b] \\ & \quad \wedge \psi(x_1, \dots, x_m, x_0, b_1, \dots, b_n)\}. \end{aligned}$$

If φ is $\psi(x_1, \dots, x_m, b_1, \dots, b_n) \wedge \eta(x_1, \dots, x_m, b_1, \dots, b_n)$ then

$$\begin{aligned} & \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \varphi(x_1, \dots, x_m, b_1, \dots, b_n)\} \\ &= b^m - [b^m - \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \psi(x_1, \dots, x_m, b_1, \dots, b_n)\} \\ & \quad - \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \eta(x_1, \dots, x_m, b_1, \dots, b_n)\}]. \end{aligned}$$

In each case it follows from the induction hypothesis and previous theorems that there is a fundamental function G on $(n+1)$ -tuples for which

$$\begin{aligned} & G(b, b_1, \dots, b_n) \\ &= \{\langle x_1, \dots, x_m \rangle \mid [x_1 \in b] \wedge \dots \wedge [x_m \in b] \wedge \varphi(x_1, \dots, x_m, b_1, \dots, b_n)\}. \end{aligned}$$

Theorem 18.35.

$$(\forall a \subseteq \omega)[[\omega < \alpha] \wedge [\alpha \in K_{II}] \wedge (\forall \beta < \alpha)[J^* \langle 0, \beta, 0 \rangle < \alpha] \rightarrow [F_a^* \alpha \times F_a^* \alpha \subseteq F_a^* \alpha]].$$

Proof. Since $\alpha \in K_{II}$, $F_a^* \alpha = F_a^* \alpha$. Furthermore if $[\beta_1 < \alpha] \wedge [\beta_2 < \alpha]$ and $\beta = \max(\beta_1, \beta_2)$ then $\beta + 1 < \alpha$ and hence by hypothesis

$$J^* \langle \beta_1, \beta_2, 1 \rangle < J^* \langle 0, \beta + 1, 0 \rangle < \alpha.$$

Therefore $F_a^* J^* \langle \beta_1, \beta_2, 1 \rangle = \{F_a^* \beta_1, F_a^* \beta_2\} \in F_a^* \alpha = F_a^* \alpha$.

Thus

$$\langle x, y \rangle \in F_a' \alpha \times F_a' \alpha \rightarrow (\exists \beta < \alpha) (\exists \gamma < \alpha) [x = F_a' \beta \wedge y = F_a' \gamma].$$

If $\eta = J' \langle \beta, \beta, 1 \rangle$ and $v = J' \langle \beta, \gamma, 1 \rangle$ then

$$[\eta < \alpha] \wedge [v < \alpha] \wedge [J' \langle \eta, v, 1 \rangle < \alpha]$$

hence

$$F_a' J' \langle \eta, v, 1 \rangle \in F_a'' \alpha = F_a' \alpha.$$

But

$$F_a' J' \langle \eta, v, 1 \rangle = \{F_a' \eta, F_a' v\} = \{\{F_a' \beta\}, \{F_a' \beta, F_a' \gamma\}\} = \langle x, y \rangle.$$

Remark. It now remains to be proved that if a is a forcing set then N_a is a model of Axioms 4 and 5. We prove Axiom 5 first. From Theorem 13.33 we must show that if

$$(\forall u \in N_a) (\forall v \in N_a) (\forall w \in N_a) [\varphi^{N_a}(u, v) \wedge \varphi^{N_a}(u, w) \rightarrow v = w]$$

then $(\forall b \in N_a) \{y \in N_a \mid (\exists x \in b) [\varphi(x, y)]^{N_a}\} \in N_a$. Since $b \in N_a \rightarrow (\exists \alpha \in \alpha_0) [b = k_\alpha]$ and $x \in b \rightarrow (\exists \alpha_1 < \alpha) [x = k_{\alpha_1}]$, our task is to prove that

$$(\forall \alpha \in \alpha_0) [\{k_\beta \mid [\beta < \alpha_0] \wedge (\exists \alpha_1 < \alpha) [\varphi(k_{\alpha_1}, k_\beta)]^{N_a}\} \in N_a].$$

Our proof procedure is to show that there is a fundamental function G on $(n+2)$ -tuples and sets b, c, b_1, \dots, b_n in N_a such that

$$G(b, c, b_1, \dots, b_n) = \{k_\beta \mid (\exists \alpha_1 < \alpha) [\varphi(k_{\alpha_1}, k_\beta)]^{N_a}\}.$$

Since N_a is closed w.r.t. the fundamental operations and since b, c, b_1, \dots, b_n are in N_a it will then follow that

$$\{k_\beta \mid (\exists \alpha_1 < \alpha) [\varphi(k_{\alpha_1}, k_\beta)]^{N_a}\} \in N_a.$$

That such a fundamental function exists follows from Theorem 18.34 if φ is a limited statement, and if $(\exists \gamma_1 \in \alpha_0) [(\exists \alpha_1 < \alpha) [\varphi(k_{\alpha_1}, k_\beta)]^{N_a} \rightarrow \beta < \gamma_1]$. We will prove that such a γ_1 does exist and that there is a limited statement $\tilde{\varphi}$ such that

$$\begin{aligned} & \{\langle k_{\alpha_1}, k_\beta \rangle \mid [a_1 \in \alpha] \wedge [\varphi(k_{\alpha_1}, k_\beta)]^{N_a}\} \\ &= \{\langle k_{\alpha_1}, k_\beta \rangle \mid [\alpha_1 \in \alpha] \wedge [\beta \in \gamma_1] \wedge \tilde{\varphi}(k_{\alpha_1}, k_\beta)\}. \end{aligned}$$

Definition 18.36. 1) $\forall' (\exists x_0) (\exists x_1) \varphi(x_0, x_1)' \in s(\alpha_0)$

$$\Phi_\varphi(\mathcal{P}, \alpha, \beta) \stackrel{\Delta}{\longleftrightarrow} [\mathcal{P} \Vdash \ulcorner \varphi(k_\alpha, k_\beta) \urcorner] \wedge [\neg \mathcal{P} \Vdash \ulcorner (\exists^\beta x_i) \varphi(k_\alpha, x_i) \urcorner']$$

2) $\forall' (\exists x_0) (\exists x_1) (\exists x_2) \psi(x_0, x_1, x_2)' \in s(\alpha_0)$

$$\Phi_\psi(\mathcal{P}, \alpha, \beta, \gamma) \stackrel{\Delta}{\longleftrightarrow} [\mathcal{P} \Vdash \ulcorner \psi(k_\alpha, k_\beta, k_\gamma) \urcorner] \wedge [\neg \mathcal{P} \Vdash \ulcorner (\exists^\gamma x_i) \psi(k_\alpha, k_\beta, x_i) \urcorner'].$$

Theorem 18.37. $\forall' (\exists x_0) (\exists x_1) \varphi(x_0, x_1)' \in s(\alpha_0)$.

1) $\Phi_\varphi(\mathcal{P}, \alpha, \beta)$ is \mathcal{M}_0 -definable.

2) $\Phi_\varphi(\mathcal{P}, \alpha, \beta) \longleftrightarrow [\mathcal{P} \Vdash \ulcorner \varphi(k_\alpha, k_\beta) \urcorner] \wedge (\forall \gamma < \beta) [\neg \mathcal{P} \Vdash \ulcorner \varphi(k_\alpha, k_\gamma) \urcorner].$

Theorem 18.38. $\forall (\exists x_0) (\exists x_1) (\exists x_2) \psi(x_0, x_1, x_2)' \in s(\alpha_0)$.

- 1) $\Phi_\psi(\mathcal{P}, \alpha, \beta, \gamma)$ is \mathcal{M}_0 -definable.
- 2) $\Phi_\psi(\mathcal{P}, \alpha, \beta, \gamma)$
 $\longleftrightarrow [\mathcal{P} \Vdash \ulcorner \varphi(k_\alpha, k_\beta, k_\gamma) \urcorner] \wedge (\forall \tau < \gamma) [\neg \mathcal{P} \Vdash \ulcorner \varphi(k_\alpha, k_\beta, k_\tau) \urcorner]$.

The proofs are left to the reader.

Definition 18.39.

- 1) $\Gamma_\varphi(\alpha) \triangleq \cup \{\beta + 1 \mid (\exists \mathcal{P}) (\exists \alpha_1 < \alpha) \Phi_\varphi(\mathcal{P}, \alpha_1, \beta)\}$.
- 2) $\Gamma_\psi(\alpha, \beta) \triangleq \cup \{\gamma + 1 \mid (\exists \mathcal{P}) (\exists \alpha_1 < \alpha) (\exists \beta_1 < \beta) \Phi_\psi(\mathcal{P}, \alpha_1, \beta_1, \gamma)\}$.

Theorem 18.40. 1) $\forall (\exists x_0) (\exists x_1) \varphi(x_0, x_1)' \in s(\alpha_0)$

$$\alpha \in \alpha_0 \rightarrow \Gamma_\varphi(\alpha) \in \alpha_0 .$$

2) $\forall (\exists x_0) (\exists x_1) (\exists x_2) \psi(x_0, x_1, x_2)' \in s(\alpha_0)$

$$\alpha, \beta \in \alpha_0 \rightarrow \Gamma_\psi(\alpha, \beta) \in \alpha_0 .$$

Proof. 1) From the definition of $\Phi_\varphi(\mathcal{P}, \alpha, \beta)$ we see that

$$(\exists \mathcal{P}) \Phi_\varphi(\mathcal{P}, \alpha, \beta_1) \wedge (\exists \mathcal{P}) \Phi_\varphi(\mathcal{P}, \alpha, \beta_2) \rightarrow \beta_1 = \beta_2 .$$

Since $\Phi_\varphi(\mathcal{P}, \alpha, \beta)$ is \mathcal{M}_0 -definable and since \mathcal{M}_0 is a model of ZF and in particular a model of Axiom 5 it follows that

$$(\forall \alpha \in \alpha_0) [\{\beta + 1 \mid (\exists \mathcal{P}) (\exists \alpha_1 \in \alpha) \Phi_\varphi(\mathcal{P}, \alpha_1, \beta)\} \in \mathcal{M}_0]$$

then from the axiom of unions

$$\cup \{\beta + 1 \mid (\exists \mathcal{P}) (\exists \alpha_1 \in \alpha) \Phi_\varphi(\mathcal{P}, \alpha_1, \beta)\} \in \mathcal{M}_0 .$$

But the union of a set of ordinals is an ordinal. Hence

$$(\forall \alpha \in \alpha_0) [\Gamma_\varphi(\alpha) \in \alpha_0] .$$

2) The proof is left to the reader.

Definition 18.41. If a is a forcing set and $\alpha, \beta \in \alpha_0$ then

- 1) $C_\varphi(\alpha) = \{k_\beta \mid (\exists \alpha_1 < \alpha) [N_a \Vdash \ulcorner \varphi(k_{\alpha_1}, k_\beta) \urcorner]\}$.
- 2) $C_\psi(\alpha, \beta) = \{k_\gamma \mid (\exists \alpha_1 < \alpha) (\exists \beta_1 < \beta) [N_a \Vdash \ulcorner \psi(k_{\alpha_1}, k_{\beta_1}, k_\gamma) \urcorner]$
 $\wedge (\forall \tau < \gamma) \neg [N_a \Vdash \ulcorner \psi(k_{\alpha_1}, k_{\beta_1}, k_\tau) \urcorner]\}$.

Theorem 18.42. If a is a forcing set and $\alpha, \beta \in \alpha_0$ then

- 1) $N_a \Vdash \ulcorner (\forall x_0) (\forall x_1) (\forall x_2) [\varphi(x_0, x_1) \wedge \varphi(x_0, x_2) \rightarrow x_1 = x_2] \urcorner$
 $\rightarrow (\exists \gamma \in \alpha_0) [C_\varphi(\alpha) \subseteq k_\gamma]$.
- 2) $N_a \Vdash \ulcorner (\forall x_0) (\forall x_1) (\exists x_2) \psi(x_0, x_1, x_2) \urcorner \rightarrow (\exists \gamma \in \alpha_0) [C_\psi(\alpha, \beta) \subseteq k_\gamma]$.

Proof. 1) If $k_\beta \in C_\varphi(\alpha)$ then $(\exists \alpha_1 < \alpha) [N_a \models \ulcorner \varphi(k_{\alpha_1}, k_\beta) \urcorner]$. Since by hypothesis

$$N_a \models \ulcorner (\forall x_0) (\forall x_1) (\forall x_2) [\varphi(x_0, x_1) \wedge \varphi(x_0, x_2) \rightarrow x_1 = x_2] \urcorner$$

it follows that

$$N_a \models \ulcorner \varphi(k_{\alpha_1}, k_\beta) \urcorner \wedge (\forall \beta_1 < \beta) \neg [N \models \ulcorner \varphi(k_{\alpha_1}, k_{\beta_1}) \urcorner]$$

therefore by Theorem 18.24

$$(\exists \mathcal{P}) (\exists \alpha_1 < \alpha) \Phi_\varphi(\mathcal{P}, \alpha_1, \beta)$$

then $\beta \in \Gamma_\varphi(\alpha)$. If $\gamma = \text{cov}(\Gamma_\varphi(\alpha))$ then $k_\beta \in k_\gamma$ and $C_\varphi(\alpha) \subseteq k_\gamma$.

2) The proof is left to the reader.

Remark. In our next result we use the fact that every wff $\varphi(x_0, x_1)$ has a prenex normal form, i.e. $\varphi(x_0, x_1)$ is equivalent to a wff $(Q_2 x_2) \dots (Q_n x_n) \psi(x_0, x_1, \dots, x_n)$ where $\psi(x_0, \dots, x_n)$ is quantifier free and each Q_i is a quantifier \forall or \exists .

Theorem 18.43. *If a is a forcing set, if*

$$N_a \models \ulcorner (\forall x_0) (\forall x_1) (\forall x_2) [\varphi(x_0, x_1) \wedge \varphi(x_0, x_2) \rightarrow x_1 = x_2] \urcorner,$$

if $\varphi(x_0, x_1)$ has prenex normal form

$$(Q_2 x_2) \dots (Q_n x_n) \psi(x_0, x_1, \dots, x_n)$$

then for each α in α_0 there exist ordinals $\gamma_1, \dots, \gamma_n$ in α_0 such that

$$(1) \quad (\forall x_0 \in k_\alpha) [(\exists x_1) N_a \models \ulcorner \varphi(x_0, x_1) \urcorner \rightarrow (\exists x_1 \in k_{\gamma_1}) [N_a \models \ulcorner \varphi(x_0, x_1) \urcorner]].$$

$$(2) \quad [\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1}] \rightarrow [[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow \tilde{\varphi}(x_0, x_1)]$$

where $\tilde{\varphi}(x_0, x_1)$ is the limited formula

$$(Q_2 x_2 \in k_{\gamma_2}) \dots (Q_n x_n \in k_{\gamma_n}) \psi(x_0, x_1, \dots, x_n).$$

Proof. (1) By Theorem 18.42 $(\exists \gamma_1 \in \alpha_0) [C_\varphi(\alpha) \subseteq k_{\gamma_1}]$. Furthermore $(\forall x_0 \in k_\alpha) (\exists \alpha_1 < \alpha) [x_0 = k_{\alpha_1}]$. If $(\exists \beta) [N_a \models \ulcorner \varphi(k_{\alpha_1}, k_\beta) \urcorner]$, then $k_\beta \in C_\varphi(\alpha) \subseteq k_{\gamma_1}$ thus

$$(\forall x_0 \in k_\alpha) (\exists x_1 \in k_{\gamma_1}) [N_a \models \ulcorner \varphi(x_0, x_1) \urcorner].$$

(2) By induction we will prove that for each $m, 1 \leq m \leq n$, there exist ordinals $\gamma_2, \dots, \gamma_m$ such that if $\langle x_0, x_1 \rangle \in k_\alpha \times k_\gamma$,

$$[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_m x_m \in k_{\gamma_m}) [\psi_m(x_0, \dots, x_n)]^{N_a}$$

where $\psi_m(x_0, \dots, x_m)$ is $(Q_{m+1} x_{m+1}) \dots (Q_n x_n) \psi(x_0, \dots, x_n)$.

If $m = 1$ the conclusion is obvious i.e.

$$[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow [\varphi(x_0, x_1)]^{N_a}.$$

Assume the result true for $m - 1$, that is, assume that there exist ordinals $\gamma_2, \dots, \gamma_{m-1}$ such that for $\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1}$

$$[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) [\psi_{m-1}(x_0 \dots x_{m-1})]^{N_a}.$$

If Q_m is \forall that is if $\psi_{m-1}(x_0, \dots, x_{m-1})$ is $(\forall x_m) \psi_m(x_0, \dots, x_m)$ and if

$$\tilde{\psi}_m(x_0, x_1, y) \stackrel{\Delta}{\longleftrightarrow} [y = 0 \wedge (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\forall x_m) \psi_m(x_0, \dots, x_m)] \vee (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) [\neg \psi_m(x_0, \dots, x_{m-1}, y)]$$

then

$$[\tilde{\psi}_m(x_0, x_1, y)]^{N_a} \longleftrightarrow [y = 0 \wedge (Q_2 x_2 \in k_{\gamma_2}) \dots \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\forall x_m \in N_a) [\psi_m(x_0, \dots, x_m)]^{N_a}] \vee (Q_2 x_2 \in k_{\gamma_2}) \dots \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) [\neg \psi_m(x_0, \dots, x_{m-1}, y)]^{N_a}.$$

Either $(Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\forall x_m \in N_a) [\psi_m(x_0, \dots, x_m)]^{N_a}$ or

$$(Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\exists x_m \in N_a) [\neg \psi_m(x_0, \dots, x_m)]^{N_a}.$$

In either case, since $0 \in N_a$ it follows that

$$(\forall x_0 \in N_a) (\forall x_1 \in N_a) (\exists x_m \in N_a) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a}$$

i.e. $N_a \models (\forall x_0) (\forall x_1) (\exists x_m) \tilde{\psi}(x_0, x_1, x_m)$. Then from Theorem 18.42

$$(\exists \gamma_m \in \alpha_0) [C_{\tilde{\psi}_m}(\alpha, \gamma_1) \subseteq k_{\gamma_m}].$$

Since $(\forall x_0 \in N_a) (\forall x_1 \in N_a) (\exists x_m \in N_a) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a}$ it follows that

$$(\forall \alpha_1 < \alpha) (\forall \beta_1 < \gamma_1) (\exists \tau) [\tilde{\psi}_m(k_{\alpha_1}, k_{\beta_1}, k_\tau)]^{N_a}.$$

Therefore there is a smallest such τ , i.e.

$$(\forall \alpha_1 < \alpha) (\forall \beta_1 < \gamma_1) (\exists \tau) [[\tilde{\psi}_m(k_{\alpha_1}, k_{\beta_1}, k_\tau)]^{N_a} \wedge (\forall \tau_1 < \tau) [\neg \tilde{\psi}_m(k_{\alpha_1}, k_{\beta_1}, k_{\tau_1})]^{N_a}]$$

then $k_\tau \in C_{\tilde{\psi}_m}(\alpha, \gamma_1) \subseteq k_{\gamma_m}$. Thus

$$(\forall x_0 \in k_\alpha) (\forall x_1 \in k_{\gamma_1}) (\exists x_m \in k_{\gamma_m}) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a}$$

that is $\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1} \rightarrow (\exists x_m \in k_{\gamma_m}) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a}$. But from the definition of $\tilde{\psi}_m$

$$(\exists x_m \in k_{\gamma_m}) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a} \wedge (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\forall x_m \in k_{\gamma_m}) [\psi_m(x_0, \dots, x_m)]^{N_a} \rightarrow (Q_2 x_2 \in k_{\gamma_2}) \dots \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\forall x_m \in N_a) [\psi_m(x_0, \dots, x_m)]^{N_a}.$$

Thus if $\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1}$

$$\begin{aligned} & [(Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\forall x_m \in N_a) [\psi_m(x_0, \dots, x_m)]^{N_a} \\ \longleftrightarrow & (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\forall x_m \in k_{\gamma_m}) [\psi_m(x_0, \dots, x_m)]^{N_a}. \end{aligned}$$

From the induction hypothesis we conclude that if $\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1}$ then

$$[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_m x_m \in k_{\gamma_m}) [\psi_m(x_0, \dots, x_m)]^{N_a}.$$

If Q_m is \exists that is if $\psi_{m-1}(x_0, \dots, x_{m-1})$ is $(\exists x_m) \psi_m(x_0, \dots, x_m)$ and if

$$\begin{aligned} & \tilde{\psi}_m(x_0, x_1, y) \stackrel{\Delta}{\longleftrightarrow} [y = 0 \wedge (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) \\ \neg(\exists x_m) \psi_m(x_0, \dots, x_m)] \vee & (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) \psi_m(x_0, \dots, x_{m-1}, y) \end{aligned}$$

then

$$\begin{aligned} [\tilde{\psi}_m(x_0, x_1, y)]^{N_a} \longleftrightarrow & [y = 0 \wedge (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) \\ & \neg(\exists x_m \in N_a) [\psi_m(x_0, \dots, x_m)]^{N_a} \vee (Q_2 x_2 \in k_{\gamma_2}) \dots \\ & \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) [\psi_m(x_0, \dots, x_{m-1}, y)]^{N_a}. \end{aligned}$$

Either

$$(Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) \neg(\exists x_m \in N_a) [\psi_m(x_0, \dots, x_m)]^{N_a}$$

or

$$(Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\exists x_m \in N_a) [\psi_m(x_0, \dots, x_m)]^{N_a}.$$

In either case, since $0 \in N_a$

$$(\forall x_0 \in N_a) (\forall x_1 \in N_a) (\exists x_m \in N_a) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a},$$

i.e.

$$N_a \models '(\forall x_0) (\forall x_1) (\exists x_m) \tilde{\psi}_m(x_0, x_1, x_m)'$$

Again by Theorem 18.42

$$(\exists \gamma_m \in \alpha_0) [C_{\tilde{\psi}_m}(\alpha, \gamma_1) \subseteq k_{\gamma_m}].$$

Since $(\forall x_0 \in N_a) (\forall x_1 \in N_a) (\exists x_m \in N_a) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a}$ it follows that

$$(\forall \alpha_1 < \alpha) (\forall \beta_1 < \gamma_1) (\exists \tau) [\tilde{\psi}_m(k_{\alpha_1}, k_{\beta_1}, k_\tau)]^{N_a}.$$

There is then a smallest such τ and for this τ we have

$$k_\tau \in C_{\tilde{\psi}_m}(\alpha, \gamma_1) \subseteq k_{\gamma_m}$$

that is

$$(\forall x_0 \in k_\alpha) (\forall x_1 \in k_{\gamma_1}) (\exists x_m \in k_{\gamma_m}) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a}.$$

Thus

$$\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1} \rightarrow (\exists x_m \in k_{\gamma_m}) [\tilde{\psi}_m(x_0, x_1, x_m)]^{N_a}.$$

As before it follows from the definition of $\tilde{\psi}_m$ that if $\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1}$

$$\begin{aligned} & [(Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\exists x_m \in N_a) [\psi_m(x_0, \dots, x_m)]]^{N_a} \\ \longleftrightarrow & (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_{m-1} x_{m-1} \in k_{\gamma_{m-1}}) (\exists x_m \in k_{\gamma_m}) [\psi_m(x_0, \dots, x_m)]^{N_a}. \end{aligned}$$

Then from the induction hypothesis if $\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1}$

$$[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_m x_m \in k_{\gamma_m}) [\psi_m(x_0, \dots, x_m)]^{N_a}.$$

In particular for $m = n$ we have that there exist ordinals $\gamma_1, \dots, \gamma_n$ such that if $\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1}$

$$[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow (Q_2 x_2 \in k_{\gamma_2}) \dots (Q_n x_n \in k_{\gamma_n}) \psi(x_0, \dots, x_n).$$

Theorem 18.44. *If a is a forcing set then N_a is a standard transitive model of the Axiom Schema of Replacement.*

Proof. If $N_a \models [(\forall x_0) (\forall x_1) (\forall x_2) \varphi(x_0, x_1) \wedge \varphi(x_0, x_2) \rightarrow x_1 = x_2]$, then $(\forall b \in N_a) (\exists \alpha \in \alpha_0) [b = k_\alpha]$ and

$$\begin{aligned} \{y \in N_a \mid (\exists x \in b) [\varphi(x, y)]^{N_a}\} &= \{k_\beta \mid (\exists x_0 \in k_\alpha) [\varphi(x_0, k_\beta)]^{N_a}\} \\ &= \{k_\beta \mid (\exists \alpha_1 < \alpha) [N_a \models \varphi(k_{\alpha_1}, k_\beta)]\} = C_\varphi(\alpha). \end{aligned}$$

By Theorem 18.42 $(\exists \gamma_1 \in \alpha_0) [C_\varphi(\alpha) \subseteq k_{\gamma_1}]$. Also there exist ordinals $\gamma_2, \dots, \gamma_n$ in α_0 and a wff $\tilde{\varphi}(x_0, x_1)$ each of whose quantifiers is restricted to one of the sets $k_{\gamma_2}, \dots, k_{\gamma_n}$, such that

$$\langle x_0, x_1 \rangle \in k_\alpha \times k_{\gamma_1} \rightarrow [[\varphi(x_0, x_1)]^{N_a} \longleftrightarrow \tilde{\varphi}(x_0, x_1)].$$

Since $J^* \langle 0, \tau, 0 \rangle < \bar{\tau}^+$ we have as a theorem in ZF

$$(\forall \delta) (\exists \gamma) [\delta < \gamma \wedge (\forall \tau < \gamma) [J^* \langle 0, \tau, 0 \rangle < \gamma] \wedge K_3^* \gamma = 0]$$

Relativizing to \mathcal{M}_0

$$(\forall \delta \in \alpha_0) (\exists \gamma \in \alpha_0) [\delta < \gamma \wedge (\forall \tau < \gamma) [J^* \langle 0, \tau, 0 \rangle < \gamma] \wedge K_3^* \gamma = 0].$$

In particular since $\text{cov}(\max(\alpha, \gamma_1, \dots, \gamma_n)) \in \alpha_0$

$$(\exists \gamma \in \alpha_0) [\text{cov}(\max(\alpha, \gamma_1, \dots, \gamma_n)) < \gamma \wedge (\forall \tau < \gamma) [J^* \langle 0, \tau, 0 \rangle < \gamma] \wedge K_3^* \gamma = 0]$$

Then

$$\{y \in N_a \mid (\exists x \in b) [\varphi(x, y)]^{N_a}\} = \{k_\beta \mid k_\beta \in k_\gamma \wedge (\exists x_0 \in k_\alpha) \tilde{\varphi}(x_0, k_\beta)\}$$

Furthermore

$$k_\alpha \subseteq k_\gamma, k_{\gamma_1} \subseteq k_\gamma, \dots, k_{\gamma_n} \subseteq k_\gamma$$

and from Theorems 15.21 and 18.35

$$\text{Tr}(k_\gamma) \wedge k_\gamma \times k_\gamma \subseteq k_\gamma.$$

Then by Theorem 18.34 there is a fundamental function G on $(n+2)$ -tuples such that

$$G(k_\gamma, k_\alpha, k_{\gamma_1}, \dots, k_{\gamma_n}) = \{k_\beta \mid k_\beta \in k_\gamma \wedge (\exists x_0 \in k_\alpha) \tilde{\varphi}(x_0, k_\beta)\}.$$

Since N_a is closed w.r.t. the fundamental operations, and since $k_\gamma, k_\alpha, k_{\gamma_1}, \dots, k_{\gamma_n}$ are in N_a it follows that

$$(\forall b \in N_a) [\{y \in N_a \mid (\exists x \in b) [\varphi(x, y)]^{N_a}\} \in N_a]$$

and hence N_a is a model of Axiom 5.

Remark. In order to show that N_a is a model of the Axiom of Powers, we prove two properties of F_a^α for $\alpha < \alpha_0$. We first define, for fixed $\alpha < \alpha_0$, three functions A_1 , A_2 , and B as follows.

$$\begin{aligned} A_1('k_\beta') &= \{\mathcal{P} \mid \mathcal{P} \Vdash 'k_\beta \subseteq k_\alpha'\}, & \beta < \alpha_0 \\ A_2('k_\beta') &= \{\langle \mathcal{P}, 'k_\gamma' \rangle \mid \mathcal{P} \Vdash 'k_\gamma \in k_\beta' \wedge \gamma < \alpha\}, & \beta < \alpha_0 \\ B('k_\beta') &= \langle A_1('k_\beta'), A_2('k_\beta') \rangle, & \beta < \alpha_0. \end{aligned}$$

By Theorem 18.15, $A_1('k_\beta')$, $A_2('k_\beta')$ and $B('k_\beta')$ are sets in \mathcal{M}_0 and A_1 , A_2 , and B are \mathcal{M}_0 -definable.

Theorem 18.45. 1) If $B('k_{\beta_1}') = B('k_{\beta_2}')$ and $F_a^{\beta_1} \subseteq F_a^\alpha$ then $F_a^{\beta_1} = F_a^{\beta_2}$.

2) There is a $\beta_0 \in \mathcal{M}_0$ such that if $\beta < \alpha_0$ and $F_a^\beta \subseteq F_a^\alpha$ then $(\exists \gamma \in \beta_0) [F_a^\gamma = F_a^\beta]$.

Proof. 1) Let \mathcal{P}_n , $n=1, 2, \dots$, be a complete sequence of forcing conditions that determines a . Since $F_a^\beta \subseteq F_a^\alpha$ there exists a \mathcal{P}_n in the complete sequence of forcing conditions, such that $\mathcal{P}_n \Vdash 'k_{\beta_1} \subseteq k_\alpha'$. Since by hypothesis $B('k_{\beta_1}') = B('k_{\beta_2}')$ it follows that $\mathcal{P}_n \Vdash 'k_{\beta_2} \subseteq k_\alpha'$ and hence $F_a^{\beta_2} \subseteq F_a^\alpha$, by Theorem 18.24.

Suppose $F_a^{\beta_1} \neq F_a^{\beta_2}$. Without loss of generality, we may assume that $F_a^{\beta_1} - F_a^{\beta_2} \neq 0$, i.e. $(\exists \gamma < \alpha) [F_a^\gamma \in F_a^{\beta_1} \wedge F_a^\gamma \notin F_a^{\beta_2}]$. Since $F_a^\gamma \in F_a^{\beta_1}$ there is a \mathcal{P}_n such that $\mathcal{P}_n \Vdash 'k_\gamma \in k_{\beta_1}'$. Since $A_2('k_{\beta_1}') = A_2('k_{\beta_2}')$, $\mathcal{P}_n \Vdash 'k_\gamma \in k_{\beta_2}'$ and hence $F_a^\gamma \in F_a^{\beta_2}$. This is a contradiction.

2) Since A_1 is \mathcal{M}_0 -definable

$$\{A_1('k_\beta') \mid \beta < \alpha_0\} \subseteq [\mathcal{P}(\text{cond})]^{\mathcal{M}_0}.$$

Similarly

$$\{A_2('k_\beta') \mid \beta < \alpha_0\} \subseteq [\mathcal{P}(\text{cond} \times \{k_\gamma \mid \gamma < \alpha\})]^{\mathcal{M}_0}.$$

Therefore $\bar{B} \triangleq \{B('k_\beta') \mid \beta < \alpha_0\}$ is a set in \mathcal{M}_0 . We then define f on \bar{B} as follows

$$f(b) = \mu_\beta(B('k_\beta') = b), \quad b \in \bar{B}$$

If $\beta_0 = \sup \{f(b) \mid b \in \bar{B}\}$ then since \bar{B} is a set in \mathcal{M}_0 , $\beta_0 \in \mathcal{M}_0$. If $\beta < \alpha_0$ and $F_a^\beta \subseteq F_a^\alpha$ then $(\exists \gamma \in \beta_0) [B('k_\beta') = B('k_\gamma')]$. Hence by 1) $F_a^\beta = F_a^\gamma$.

Theorem 18.46. *If a is a forcing set, then N_a is a standard transitive model of the Axiom of Powers.*

Proof. $(\forall b \in N_a) (\exists \alpha \in \alpha_0) [b = F_a' \alpha]$. Let β_0 be an ordinal with the property prescribed by Theorem 18.45,2) and let $\beta = \text{cov}(\beta_0)$. Then $F_a' \beta = F_a'' \beta$, and by Theorem 18.45,2) $[\mathcal{P}(F_a' \alpha)]^{N_a} \subseteq F_a' \beta$. Therefore

$$[\mathcal{P}(F_a' \alpha)]^{N_a} = \{x \in F_a' \beta \mid x \subseteq F_a' \alpha\}.$$

Since N_a is a model of the Axiom of Replacement, it follows that $[\mathcal{P}(F_a' \alpha)]^{N_a} \in N_a$, i.e. N_a is a model of the Axiom of Powers.

Theorem 18.47. *If a is a forcing set then N_a is a standard transitive model of $\text{ZF} + \text{AC} + \text{GCH} + V = L_a$.*

Proof. From Theorems 17.7, 18.44 and 18.46 it follows that N_a is a standard transitive model of ZF. Therefore $\mathcal{M}_0 \subseteq N_a$. Then

$$\alpha \in \mathcal{M}_0 \rightarrow \alpha \in N_a.$$

By Theorem 17.9

$$\alpha \in N_a \rightarrow \alpha \in \mathcal{M}_0.$$

Thus $\text{On}^{N_a} = \alpha_0$ and hence

$$L_a^{N_a} = \{x \mid (\exists \alpha \in \alpha_0) [x = F_a' \alpha]\} = N_a.$$

Therefore N_a is a standard transitive model of $\text{ZF} + \text{AC} + \text{GCH} + V = L_a$.

Theorem 18.48. *There exists an a such that N_a is a standard transitive model of $\text{ZF} + \text{AC} + \text{GCH} + V \neq L$.*

Proof. By Theorem 18.23 there exists a forcing set a not in \mathcal{M}_0 . Since $a \in N_a$ it follows that $N_a \neq \mathcal{M}_0$. But then

$$L^{N_a} = \{x \mid (\exists \alpha \in \alpha_0) [x = F' \alpha]\} = \mathcal{M}_0 \neq N_a = V^{N_a}.$$

Thus, with Theorem 18.47, N_a is a standard transitive model of $\text{ZF} + \text{AC} + \text{GCH} + V \neq L$.

Remark. Only in the proof of Theorem 18.47 have we used the fact that \mathcal{M}_0 is a submodel of every standard transitive model of ZF. We next show that Theorem 18.47 can also be proved without using this fact. By proving

$$\alpha \in \mathcal{M}_0 \rightarrow \alpha \in N_a$$

without using the minimality of \mathcal{M}_0 we will establish a more general theory, for it will then follow that all of the results of the section hold if \mathcal{M}_0 is replaced by any \mathcal{M} that is a countable standard transitive model of $\text{ZF} + \text{AC} + \text{GCH} + V = L$. For this proof we introduce a collection of functions similar to the fundamental terms except that they are generated

from J -maps instead of fundamental operations. For this reason we call them J -functions.

Definition 18.49. H is a J -function on n -tuples of type zero iff $H \in \mathcal{F}_n \text{On}^n$ and

$$(\exists i \leq n) (\forall \alpha_1) \dots (\forall \alpha_n) [H(\alpha_1, \dots, \alpha_n) = \alpha_i \wedge i \geq 1].$$

H is a J -function on n -tuples of type $m + 1$ iff H is a J -function on n -tuples of type m or there exist J -functions on n -tuples, H_1 and H_2 , of type m and an integer $0 \leq i \leq n$ such that

$$(\forall \alpha_1) \dots (\forall \alpha_n) H(\alpha_1, \dots, \alpha_n) = J^{\langle H_1(\alpha_1, \dots, \alpha_n), H_2(\alpha_1, \dots, \alpha_n), i \rangle}.$$

Theorem 18.50. *If H is a J -function on n -tuples then*

$$\min(\alpha_1, \dots, \alpha_n) \leq H(\alpha_1, \dots, \alpha_n).$$

Proof. The result follows by induction on the type of H . Details are left to the reader.

Theorem 18.51. *For each fundamental function G on n -tuples there exists a J -function H on n -tuples for which*

$$\omega < \alpha_1 \wedge \dots \wedge \omega < \alpha_n \rightarrow G(F_a^{\omega} \alpha_1, \dots, F_a^{\omega} \alpha_n) = F_a^{\omega} H(\alpha_1, \dots, \alpha_n).$$

Proof (by induction on the type of G). If G is of type zero then

$$(\exists i \leq n) (\forall a_1) \dots (\forall a_n) [G(a_1, \dots, a_n) = a_i \wedge i \geq 1].$$

If we define H by

$$H(\alpha_1, \dots, \alpha_n) = \alpha_i$$

then H is a J -function on n -tuples and

$$G(F_a^{\omega} \alpha_1, \dots, F_a^{\omega} \alpha_n) = F_a^{\omega} \alpha_i = F_a^{\omega} H(\alpha_1, \dots, \alpha_n).$$

As our induction hypothesis we assume the result true for all fundamental functions of type m . If G is a fundamental function of type $m + 1$ then G is a fundamental function of type m , in which case the result follows from the induction hypothesis, or there exist fundamental functions G_1 and G_2 of type m and a fundamental operation \mathcal{F}_i such that

$$G(a_1, \dots, a_n) = \mathcal{F}_i(G_1(a_1, \dots, a_n), G_2(a_1, \dots, a_n)).$$

From the induction hypothesis there exist J -functions H_1 and H_2 on n -tuples for which

$$\begin{aligned} [\omega < \alpha_1] \wedge \dots \wedge [\omega < \alpha_n] \rightarrow G_1(F_a^{\omega} \alpha_1, \dots, F_a^{\omega} \alpha_n) &= F_a^{\omega} H_1(\alpha_1, \dots, \alpha_n) \\ \wedge G_2(F_a^{\omega} \alpha_1, \dots, F_a^{\omega} \alpha_n) &= F_a^{\omega} H_2(\alpha_1, \dots, \alpha_n). \end{aligned}$$

From Theorem 18.50

$$[\omega < \alpha_1] \wedge \cdots \wedge [\omega < \alpha_n] \rightarrow [\omega < H_1(\alpha_1, \dots, \alpha_n)] \wedge [\omega < H_2(\alpha_1, \dots, \alpha_n)].$$

Therefore if

$$H(\alpha_1, \dots, \alpha_n) = J^* \langle H_1(\alpha_1, \dots, \alpha_n), H_2(\alpha_1, \dots, \alpha_n), i \rangle$$

then $\omega + 1 < H(\alpha_1, \dots, \alpha_n)$ and hence

$$\begin{aligned} G(F_a^* \alpha_1, \dots, F_a^* \alpha_n) &= \mathcal{F}_i(G_1(F_a^* \alpha_1, \dots, F_a^* \alpha_n), G_2(F_a^* \alpha_1, \dots, F_a^* \alpha_n)) \\ &= \mathcal{F}_i(F_a^* H_1(\alpha_1, \dots, \alpha_n), F_a^* H_2(\alpha_1, \dots, \alpha_n)) \\ &= F_a^* H(\alpha_1, \dots, \alpha_n). \end{aligned}$$

Definition 18.52.

- 1) $g_1(\alpha, \beta, \gamma) \triangleq J \langle \alpha, J^* \langle J^* \langle \alpha, \beta, 3 \rangle, \gamma, 3 \rangle, 3 \rangle.$
- 2) $g_2(\alpha, \beta) \triangleq g_1(\text{cov}(\max(\alpha, \beta)), \alpha, \beta).$
- 3) $g_3(\alpha) = g_2(\alpha, J^* \langle \alpha, \alpha, 1 \rangle).$

Theorem 18.53.

- 1) $\max(\alpha, \beta) \geq \omega \rightarrow g_1(\alpha, \beta, \gamma) > \omega + 1.$
- 2) $\max(\alpha, \beta) \geq \omega \rightarrow g_2(\alpha, \beta) > \omega + 1.$
- 3) $\alpha \geq \omega \rightarrow g_3(\alpha) > \omega + 1.$

The proofs are left to the reader.

Theorem 18.54. *If $a \subseteq \omega$ then*

- 1) $\max(\alpha, \beta) \geq \omega \rightarrow F_a^* g_1(\alpha, \beta, \gamma) = F_a^* \alpha - [[F_a^* \alpha - F_a^* \beta] - F_a^* \gamma],$
- 2) $\max(\alpha, \beta) \geq \omega \rightarrow F_a^* g_2(\alpha, \beta) = F_a^* \alpha \cup F_a^* \beta,$
- 3) $\alpha \geq \omega \rightarrow F_a^* g_3(\alpha) = F_a^* \alpha \cup \{F_a^* \alpha\}.$

Proof. 1) If $\max(\alpha, \beta) \geq \omega$ then

$$J^* \langle \alpha, \beta, 3 \rangle > \omega + 1, J^* \langle J^* \langle \alpha, \beta, 3 \rangle, \gamma, 3 \rangle > \omega + 1$$

and

$$g_1(\alpha, \beta, \gamma) > \omega + 1.$$

Therefore

$$\begin{aligned} F_a^* g_1(\alpha, \beta, \gamma) &= F_a^* \alpha - F_a^* J^* \langle J^* \langle \alpha, \beta, 3 \rangle, \gamma, 3 \rangle \\ &= F_a^* \alpha - [F_a^* J^* \langle \alpha, \beta, 3 \rangle - F_a^* \gamma] \\ &= F_a^* \alpha - [[F_a^* \alpha - F_a^* \beta] - F_a^* \gamma]. \end{aligned}$$

- 2) If $\gamma = \text{cov}(\max(\alpha, \beta))$ then

$$[F_a^* \alpha \in F_a^* \gamma = F_a^* \gamma] \wedge [F_a^* \beta \in F_a^* \gamma = F_a^* \gamma].$$

Since $F_a^* \gamma$ is transitive

$$[F_a^* \alpha \subseteq F_a^* \gamma] \wedge [F_a^* \beta \subseteq F_a^* \gamma].$$

Therefore, from 1) above

$$\begin{aligned} F_a^* g_2(\alpha, \beta) &= F_a^* g_1(\gamma, \alpha, \beta) \\ &= F_a^* \gamma - [[F_a^* \gamma - F_a^* \alpha] - F_a^* \beta] \\ &= F_a^* \alpha \cup F_a^* \beta. \end{aligned}$$

3) From 2) above

$$\begin{aligned} F_a^* g_3(\alpha) &= F_a^* g_2(\alpha, J^* \langle \alpha, \alpha, 1 \rangle) \\ &= F_a^* \alpha \cup F_a^* J^* \langle \alpha, \alpha, 1 \rangle \\ &= F_a^* \alpha \cup \{F_a^* \alpha\}. \end{aligned}$$

Theorem 18.55. $J^* \langle 0, \alpha, 0 \rangle < g_3(\alpha)$.

Proof. If $\gamma = \text{cov}(\max(\alpha, J^* \langle \alpha, \alpha, 1 \rangle))$ then

$$\max(0, \alpha) < \max(\gamma, J^* \langle J^* \langle \gamma, \alpha, 3 \rangle, J^* \langle \alpha, \alpha, 1 \rangle, 3 \rangle).$$

Therefore

$$\begin{aligned} J^* \langle 0, \alpha, 0 \rangle &< J^* \langle \gamma, J^* \langle J^* \langle \gamma, \alpha, 3 \rangle, J^* \langle \alpha, \alpha, 1 \rangle, 3 \rangle, 3 \rangle \\ &= g_1(\gamma, \alpha, J^* \langle \alpha, \alpha, 1 \rangle) \\ &= g_2(\alpha, J^* \langle \alpha, \alpha, 1 \rangle) \\ &= g_3(\alpha). \end{aligned}$$

Theorem and Definition 18.56. *There exists a unary J-function g_4 with the property that*

$$[\omega < \alpha \in K_{II}] \wedge (\forall \beta < \alpha) [J^* \langle 0, \beta, 0 \rangle < \alpha] \rightarrow F_a^* g_4(\alpha) = F_a^* \alpha \cap \text{On}.$$

Proof. By Theorems 15.21 and 18.35

$$[\omega < \alpha \in K_{II}] \wedge (\forall \beta < \alpha) [J^* \langle 0, \beta, 0 \rangle < \alpha]$$

implies

$$\text{Tr}(F_a^* \alpha) \wedge [F_a^* \alpha \times F_a^* \alpha \subseteq F_a^* \alpha].$$

From Theorem 18.34 it then follows that there is a unary fundamental function G such that

$$G(F_a^* \alpha) = \{x \mid [x \in F_a^* \alpha] \wedge \text{Ord}(x)\}.$$

By Theorem 18.51 there exists a J -function g_4 such that

$$F_a^* g_4 \alpha = \{x \mid [x \in F_a^* \alpha] \wedge \text{Ord}(x)\} = F_a^* \alpha \cap \text{On}.$$

Definition 18.57. $G'x = \mathcal{D}(x)$ if $\mathcal{D}(x) \leq \omega$
 $= g_3(x'(\cup \mathcal{D}(x)))$ if $\omega < \mathcal{D}(x) \in K_I$
 $= g_4(\cup \mathcal{W}(x))$ otherwise.

$$F_0 \mathcal{F}_n \text{ On } \wedge (\forall \alpha) [F_0' \alpha = G'(F_0 \Gamma \alpha)].$$

Theorem 18.58. $F_0' \alpha = \alpha$ if $\alpha \leq \omega$
 $= g_3(F_0' \beta)$ if $\omega < \alpha = \beta + 1$
 $= g_4(\cup F_0'' \alpha)$ if $\omega < \alpha \in K_{II}$.

Proof. $\mathcal{D}(F_0 \Gamma \alpha) = \alpha$. Therefore if $\alpha \leq \omega$ then

$$F_0' \alpha = G'(F_0 \Gamma \alpha) = \mathcal{D}(F_0 \Gamma \alpha) = \alpha.$$

If $\omega < \alpha = \beta + 1$ then

$$F_0' \alpha = G'(F_0 \Gamma \alpha) = g_3((F_0 \Gamma \alpha)' \beta) = g_3(F_0' \beta).$$

If $\omega < \alpha \in K_{II}$ then

$$F_0' \alpha = G'(F_0 \Gamma \alpha) = g_4(\cup (\mathcal{W}(F_0 \Gamma \alpha))) = g_4(\cup F_0'' \alpha).$$

Theorem 18.59. $\omega < \alpha \in K_{II} \rightarrow (\forall \beta < F_0'' \alpha) [J' \langle 0, \beta, 0 \rangle < F_0'' \alpha]$.

Proof. $\beta < F_0'' \alpha \rightarrow (\exists \eta < \alpha) [\beta = F_0' \eta]$

$$J' \langle 0, \beta, 0 \rangle = J' \langle 0, F_0' \eta, 0 \rangle < g_3(F_0' \eta) = F_0'(\eta + 1) < F_0'' \alpha.$$

Theorem 18.60. F_0 is a strictly monotonic ordinal function.

Proof (by induction on α). If $\beta < \alpha \leq \omega$ then $F_0' \beta = \beta < \alpha = F_0' \alpha$. If $\omega < \alpha = \eta + 1$ then

$$\beta < \alpha \rightarrow \beta < \eta \vee \beta = \eta.$$

If $\beta < \eta$ then since $\eta < \alpha$ we have by our induction hypothesis

$$F_0' \beta < F_0' \eta.$$

But

$$F_0' \alpha = g_3(F_0' \eta) > J' \langle 0, F_0' \eta, 0 \rangle \geq F_0' \eta$$

i.e.

$$F_0' \beta < F_0' \eta < F_0' \alpha.$$

If $\omega < \alpha \in K_{II}$ then

$$F_0' \alpha = g_4(\cup F_0'' \alpha).$$

But since g_4 is a J -function

$$g_4(\cup (F_0'' \alpha)) \geq \cup (F_0'' \alpha)$$

i.e.

$$\cup (F_0'' \alpha) \leq F_0' \alpha.$$

If $\beta < \alpha$ then $\beta < \beta + 1 < \alpha$ and, by the induction hypothesis, $F_0' \beta < F_0'(\beta + 1) \in F_0'' \alpha$. Thus $F_0' \beta \in \cup(F_0'' \alpha) \subseteq g_4(\cup(F_0'' \alpha)) = F_0' \alpha$. Therefore

$$\beta < \alpha \rightarrow F_0' \beta < F_0' \alpha.$$

Theorem 18.61. *If $[0 < i < 9] \wedge [\omega + 1 < J' \langle \alpha, \beta, i \rangle]$ then*

$$\text{rank}(F_a' J' \langle \alpha, \beta, i \rangle) \leq [1 + \max(\text{rank}(F_a' \alpha), \text{rank}(F_a' \beta))].$$

Proof. If $i = 1$ then $F_a' J' \langle \alpha, \beta, i \rangle = \{F_a' \alpha, F_a' \beta\}$. Hence

$$\text{rank}(F_a' J' \langle \alpha, \beta, i \rangle) = 1 + \max(\text{rank}(F_a' \alpha), \text{rank}(F_a' \beta)).$$

If $1 < i < 9$ then $F_a' J' \langle \alpha, \beta, i \rangle \subseteq F_a' \alpha$ and

$$\text{rank}(F_a' J' \langle \alpha, \beta, i \rangle) \leq \text{rank}(F_a' \alpha) < 1 + \max(\text{rank}(F_a' \alpha), \text{rank}(F_a' \beta)).$$

Definition 18.62. $\tilde{j}(\alpha) = \alpha$ if $\alpha \leq \omega$

$$= J' \langle 0, \tilde{j}(\beta), 1 \rangle \quad \text{if } \omega < \alpha = \beta + 1$$

$$= \bigcup_{\beta < \alpha} \tilde{j}(\beta) \quad \text{if } \omega < \alpha \in K_{II}.$$

Theorem 18.63. \tilde{j} is a strictly monotonic ordinal function.

Proof. If $\alpha \leq \omega$ then $\tilde{j}(\alpha) = \alpha$. If $\omega < \alpha = \beta + 1$ then $\tilde{j}(\beta) < J' \langle 0, \tilde{j}(\beta), 1 \rangle = \tilde{j}(\alpha)$. If $\omega < \alpha \in K_{II}$ then

$$\tilde{j}(\alpha) = \bigcup_{\beta < \alpha} \tilde{j}(\beta),$$

$$\tilde{j}(\beta) < \tilde{j}(\beta + 1) \leq \tilde{j}(\alpha).$$

Theorem 18.64.

$$1) \quad [\beta < \tilde{j}(\alpha)] \wedge [\gamma = F_a' \beta] \rightarrow \gamma < \alpha.$$

$$2) \quad [\beta < \tilde{j}(\alpha)] \wedge [\gamma \in F_a' \beta] \rightarrow \gamma < \alpha.$$

Proof. 1) (By induction on α .) If $\alpha \leq \omega$ then $\tilde{j}(\alpha) = \alpha$. Therefore

$$\begin{aligned} \beta < \tilde{j}(\alpha) &\rightarrow \beta < \omega \\ &\rightarrow F_a' \beta = \beta < \tilde{j}(\alpha) = \alpha. \end{aligned}$$

If $\omega < \alpha = \eta + 1$ then

$$\tilde{j}(\alpha) = J' \langle 0, \tilde{j}(\eta), 1 \rangle.$$

Therefore if $\beta < \tilde{j}(\alpha) \wedge \beta = J' \langle \beta_1, \beta_2, i \rangle$ then

$$i = 0 \rightarrow F_a' \beta = F_a'' \beta,$$

$$i = 1 \rightarrow F_a' \beta = \{F_a' \beta_1, F_a' \beta_2\},$$

$$i > 1 \rightarrow F_a' \beta \subseteq F_a' \beta_1.$$

If $i = 0$ and $\gamma = F_a^* \beta$ then $F_a^* \tau$ is an ordinal for all $\tau < \beta$. But $F_a^*(\omega + 9) = \{1, \omega\} \notin \text{On}$. Therefore $\beta < \omega + 9$. Since $i = 0$ we have that $\beta \leq \omega$, hence

$$F_a^* \beta = \beta < \alpha.$$

If $\gamma = \{F_a^* \beta_1, F_a^* \beta_2\}$ then $\gamma \leq 2 < \alpha$. If $i > 1 \wedge \gamma = F_a^* \beta$ then, since by Theorem 15.20,

$$F_a^* \beta \subseteq F_a^{**} \beta,$$

it follows that $\tau < \gamma = F_a^* \beta \subseteq F_a^* \beta_1 \rightarrow (\exists \beta_3 < \beta_1) [\tau = F_a^* \beta_3]$. Furthermore

$$i > 1 \wedge \beta < \tilde{j}(\alpha) \rightarrow \beta_1 < \tilde{j}(\eta).$$

From the induction hypothesis

$$\beta_1 < \tilde{j}(\eta) \wedge \tau = F_a^* \beta_3 \rightarrow \tau < \eta$$

i.e. $\tau < \gamma \rightarrow \tau < \eta$. Thus

$$\gamma = F_a^* \beta \rightarrow \gamma \leq \eta < \alpha.$$

If $\omega < \alpha \in K_{II}$ then

$$\tilde{j}(\alpha) = \bigcup_{\eta < \alpha} \tilde{j}(\eta).$$

Therefore

$$\beta < \tilde{j}(\alpha) \rightarrow (\exists \eta < \alpha) [\beta < \tilde{j}(\eta)].$$

For the induction hypothesis

$$\gamma = F_a^* \beta \rightarrow \gamma < \eta < \alpha.$$

2) Since by Theorem 15.20

$$F_a^* \beta \subseteq F_a^{**} \beta,$$

$$\gamma \in F_a^* \beta \rightarrow (\exists \eta < \beta) [\gamma = F_a^* \eta].$$

Then $[\eta < \beta] \wedge [\beta < \tilde{j}(\alpha)]$ implies $\eta < \tilde{j}(\alpha)$. But from

$$[\eta < \tilde{j}(\alpha)] \wedge [\gamma = F_a^* \eta].$$

and 1) above, we conclude that

$$\gamma < \alpha.$$

Theorem 18.65.

$$[\omega < \alpha \in K_{II}] \wedge [\beta < \tilde{j}(\alpha)] \wedge [\gamma < \tilde{j}(\alpha)] \rightarrow [J^* \langle \beta, \gamma, 1 \rangle < \tilde{j}(\alpha)].$$

Proof.

$$[\beta < \tilde{j}(\alpha)] \wedge [\gamma < \tilde{j}(\alpha)] \rightarrow (\exists \eta < \alpha) [[\beta < \tilde{j}(\eta)] \wedge [\gamma < \tilde{j}(\eta)]]$$

$$J^* \langle \beta, \gamma, 1 \rangle < J^* \langle 0, \tilde{j}(\eta), 1 \rangle = \tilde{j}(\eta + 1) < \tilde{j}(\alpha).$$

Theorem 18.66.

$$[\omega < \alpha \in K_{II}] \wedge [\beta < \tilde{j}(\alpha)] \rightarrow [g_3(\beta) < \tilde{j}(\alpha)] \wedge [g_4(\beta) < \tilde{j}(\alpha)].$$

Proof. Since g_3 and g_4 are J -functions this follows from the previous theorem by induction on type.

$$\textbf{Theorem 18.67. } [\omega < \alpha \in K_{II}] \wedge [\beta < \alpha] \rightarrow F'_0 \beta < \tilde{j}(\alpha).$$

Proof (by induction on β). If $\beta < \omega$ then

$$F'_0 \beta = \beta < \alpha = \tilde{j}(\alpha).$$

If $\omega < \beta = \eta + 1$ then $\eta < \beta$ and hence by the induction hypothesis

$$F'_0 \eta < \tilde{j}(\alpha).$$

But from the preceding theorem

$$F'_0 \beta = g_3(F'_0 \eta) < \tilde{j}(\alpha).$$

If $\omega < \beta \in K_{II}$ then

$$F'_0 \beta = g_4(\cup F''_0 \beta).$$

But from the induction hypothesis and the fact that $\beta \in K_{II}$

$$F''_0 \beta = \bigcup_{\gamma < \beta} F'_0 \gamma \leq \bigcup_{\gamma < \beta} \tilde{j}(\gamma) = \tilde{j}(\beta).$$

Since $\beta < \alpha$

$$\tilde{j}(\beta) < \tilde{j}(\alpha)$$

and hence by Theorem 18.66

$$F'_0 \beta = g_4(\cup F''_0 \beta) < \tilde{j}(\alpha).$$

Theorem 18.68.

$$[\omega < \alpha \in K_{II}] \wedge [\beta < \alpha] \wedge [\gamma \in F'_a F'_0 \beta \vee \gamma = F'_a F'_0 \beta] \rightarrow \gamma < \alpha.$$

Proof. From Theorem 18.67

$$F'_0 \beta < \tilde{j}(\alpha).$$

Hence by Theorem 18.64

$$\gamma \in F'_a F'_0 \beta \rightarrow \gamma < \alpha,$$

$$\gamma = F'_a F'_0 \beta \rightarrow \gamma < \alpha.$$

$$\textbf{Theorem 18.69. } \alpha \in K_{II} \rightarrow \cup (F''_0 \alpha) \in K_{II}.$$

Proof. $\alpha \in K_{II} \rightarrow \alpha \neq 0 \wedge \cup (F''_0 \alpha) \neq 0$. If $\cup (F''_0 \alpha) = \beta + 1$ then $\beta \in \cup (F''_0 \alpha)$. Hence

$$(\exists \gamma < \alpha) [\beta \in F'_0 \gamma].$$

Since F_0 is strictly monotonic

$$\beta \in F_0^* \gamma < F_0^*(\gamma + 1) \in F_0^{**} \alpha$$

i.e.

$$\begin{aligned} F_0^* \gamma \in \cup (F_0^{**} \alpha) &= \beta + 1, \\ \beta < F_0^* \gamma &< \beta + 1. \end{aligned}$$

From this contradiction we conclude that $\cup (F_0^{**} \alpha) \in K_{II}$.

Theorem 18.70. $F_a^* F_0^* \alpha = \alpha$.

Proof (by induction on α). If $\alpha \leq \omega$ then $F_0^* \alpha = \alpha$ and $F_a^* \alpha = \alpha$ hence

$$F_a^* F_0^* \alpha = \alpha.$$

If $\omega < \alpha = \beta + 1$ then from the induction hypothesis and Theorem 18.54

$$F_a^* F_0^* \alpha = F_a^* g_3(F_0^* \beta) = F_a^* F_0^* \beta \cup \{F_a^* F_0^* \beta\} = \beta \cup \{\beta\} = \alpha.$$

If $\omega < \alpha \in K_{II}$ then

$$F_a^* F_0^* \alpha = F_a^* g_4(\cup F_0^{**} \alpha) = F_a^*(\cup F_0^{**} \alpha) \cap \text{On}.$$

Since $\alpha \in K_{II}$, $\cup F_0^{**} \alpha \in K_{II}$. Hence

$$F_a^*(\cup F_0^{**} \alpha) = F_a^{**}(\cup F_0^{**} \alpha).$$

Therefore

$$\begin{aligned} F_a^*(\cup F_0^{**} \alpha) \cap \text{On} &= \{\gamma \mid (\exists \beta < \cup F_0^{**} \alpha) [\gamma = F_a^* \beta]\} \\ &= \{\gamma \mid (\exists \tau < \alpha) [\gamma \in F_a^* F_0^* \tau]\}. \end{aligned}$$

Since by the induction hypothesis $\tau < \alpha \rightarrow F_a^* F_0^* \tau = \tau$ we conclude that

$$F_a^* F_0^* \alpha = \{\gamma \mid \gamma < \alpha\} = \alpha.$$

Theorem 18.71. $(\forall a \subseteq \omega) [\alpha \in \mathcal{M}_0 \rightarrow \alpha \in N_a]$.

Proof. Since F_0 is defined from J -functions $\alpha \in \mathcal{M}_0 \rightarrow F_0^* \alpha \in \mathcal{M}_0$.

$$F_0^* \alpha \in \mathcal{M}_0 \rightarrow F_a^* F_0^* \alpha \in N_a.$$

But $F_a^* F_0^* \alpha = \alpha$.

Remark. From Theorem 18.71 we see that for each $a \subseteq \omega$ the ordinals in N_a are precisely the ordinals in \mathcal{M}_0 . Furthermore with Theorem 18.71 we have eliminated from this section the need to know that \mathcal{M}_0 is a submodel of every standard transitive model of ZF. Consequently all of the results of this section hold if \mathcal{M}_0 is replaced any \mathcal{M} , that is a countable standard transitive model of ZF + AC + GCH + $V = L$.

19 Languages, Structures, and Models

In Sections 2–15 we developed Zermelo-Fraenkel set theory in a very simple language and we proved Gödel's consistency results. In Sections 16–18 we enriched the original language by adding individual constants. In this enriched language we developed a rather powerful technique, Cohen's forcing, with which we proved one of Cohen's independence results. In this section we will introduce a language that is richer than any we have used heretofore. We will briefly outline the development of Zermelo-Fraenkel theory in this language and define certain concepts for later use. One of our objectives is to explain an earlier claim that the theories $ZF\{\alpha\}$ are slightly different formulations of Zermelo-Fraenkel set theory than that of Sections 2–15. Our main objective, however, is the introduction of a language \mathcal{L} that is of great value in the development of a very general method for attacking certain problems in set theory.

The language \mathcal{L} consists of the following:

individual variables:	$x_0, x_1, \dots, x_i, \dots;$	$i < \omega,$
class variables:	$A_0, A_1, \dots, A_i, \dots;$	$i < \omega,$
individual constants:	$k_0, k_1, \dots, k_\alpha, \dots;$	$\alpha < \alpha_0,$
n-ary predicate symbols:	$R_0^n, R_1^n, \dots, R_i^n, \dots;$	$i < \omega, n \geq 1,$
logical symbols:	$\neg, \vee, \wedge, \rightarrow, \leftrightarrow, \exists, \forall,$	
auxiliary symbols:	$[], ().$	

We will use

$$a, b, c, \dots, x, y, z$$

as meta-variables whose range is the collection of individual variables of the language. We will use

$$s, t, u$$

as meta-variables whose range is the collection of terms. By a term we mean an individual variable, a class variable, or an individual constant. We will also use

$$\varphi, \psi, \eta$$

as metavariables that range over the collection of well formed formulas (wffs). This collection we define inductively in the following way:

- 1) If s and t are terms then $s \in t$ is a wff.
- 2) If φ and ψ are wffs then $\neg \varphi$, $\varphi \vee \psi$, $\varphi \wedge \psi$, $\varphi \rightarrow \psi$, $\varphi \longleftrightarrow \psi$ are wffs.
- 3) If φ is a wff and x is an individual variable then $(\forall x) \varphi$ and $(\exists x) \varphi$ are wffs.

From the language \mathcal{L} we obtain a theory of sets, $ZF_{\mathcal{L}}$, by adjoining the logical axioms, rules of inference, and nonlogical axioms of ZF. However in order for the nonlogical axioms to be wffs of the language we must have \in as one of our binary predicate symbols. We assume it to be R_0^2 .

In Section 3, Definition 3.1 and Theorem 3.2 can be extended to terms in the obvious way. However, since Axiom 1 tells us nothing about individual constants or class variables, the proof of Theorem 3.3 remains valid only for sets. This theorem is extended to terms in Section 4 by means of Definition 4.2 from which we have

$$s \in t \longleftrightarrow (\exists x \in t) [x = s].$$

A serious difficulty arises with Theorem 3.4. Since each

$$R_i^n(t_1, \dots, t_n)$$

is an atomic formula we must establish that

$$t_1 = s_1 \wedge \dots \wedge t_n = s_n \rightarrow [R_i^n(t_1, \dots, t_n) \longleftrightarrow R_i^n(s_1, \dots, s_n)]. \quad (1)$$

With the aid of Axiom 1 this is easily done for R_0^2 . We however have no means of establishing (1) for the remaining relational symbols.

We could require that equality be one of our binary predicate symbols and take (1) as an equality axiom. We however choose to abandon Theorem 3.4. Similarly we abandon Theorem 4.8 and 4.12.

Definitions 4.1–4.3 and Theorem 4.4 are easily modified to our new language. We then extend the notion of term to include class symbols or as an alternative we can introduce a formal notion of class: By a class we mean an individual variable or a class variable or an individual constant or a class symbol.

The remaining results of Section 4 are unaltered except that Theorem 4.9 can be extended to individual constants and Definition 4.15 should be broadened:

Definition. A class A is *definable* in \mathcal{L} using the terms t_1, \dots, t_n iff there is a wff $\varphi(x, t_1, \dots, t_n)$ containing no free variables other than those indicated, such that

$$A = \{x \mid \varphi(x, t_1, \dots, t_n)\}.$$

A class A is definable in \mathcal{L} iff there is a wff $\varphi(x)$ containing no free variables other than x and no individual constants, such that

$$A = \{x \mid \varphi(x)\} .$$

All of the material in Section 5–11 can be retained essentially as it is. In Section 12 we must alter the notion of structure and model to take into account the individual constants and the new relational symbols.

By a structure for \mathcal{L} , denoted by $[A, g]$, we mean a nonempty class A called the universe of the structure, together with a function g that assigns to each individual constant an element of A or a subclass of A i.e.

$$(\forall \alpha < \alpha_0) [[g(k_\alpha) \in A] \vee [g(k_\alpha) \subseteq A]],$$

and assigns to each n -ary relational symbol an n -ary relation on A i.e.

$$(\forall i < \omega) (\forall n \geq 1) [g(R_i^n) \subseteq A^n] .$$

The structure is a standard structure iff

$$g(\epsilon) = E \cap A^2 .$$

For convenience in defining satisfaction we introduce the following notations:

$$\bar{x}_i \triangleq x_i ,$$

$$\bar{k}_\alpha \triangleq g(k_\alpha) ,$$

$$\bar{R}_i^n \triangleq g(R_i^n) ,$$

$$[A; \bar{k}_0, \bar{k}_1, \dots, \bar{R}_0^n, \bar{R}_1^n, \dots] \triangleq [A, g] .$$

Definition. If $\mathcal{A} = [A; \bar{k}_0, \bar{k}_1, \dots; \bar{R}_0^n, \bar{R}_1^n, \dots]$ is a structure for \mathcal{L} then

- 1) $\mathcal{A} \models R_i^n(t_1, \dots, t_n) \longleftrightarrow \bar{t}_1 \in A \wedge \dots \wedge \bar{t}_n \in A \wedge (\bar{t}_1, \dots, \bar{t}_n) \in \bar{R}_i^n .$
- 2) $\mathcal{A} \models \neg \psi \longleftrightarrow \neg [\mathcal{A} \models \psi] .$
- 3) $\mathcal{A} \models \psi \wedge \eta \longleftrightarrow \mathcal{A} \models \psi \wedge \mathcal{A} \models \eta .$
- 4) $\mathcal{A} \models (\forall x) \psi(x) \longleftrightarrow (\forall x \in A) [\mathcal{A} \models \psi(x)] .$

By a model of our theory we mean a structure \mathcal{A} that satisfies each of the axioms. We will refer to any such model as a model of Zermelo-Fraenkel set theory. In contexts where the language is of interest we refer to a model of ZF in the language \mathcal{L} .

The remaining details of the development we leave to the reader.

Bibliography

- Bar-Hillel, Y. (Ed.): *Essays on the foundations of mathematics*. Jerusalem: Magnes Press 1966.
- Benacerraf, P., Putnam, H.: *Philosophy of mathematics selected readings*. Englewood Cliffs: Prentice-Hall Inc. 1964.
- Bernays, P., Fraenkel, A. A.: *Axiomatic set theory*. Amsterdam: North-Holland 1958.
- Cantor, G.: *Beiträge zur Begründung der transfiniten Mengenlehre (Erster Artikel)* *Math. Ann.* **46**, 481—512 (1895); — *(Zweiter Artikel)* *Math. Ann.* **49**, 207—246 (1897).
- *Contributions to the founding of the theory of transfinite numbers*, English translation by Philip E. B. Jourdain. Reprinted by Dover Publications Inc. (1915).
- Church, A.: *Introduction to mathematical logic*. Princeton: Princeton University Press 1956.
- Cohen, P.: *The independency of the continuum hypothesis*. *Proc. Nat. Acad. Sci. U.S.* **50**, 1143—1148 (1963).
- *Set theory and the continuum hypothesis*. Amsterdam: W. A. Benjamin Inc. 1966.
- Fraenkel, A. A., Bar-Hillel, Y.: *Foundations of set theory*. Amsterdam: North-Holland 1958.
- Gödel, K.: *The consistency of the continuum hypothesis*. Princeton: Princeton University Press 1940.
- *What is Cantor's continuum problem?* *Am. Math. Monthly* **54**, 515—525 (1947).
- van Heijenoort, J.: *From Frege to Gödel*. Cambridge: Harvard University Press 1967.
- Quine, W. V.: *Set theory and its logic*. Cambridge: Belknap Press 1969.
- Rubin, J. E.: *Set theory for the mathematician*. San Francisco: Alden-Day 1967.
- Schoenfield, J. R.: *Mathematical logic*. Reading: Addison Wesley 1967.

Problem List

1. Let A be an infinite set. Prove that the cardinality of the set of all automorphisms of A , i.e. one-to-one mappings of A onto A , is $\overline{2^A}$. (Hint: Divide A into A_1, A_2, A_3 so that $\overline{A_1} = \overline{A_2} = \overline{A_3} = \overline{A}$. For each $B \subseteq A_2$ find an automorphism π for which $\pi(A_1 \cup B) = A_3 \cup (A_2 - B)$.

2. Let A be a countable infinite set and $<_1$ be an order relation on A (Definition 6.18). Let R_0 be the set of rationals in the interval $(0, 1)$. Find a one-to-one order-preserving map τ from A into R_0 . (Hint: Let $A = \{a_0, a_1, \dots\}$. Define $\tau(a_i)$ assuming that $\tau(a_0), \dots, \tau(a_{i-1})$ have been defined.

3. Let A_1 and A_2 be infinite countable sets. Let $<_1$ Or $A_1, <_2$ Or A_2 , and both structures satisfy

- a) $(\forall x)(\exists y)[y < x]$
- b) $(\forall x)(\exists y)[x < y]$
- c) $(\forall x)(\forall y)(\exists z)[x < y \rightarrow x < z < y]$.

Prove: $(\exists f)f \text{ Isom}_{<_1, <_2}(A_1, A_2)$. (Hint: Let $A_1 = \{a_0, a_1, \dots\}$ and $A_2 = \{b_0, b_1, \dots\}$. Define $\tau_1 \text{ Isom}_{<_1, <_2}(A_1, A_2)$ and $\tau_2 \text{ Isom}_{<_2, <_1}(A_2, A_1)$ inductively in the order $\tau_1(a_0), \tau_2(b_0), \tau_1(a_1), \tau_2(b_1), \dots$, such that $\tau_1 \circ \tau_2$ and $\tau_2 \circ \tau_1$ are identity functions on A_2 and A_1 respectively.

4. Let $W_\alpha = \{\langle \alpha_0, \dots, \alpha_n \rangle \mid n < \omega \wedge (\forall i \leq n) [\alpha_i < \alpha]\}$. Let $<_\alpha$ be the lexicographical ordering on W_α . Prove that if α is finite $\langle W_\alpha, <_\alpha \rangle$ is isomorphic to $R_1 \times \omega$ where R_1 is the set of all rationals in the interval $[0, 1)$ and $R_1 \times \omega$ is ordered lexicographically relative to the natural order on R_1 and on ω . What is the order type of $\langle W_\alpha, <_\alpha \rangle$ if $\alpha \geq \omega$?

5. Let \mathcal{R} be the set of all real numbers and let f be a mapping from \aleph_1 into \mathcal{R} that is monotone increasing. Prove: $(\exists \alpha < \aleph_1) (\forall \beta > \alpha) [f(\beta) = f(\alpha)]$. (Hint: \mathcal{R} is separable and hence $\text{cf}(f'' \aleph_1) = \aleph_0$.

6. Let \mathcal{R} be the set of all real numbers and let f be a continuous mapping from \aleph_1 into \mathcal{R} . Prove: $(\exists \alpha < \aleph_1) (\forall \beta > \alpha) [f(\beta) = f(\alpha)]$.

7. Let \mathcal{R} be the set of all real numbers and let $f: \aleph_1 \xrightarrow{1-1} \mathcal{R}$. $\forall \alpha, \beta < \aleph_1$ define $\alpha \ll \beta$ iff $\alpha < \beta \wedge f(\alpha) < f(\beta)$. Prove the following:

- a) \ll is a well-founded partial ordering on \aleph_1 .
- b) If $A \subseteq \aleph_1$, and $(\forall x, y \in A) [x \ll y \vee x = y \vee y \ll x]$ then A is countable.
- c) If $A \subseteq \aleph_1$ and $(\forall x, y \in A) [x = y \vee \neg[x \ll y \vee y \ll x]]$ then A is countable.

8. Let $A \subseteq [0, 1]$. $\forall x \in [0, 1]$, x is a κ -accumulation point of A iff

$$\forall N(x) [\overline{\overline{N(x) \cap A}} \geq \kappa].$$

(Here $N(x)$ is a neighborhood of x in the usual topology on $[0, 1]$.)

a) Prove that $\{x \in [0, 1] \mid x \text{ is an } \aleph_1\text{-accumulation point of } A\}$ is a closed set that is dense in itself.

b) Prove that $\{x \in [0, 1] \mid x \text{ is an } \aleph_2\text{-accumulation point of } A\}$ is a closed set that is dense in itself.

9. If $\text{cf}(\aleph_\alpha) < \aleph_\alpha$ and $(\forall \lambda < \alpha) (\exists \nu < \alpha) [2^{\aleph_\lambda} < 2^{\aleph_\nu}]$ and if $\lambda = \sup_{\xi < \aleph_\alpha} \overline{\overline{\aleph_\xi^{\text{cf}(\aleph_\xi)}}$ prove that $\overline{\overline{2^{\aleph_\alpha}}} = \overline{\overline{\lambda^{\text{cf}(\lambda)}}$.

10. If $\text{cf}(\aleph_\alpha) < \aleph_\alpha$, if $\lambda < \alpha$, and if $(\forall \nu < \alpha) [\lambda \leq \nu \rightarrow \overline{\overline{2^{\aleph_\lambda}}} = \overline{\overline{2^{\aleph_\nu}}}]$ prove that $\overline{\overline{2^{\aleph_\alpha}}} = \overline{\overline{2^{\aleph_\lambda}}}$.

11. Prove that if $\aleph_\alpha > \aleph_\beta \geq \text{cf}(\aleph_\alpha)$ and $(\exists \gamma < \alpha) [\aleph_\alpha \leq \overline{\overline{\aleph_\gamma^{\aleph_\beta}}}]$ then

$$\overline{\overline{\aleph_\alpha^{\aleph_\beta}}} = \overline{\overline{\aleph_{\gamma_0}^{\aleph_\beta}}}$$

where $\gamma_0 = \mu_\gamma(\gamma < \alpha \wedge \aleph_\alpha \leq \overline{\overline{\aleph_\gamma^{\aleph_\beta}}})$.

12. Let $\aleph_\alpha > \aleph_\beta \geq \text{cf}(\aleph_\alpha)$ and $(\forall \gamma < \alpha) [\overline{\overline{\aleph_\gamma^{\aleph_\beta}}} < \aleph_\alpha]$. Prove that

$$\overline{\overline{\aleph_\alpha^{\text{cf}(\aleph_\alpha)}}} = \overline{\overline{\aleph_\alpha^{\aleph_\beta}}}.$$

13. Prove: If $\overline{\overline{2^{\text{cf}(\aleph_\alpha)}}} < \aleph_\alpha$, if $(\exists \beta < \alpha) [\aleph_\alpha \leq \overline{\overline{\aleph_\beta^{\text{cf}(\aleph_\beta)}}} \wedge \text{cf}(\aleph_\beta) < \text{cf}(\aleph_\alpha)]$ and if $\lambda = \mu_\gamma(\text{cf}(\aleph_\gamma) \leq \text{cf}(\aleph_\alpha) \wedge \aleph_\alpha \leq \overline{\overline{\aleph_\gamma^{\text{cf}(\aleph_\gamma)}}})$, then $\overline{\overline{\aleph_\alpha^{\text{cf}(\aleph_\alpha)}}} = \overline{\overline{\aleph_\lambda^{\text{cf}(\aleph_\lambda)}}}$. (Hint: If $\nu = \mu_\gamma(\aleph_\alpha \leq \overline{\overline{\aleph_\gamma^{\text{cf}(\aleph_\alpha)}}})$ then $\overline{\overline{\aleph_\alpha^{\text{cf}(\aleph_\alpha)}}} = \overline{\overline{\aleph_\nu^{\text{cf}(\aleph_\nu)}}}$ and $\lambda = \nu$.)

14. Let F^{α_0} be a model of ZF. Prove that $\{\ulcorner \varphi \urcorner \mid F^{\alpha_0} \models \varphi\} \in L$.

15. A set a is L -finite iff $(\forall x \in L) [x \subseteq a \rightarrow x \text{ is finite}]$. Assuming that $\overline{\overline{\mathcal{P}(\omega)^L}} = \omega$, prove that $(\exists x \subseteq \omega)$, x and $\omega - x$ are each L -finite.

16. Find a model N_a of $\text{ZF} + \text{AC} + V \neq L$ such that the forcing set a is L -finite. (Hint: Make a suitable modification of the definition of forcing set.)

17. If $\vdash^1 \ulcorner \varphi \urcorner \leftrightarrow (\exists \mathcal{P}) [\mathcal{P} \Vdash \ulcorner \varphi \urcorner]$ and $\vdash^2 \ulcorner \varphi \urcorner \leftrightarrow (\forall \mathcal{P}) [\mathcal{P} \Vdash \ulcorner \varphi \urcorner]$ what logical rules do \vdash^1 and \vdash^2 satisfy.

18. A sentence is called arithmetical if every quantifier in it is restricted to ω . Let φ be an arithmetic sentence and let F^{α_0} be a model of ZF. Prove

$$\varphi \leftrightarrow F^{\alpha_0} \models \varphi$$

19. A sentence is called a $\mathcal{P}(\omega)$ -sentence if every quantifier in it is restricted to $\mathcal{P}(\omega)$. Assuming the existence of the minimal model \mathcal{M}_0 find a $\mathcal{P}(\omega)$ -sentence φ such that

$$L \models \varphi \leftrightarrow \mathcal{M}_0 \models \varphi$$

is false.

20. In a forcing model M of $V \neq L$, find $a \subseteq \omega$ such that

a) $L_a \neq M$.

b) a and $\omega - a$ are L -finite.

Appendix

Let M , SM , and $\text{Consis}(\text{ZF})$ be statements that assert respectively.

1. There exists a set that is a model of ZF.
2. There exists a set that is a standard model of ZF.
3. ZF is consistent.

Furthermore, let $\text{Consis}(\text{ZF})$ be so chosen that it is absolute w.r.t. every standard transitive model of ZF.

Theorem. $\neg \vdash_{\text{ZF}} M \rightarrow SM$.

Proof. Suppose

1. $\vdash_{\text{ZF}} M \rightarrow SM$.

It is known that

2. $\vdash_{\text{ZF}} M \leftrightarrow \text{Consis}(\text{ZF})$.

Consequently, from 1. and 2.

3. $\text{Consis}(\text{ZF}) \vdash_{\text{ZF}} SM$.

There exists a minimal standard transitive model of ZF, \mathcal{M}_0 . Clearly

4. $\mathcal{M}_0 \models \text{Consis}(\text{ZF})$.

Then relativizing 3. to \mathcal{M}_0 , using the fact that $\text{Consis}(\text{ZF})$ is absolute w.r.t. \mathcal{M}_0 , we have

$$\mathcal{M}_0 \models SM.$$

This is a contradiction.

Index

- absoluteness 112—123
- addition, ordinal 49
- Aleph Hypothesis (AH) 93, 96
- almost universal 135
- arithmetic, ordinal 49—62
- auxiliary symbols 4, 236
- Axiom of Abstraction 9, 16
 - of Choice (AC) 2, 87, 87—101
 - of Constructibility 152—174
 - of Extensionality 7, 123
 - of Foundation 18
 - of Infinity 39, 123, 129
 - of Pairing 14, 123—126
 - of Powers 15, 123, 127—128
 - of Regularity 18, 19, 130
 - of Unions 15, 123, 126—127
 - Schema of Replacement 17, 128
 - Schema of Separation 16—17
- axioms, logical 5

- Benacerraf, Paul** 2
- Bernays, Paul (1888) 3
- Bernstein, Felix 71
- Burali-Forti paradox 37

- Cantor, Georg (1845—1918) 1—3, 32, 37, 71—73, 88, 90, 92
- Cantor's Law of Trichotomy 88
 - notion of cardinal number 73
 - notion of ordinal number 32
 - notion of set 2
- Cantor-Schröder-Bernstein Theorem 71, 90
- cardinal, inaccessible 85—86
 - numbers 1, 71—101
 - —, Cantor's notion of 73
 - —, class of 79
 - —, cofinality of 82
 - , regular 85
 - , singular 85
 - , weakly inaccessible 85—86
- character of cofinality 84
- class, almost universal 135
 - , definable 237
 - , M-constructible 140
 - of cardinal numbers 79
 - of closed wffs 191
 - of closed limited formulas 191
 - of limit ordinals 38
 - of prime formulas 178
 - , proper 3, 12
 - of quasi formulas 179
 - of statements 203
 - symbol 9
 - of wffs 189
- class(es) 3, 9—20
 - , ordinal 32
- closed limited formulas 190—191
 - unlimited statements 203
 - wffs 190—191
- cofinality, character of 84
 - of ordinals 82
- Cohen, Paul 2, 3, 73, 96
- complete sequence of forcing conditions 210
- conjunction 4
- constant, predicate 4
- continuum Hypothesis (CH) 2, 92

- Dedekind, Richard (1831—1916)** 1, 2
- definable class 237
 - , set 13
- disjunction 4

- end of string 184
- equality 6—8
- equivalence 4
 - , of sets 71
- equivalent, logically 5
- existential quantification 4
- extensionality 7—8

- finite induction 39
 - recursion 43
 - set 75
- forcing conditions 204
 - —, complete sequence of 210
 - for unlimited statements 205
 - , for limited statements 205—206
 - set 211
- formula, model of 103
 - , parity of 204
 - , quasi 179
 - , quasi limited 179
 - , subquasi 187
 - , type of 203
 - , unlimited 203
 - , well formed 4, 175, 189, 237
- formulas, logically equivalent 5
 - , prime 5, 178
 - , well formed 4
 - , well formed parts of 5
- foundational relation(s) 27
- Fraenkel, Abraham A. (1891) 3, 16, 17
- Frege, Gottlob (1848—1925) 1, 2, 9, 32, 73
- function(s) 21—31
- function, ordinal 44
 - , satisfaction 192
- fundamental function 214
 - operations 133
- Generalized Continuum Hypothesis (GCH)** 2, 92—100
- Gödel-Bernays set theory (GB)** 3
 - , Kurt (1906) 2, 3
 - model 148—174
 - numbers 176
- implication 4
- inaccessible cardinal 85—86
- individual variables 4
- induction, finite 39
 - , transfinite 35
- inference, rules of 5
- infinite set 75
- J-function** 228
- König** 100
- language(s) 4—5, 236—238
- limit ordinal 37
 - ordinals, class of 38
- limited quantifier 178
 - statements, forcing for 205—20
- Lindenbaum, A. 93
- logic 4—5
- logical axioms 5
 - symbols 4
- logically equivalent 5
- \mathcal{M} -constructible 140
- \mathcal{M}_0 -definable 201
- membership 3
- minimal model 197
- model of a formula 103
 - , minimal 197
 - of ZF 102, 130—142
 - , standard transitive 107
- models 102—111
- Montague, Richard 17
- Mostowski, A. 107
- multiplication, ordinal 53
- negation 4
- n-tuple, ordered 14
 - , unordered 14
- number(s), cardinal 1, 71—86
 - ordinal 1
- order of a set 149
 - , partial 26
 - relation 26
- ordered n-tuples 14
 - pairs 14
- ordinal addition 49
 - arithmetic 49—62
 - classes 32
 - function 44
 - , limit 37
 - multiplication 53
 - numbers 1, 32—62
 - —, Cantor's notion of 32
- ordinals, cofinality of 82
- pair, ordered 14
 - , unordered 14
- Paradox, Berali-Forti 37
 - , Russell's 2, 9, 10
- parity of a formula 204
- partial ordering 26
- Peano postulates 38
- Poincare, Henri (1854—1912) 1
- power set 15
- predicate constant 4
 - symbol 3

- prime formulas 5, 178
- proper class (s) 3, 12
- Putnam, Hilary 2
- quantification, existential** 4
 - , limited 178
 - , universal 4
- quasi formulas 179
 - —, string of 182
 - limited formulas 179
- Quine, Van Orman 8
- rank 66, 68
- recursion, finite 43
 - , transfinite 42
- regular cardinal 85
- relational systems 26—31, 102
- relations 21—31
 - , foundational 27
 - , order 26
 - , well founded 27
 - , well ordering 27
- R-minimal element 27
- Rubin, H. 96
- rules for well formed formulas 4
 - of inference 5
- Russell, Bertrand (1872—1970) 1—3, 9, 32, 73
- Russell class 13
- Russell's Paradox 2, 9, 10
- satisfaction functions** 192
- Schröder, Ernst 71
- Scott, Dana 8
- set, definable 13
 - finite 75
 - , forcing 211
 - infinite 75
 - , order of 149
 - , power 15
 - , supertransitive 66
 - theory, Gödel-Bernays (GB) 3
 - —, Zermelo-Fraenkel (ZF) 3, 5, 9
 - , transitive 32
 - union 15
 - , well founded 67
- set(s) 3, 12
 - , Cantor's notion of 2
- sets, equivalent 71
- Sierpinski, W. 93—94
- singular cardinal 85
- Skolem, Thoralf (1887—1963) 3
- Standard Model Hypothesis 197
- standard structure 103
 - transitive model 107
- statements 203
 - class of 203
- string, end of 184
 - of quasi formulas 182
- structure 102
 - , standard 103
- subclass 15
- subquasi formula 187
- supertransitive set 66
- symbols, auxiliary 4, 236
- symbol, class 9
- symbols, logical 4
- symbol(s), logical 4
 - , predicate 4
- Tarski, A.** 93
- term 12, 236
- theory of types 3
- transfinite induction 35
 - recursion 42
- transitive set 32
- type of a formula 203
- types, theory of 3
- union, of a set** 15
- universal quantification 4
- unlimited statements 203
- unlimited statements, forcing for 205
- weakly inaccessible cardinals 85—86
- unlimited formulas 203
- unordered n-tuple 14
 - pair(s) 14
- van Heijenoort, Jean 2
- variables, individual 4
- von Neumann, John (1903—1957) 3, 32
- well formed formulas** 4, 175, 189, 237
 - — —, rules for 4
 - — — in the wider sense 10
 - founded relation(s) 27
 - — set 67
- Well Ordering Principle 88
- well ordering relations 27
- Whitehead, Alfred North (1861—1947) 3
- Zermelo, Ernst (1871—1953) 1, 3, 16—18, 98
- Zermelo-Fraenkel set theory (ZF) 3, 5, 9
- Zorn's Lemma 88

Index of Symbols

- GB 3
 ZF 3
 \in 3
 \neg 4
 $\rightarrow, \leftrightarrow$ 4
 \forall, \exists 4
 \vee, \wedge 4
 wff(s) 4
 x_0, x_1, \dots 4
 a, b, c, \dots, x, y, z 4
 φ, ψ, η 4
 \vdash 5
 \vdash_{LA} 5
 $a = b$ 6
 $\{x \mid \varphi(x)\}$ 10
 φ^* 11
 A, B, C 12
 $A = B$ 12
 $\mathcal{M}(A)$ 13
 $\mathcal{P}_r(A)$ 13
 Ru 13
 $\{a_1, \dots, a_n\}$ 14
 $\langle a_1, \dots, a_n \rangle$ 14
 $\cup(A)$ 15
 $\mathcal{P}(a)$ 15
 $A \subseteq B$ 15
 $A \subset B$ 15
 $A \cup B$ 15
 $A \cap B$ 15
 $A - B$ 18
 0 18
 $\{x \in a \mid \varphi(x)\}$ 18
 V 19
 $A \times B$ 21
 A^2 21
 A^{-1} 21
 $\mathcal{R}el(A)$ 21
 $\mathcal{U}n(A)$ 21
 $\mathcal{U}n_2(A)$ 21
 $\mathcal{F}nc(A)$ 21
 $\mathcal{F}nc_2(A)$ 21
 $\mathcal{D}(A)$ 21
 $\mathcal{W}(A)$ 21
 $A \vdash B$ 22
 $A \dashv B$ 22
 $A \circ B$ 22
 $\exists!$ 24
 $A \cdot b$ 24
 $A \mathcal{F}n B$ 25
 $A \mathcal{F}n_2 B$ 25
 $F : A \rightarrow B$ 25
 $F : A \xrightarrow{\text{onto}} B$ 25
 $F : A \xrightarrow{1-1} B$ 25
 $F : A \xrightarrow[1-1]{\text{onto}} B$ 25
 $R \text{Or } A$ 26
 $R \text{Po } A$ 26
 $R^{-1} A$ 26
 $R \text{Fr } A$ 27
 E 27
 $R \text{Wfr } A$ 27
 $R \text{We } A$ 27
 $R \text{Wfwe } A$ 27
 $H \text{Isom}_{R_1, R_2}(A_1, A_2)$ 29
 I 29
 $\text{Tr}(A)$ 32
 $\text{Ord}(A)$ 32
 α, β, γ 35
 $(\forall \alpha) \varphi(\alpha)$ 35
 $(\exists \alpha) \varphi(\alpha)$ 35
 $\alpha < \beta$ 35
 $\alpha \leq \beta$ 35
 $\max(\alpha, \beta)$ 35
 $\alpha + 1$ 37
 K_I, K_{II} 37
 ω 37
 i, j, k, \dots, m, n 38
 $(\forall i) \varphi(i)$ 38
 $(\exists i) \varphi(i)$ 38
 $\cap(A)$ 40
 $\sup(A), \inf(A)$ 40
 $\sup(A), \inf(A)$ 40
 $\begin{matrix} < \beta \\ > \beta \end{matrix}$
 $\mu_\alpha(\varphi(\alpha))$ 40
 $\text{Orf}(G)$ 44
 $\text{Smo}(G)$ 44
 Le 48
 R_0 48
 J_0 48
 $\alpha + \beta$ 49
 $\alpha - \beta$ 52
 α^β 53
 α^b 57
 $\text{TrCl}(a)$ 63
 $\text{Cl}(R, A)$ 65
 $\text{Cl}_2(R, A)$ 65
 $\text{St}(A)$ 66
 $R(\alpha)$ 66
 $\text{Wf}(a)$ 67
 $\text{rank}(a)$ 68
 $a \simeq b$ 71
 \bar{a} 73
 $\text{Fin}(a), \text{Inf}(a)$ 75
 N 79
 N' 81
 \aleph, \aleph_α 81
 a^b 81
 $\text{cof}(\alpha, \beta)$ 82
 $\text{cf}(\alpha)$ 84
 $\text{Reg}(\alpha)$ 85
 $\text{Sing}(\alpha)$ 85
 $\text{Inacc}_w(\aleph_\alpha)$ 86
 $\text{Inacc}(\aleph_\alpha)$ 86
 AC 87
 WOP 88
 $\text{ch}(B, A)$ 88
 $\max_A(a)$ 88
 GCH, CH 92
 AH 96
 $\prod_{x \in a} c \cdot x$ 98
 a^+ 99
 $[A, R] \models \varphi$ 102
 $A \models \varphi$ 103
 φ^A 103

$STM(A, \varphi)$	107	ic, vc, pf, qf, qlf	202
$\varphi Abs A$	112	$wff, wflf, cf, clf$	202
x^A	117	$typ(\alpha_0; ' \varphi')$	203
$\{x \mid \varphi(x)\}^A$	117	$par(\alpha_0; ' \varphi')$	204
$BAbs A$	118	$\alpha < \beta$	204
$Cnv_2(A), Cnv_3(A)$	133	$cond$	204
$\mathcal{F}_1, \dots, \mathcal{F}_8$	133–134	$\mathcal{P} \subseteq \mathcal{P}'$	204
Q_4, \dots, Q_8	140	$\mathcal{P} \Vdash ' \varphi'$	205
$J \langle \alpha, \beta, m \rangle$	144	$shs(' \varphi')$	207
S	145	$forc(g, ' \varphi')$	208
K_1, K_2, K_3	145	$Forc(\mathcal{P}, ' \varphi')$	208
F	147	$\Phi_\varphi(\mathcal{P}, \alpha, \beta)$	220
L	148	$\Phi_\psi(\mathcal{P}, \alpha, \beta, \gamma)$	220
F_a, L_a	149	$\Gamma_\varphi(\alpha)$	221
Od, Od_a	149	$\Gamma_\psi(\alpha, \beta)$	221
$V = L$	152	$C_\varphi(\alpha)$	221
C, C_a	156	$C_\psi(\alpha, \beta)$	221
$ZF\{\alpha_0\}$	175	$A_1('k_\beta')$	226
$' \varphi'$	176	$A_2('k_\beta')$	226
va	177	$B('k_\beta')$	226
$ic(\alpha_0), vc(\alpha_0)$	177	$g_1(\alpha, \beta, \gamma)$	229
$pf(\alpha_0)$	178	$g_2(\alpha, \beta)$	229
$(\exists^\beta x) \varphi(x)$	178	$g_3(\alpha)$	229
R_3, R_4, R_5, R_6	179	F_0	231
		$\tilde{j}(\alpha)$	232
$qf(\alpha_0), qlf(\alpha_0)$	179		
$Sqf(\alpha_0; f), Sqlf(\alpha_0; f)$	182		
$end(f)$	184		
$Subqf(\alpha_0; ' \varphi', ' \psi')$	187		
$Subqlf(\alpha_0; ' \varphi', ' \psi')$	187		
$Ob(\alpha_0; 'x_i', ' \psi')$	188		
$Ob_i(\alpha_0; 'x_i', ' \varphi')$	188		
$O(\alpha_0; 'x_i', ' \varphi')$	188		
$wff(\alpha_0), wflf(\alpha_0)$	189		
$Sub(\alpha_0; ' \varphi')$	190		
$Sub_i(\alpha_0; ' \varphi')$	190		
$cwf(\alpha_0), clf(\alpha_0)$	191		
$S_j^i ' \varphi'$	191		
$Sm(\alpha_0; n)$	192		
$Sat(g, b, n)$	192		
$b \models ' \varphi'$	193		
$wff(1, 2; 0, 3)$	195		
ax_5	195		
$SM(b, ZF), STM(b, ZF)$	195		
\mathcal{M}_0	196		
N_a	198		
$cov(\alpha)$	198		

Graduate Texts in Mathematics

For information

A student approaching mathematical research is often discouraged by the sheer volume of the literature and the long history of the subject, even when the actual problems are readily understandable. The new series, Graduate Texts in Mathematics, is intended to bridge the gap between passive study and creative understanding; it offers introductions on a suitably advanced level to areas of current research. These introductions are neither complete surveys, nor brief accounts of the latest results only. They are textbooks carefully designed as teaching aids; the purpose of the authors is, in every case, to highlight the characteristic features of the theory.

Graduate Texts in Mathematics can serve as the basis for advanced courses. They can be either the main or subsidiary sources for seminars, and they can be used for private study. Their guiding principle is to convince the student that mathematics is a living science.

Vol. 1 TAKEUTI/ZARING: Introduction to Axiomatic Set Theory. VII, 250 pages. Soft cover US \$ 9.50, DM 35,—; hard cover US \$ 13.50

Vol. 2 OXToby: Measure and Category. VIII, 95 pages. Soft cover US \$ 7.50, DM 28,—; hard cover US \$ 10.50

Vol. 3 SCHAEFER: Topological Vector Spaces. XI, 294 pages. Soft cover US \$ 9.50, DM 35,—; hard cover US \$ 13.50

Vol. 4 HILTON/STAMMBACH: A Course in Homological Algebra. In preparation

Vol. 5 MAC LANE: Categories. For the Working Mathematician. In preparation

Gaisi Takeuti

Professor Takeuti is Professor of Mathematics at the University of Illinois, Urbana, Illinois. He received his Ph. D. in mathematics from the University of Tokyo in 1956. He was Professor of Mathematics at Tokyo University of Education until 1963 when he joined the mathematics staff of the University of Illinois. Professor Takeuti has an extensive list of papers in logic and set theory published in Japanese and American journals.

Wilson M. Zaring

Dr. Zaring is Associate Professor of Mathematics at the University of Illinois, Urbana, Illinois. He received his Ph. D. in mathematics from the University of Kentucky in 1955 at which time he joined the mathematics staff at the University of Illinois. Since 1964, he has served as Director of the University of Illinois' Academic Year Institute. He is the author of a text in analysis.